



IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 5 Issue: IX Month of publication: September 2017 DOI: http://doi.org/10.22214/ijraset.2017.9198

www.ijraset.com

Call: 🕥 08813907089 🔰 E-mail ID: ijraset@gmail.com



Computational Approach for Finding Pythagoras Octagon Using Programming language MATLAB

S.N.R.G. Bharat Iragavarapu¹, Kota Sai Amulya²

¹Department of Mathematics, ²Department of Electrical and Electronics Engineering, G.V.P. College of Engineering (Autonomous), Visakhapatnam, A.P., India

Abstract: In this paper, using computer programming language MATLAB, for any natural number n, we determine the Pythagoras octagon (a, b, c, d, e, f, g, h) where h denotes the length of the hypotenuse and is \leq n, when one of a, b, c, d, e, f or g is given thereby the number of such Pythagoras octagons are also known.

Keywords: Pythagoras theorem, octagon, Pythagoras octagon, natural numbers, programming language.

I. INTRODUCTION

In [1, 2, 3, 4, 5], we found the remaining side lengths of the Pythagoras triangle, quadrilateral, pentagon, hexagon and heptagon, if one of the side lengths is known. Now in this paper, we exhibit all possible Pythagoras octagons, knowing only one side length that is not hypotenuse. For example, suppose we take the length of one side as 9 and the maximum limit n as 12 then all the possible Pythagoras octagons are (1, 1, 2, 3, 3, 4, 9, 11), (2,3,3, 3,4,4, 9,12), (1,1,2,4,4,5, 9,12), (1,2,2,2,5,5,9,12), (1,2,2,3,3,6,9,12), (1,1,2,2,2,7,9,12). Because the above all combinations satisfies extension of Pythagoras theorem. This process is very difficult if one side length is sufficiently large and n is also large. Now our aim is to find the number of Pythagoras octagons using programming language Matlab.

II. MAIN RESULT

- A. Algorithm
- 1) Step-1: START.
- 2) Step-2: Enter the length of one side of Pythagoras Octagon 'c'.
- 3) Step-3: Read 'c' value.
- 4) Step-4: Enter the maximum limit of hypotenuse of Pythagoras Heptagon 'n'.
- 5) Step-5: Read 'n' value.
- 6) Step-6: Initialise the variables a, b, d, e, f, g, k.
- 7) Step-7: Give k value as 0.
- 8) Step-8: If c>n, go to step9, else go to step 10.
- 9) Step-9: Display that the side length exceeds maximum limit.
- 10) Step-10: Initialise a for loop with condition h=4, h<=n, h increases by 1; If condition fails, go to step 19.
- 11) Step-11: Initialize a for loop with condition a=1, a<=h, increment a by1; If condition fails, go to step 10.
- 12) Step-12: Initialize a for loop with condition b=1, b<=a, increment b by1; If condition fails, go to step 11.
- 13) Step-13: Initialize a for loop with condition d=1, d<=b, increment d by1; If condition fails, go to step 12.
- 14) Step-14: Initialize a for loop with condition e=1, e<=d, increment e by 1; If condition fails, go to step 13.
- 15) Step-15: Initialize a for loop with condition f=1, f<=e, increment f by1; If condition fails, go to step 14.
- 16) Step-16: Initialize a for loop with condition g=1, $g \le f$, increment g by1; If condition fails, go to step 15.
- 17) Step-17:If $a^2+b^2+c^2+d^2+e^2+f^2+g^2=h^2$ go to step 18, else, go to step 16.
- 18) Step-18:Display values of a, b, c, d, e, f, g, h.
- 19) Step-19:Increment count k.
- 20) Step-20: End for-loops.
- 21) Step-21: Display k value.
- 22) Step-22:STOP.



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor:6.887

Volume 5 Issue IX, September 2017- Available at www.ijraset.com

B. Result Analysis

We are required to display all the possible combinations of a Pythagoras octagons by taking one side as parameter which is not a hypotenuse. This can be achieved by the following steps.

- 1) Step-1: Write all the possible combinations that are possible to form a Pythagoras Octagon by keeping a maximum limit to the hypotenuse
- 2) Step-2: Arrange the side lengths in the combinations in ascending order and count the number of combinations and display all the combinations and count.

To illustrate how this works, let us perform this process with one side length 19 and the maximum limit as 21. Step 1: Write all the possible combinations to form Pythagoras Octagon with the given number 18 are.....

a	b	с	d	е	f	g	h
2	2	2	3	3	3	19	20
1	1	1	2	4	4	19	20
1	1	2	2	2	5	19	20
1	2	3	4	5	5	19	21
1	3	3	3	4	6	19	21
2	2	2	4	4	6	19	21
1	1	1	4	5	6	19	21
1	1	2	3	4	7	19	21
1	1	1	2	3	8	19	21
			T 4 1	NEI			

TABLE I

In the above table every combination is a Pythagoras octagon and satisfies the extension of Pythagoras theorem and we can form total 9 Pythagoras octagons if one side length is 19 and maximum limit n as 21.

If the side length is exceed the maximum limit n then result displays no Pythagoras Octagons. For example suppose we take the length of one side as 50 and the maximum limit n as 45 then the result is no Octagons

III.OUTPUTS

>>	>> hexoct										
Enter the length of any one of the sides of Pythogoras Octagon 4											
Enter the maximum limit for the hypotenuse of the Pythogoras Octagon 8											
1	1	1	1	1	2	4	5				
1	1	1	2	2	з	4	6				
1	1	2	3	з	3	4	7				
1	2	2	2	2	4	4	7				
1	1	1	1	2	4	5	7				
1	2	з	з	з	4	4	8				
2	2	2	2	4	4	4	8				
1	1	1	2	4	4	5	8				
The	The total number of Pythogoras Octagons possible										
k =	8										

Fig. 1 One side length=4, Maximum limit=8



International Journal for Research in Applied Science & Engineering Technology (IJRASET)

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor:6.887 Volume 5 Issue IX, September 2017- Available at www.ijraset.com

MATLAB R2015a

HOME	PLOTS	APPS	EDITOR	PUBLISH	ł VI	EW				
w Open Si	ave ✓ Print ✓	🔍 Find 💌	Comment 🤌 Indent 📑		Breakpoints	Run	Run and Advance	Run Section	Run and Time	
	FILE	NAVIGATE		DIT	BREAKPOINTS			RUN		
	C: MATLA	AB MATLAB Pr	roduction Serve	er ▶ R2015a ▶	bin 🕨					
mmand Winde	DW									
>> hexoct										
	e rengen or a	ny one of th	ne sides of	E Pythogora	as Octagon	65				
	e maximum lim:						n 56			
							n 56			
Enter th							n 56			
							n 56			
Enter th							n 56			
Enter th ans =	e maximum lim	it for the h	nypotenuse				n 56			
Enter th ans =		it for the h	nypotenuse				n 56			
Enter th ans =	e maximum lim	it for the h	nypotenuse				n 56			
Enter th ans =	e maximum lim	it for the h	nypotenuse				n 56			
Enter th ans = The maxim	e maximum lim	it for the h	nypotenuse				n 56			
Enter th ans =	e maximum lim	it for the h	nypotenuse				n 56			
Enter th ans = The maxim ans =	e maximum lim um limit exce	it for the h eds the side	ypotenuse : length	of the Pyt			n 56			
Enter th ans = The maxim ans =	e maximum lim	it for the h eds the side	ypotenuse : length	of the Pyt			n 56			
Enter th ans = The maxim ans =	e maximum lim um limit exce	it for the h eds the side	ypotenuse : length	of the Pyt			n 56			
Enter th ans = The maxim ans =	e maximum lim um limit exce	it for the h eds the side	ypotenuse : length	of the Pyt			n 56			
Enter th ans = The maxim ans = The total	e maximum lim um limit exce	it for the h eds the side	ypotenuse : length	of the Pyt			n 56			
Enter th ans = The maxim ans =	e maximum lim um limit exce	it for the h eds the side	ypotenuse : length	of the Pyt			n 56			
Enter th ans = The maxim ans = The total	e maximum lim um limit exce	it for the h eds the side	ypotenuse : length	of the Pyt			n 56			
Enter th ans = The maxim ans = The total	e maximum lim um limit exce	it for the h eds the side	ypotenuse : length	of the Pyt			n 56			

Fig. 2One side length=65, Maximum limit=56

>> hexoct

Enter the length of any one of the sides of Pythogoras Octagon 15

Enter the maximum limit for the hypotenuse of the Pythogoras Octagon 20

1	2	2	2	3	3	15	16	2	3	3	4	7	7	15	19	1	1	3	6	8	8	15	20
1	2	3	3	4	5	15	17	2	3	3	5	5	8	15	19	1	4	4	5	6	9	15	20
1	1	1	3	4	6	15	17	3	3	3	3	6	8	15	19	2	2	5	5	6	9	15	20
1	4	4	4	5	5	15	18	1	1	3	5	6	8	15	19	2	3	3	6	6	9	15	20
2	2	4	5	5	5	15	18	1	2	3	3	7	8	15	19	2	3	4	4	7	9	15	20
2	3	3	4	5	6	15	18	1	2	3	4	5	9	15	19	1	2	2	6	7	9	15	20
1	1	3	4	6	6	15	18	1	1	1	4	6	9	15	19	1	2	3	4	8	9	15	20
1	1	4	4	4	7	15	18	1	1	3	3	4	10	15	19	1	2	2	2	9	9	15	20
1	2	2	4	5	7	15	18	2	5	5	6	6	7	15	20	3	3	4	4	5	10	15	20
2	2	3	3	3	8	15	18	3	3	6	6	6	7	15	20	1	2	3	5	6	10	15	20
1	1	1	4	4	8	15	18	3	4	4	6	7	7	15	20	1	1	1	6	6	10	15	20
1	1	2	2	5	8	15	18	1	2	6	6	7	7	15	20	2	2	3	3	7	10	15	20
3	4	5	5	5	6	15	19	3	4	5	5	6	8	15	20	2	3	3	4	4	11	15	20
4	4	4	4	6	6	15	19	1	3	4	6	7	8	15	20	1	2	2	3	6	11	15	20
1	3	4	5	6	7	15	19	2	3	3	5	8	8	15	20	1	1	2	3	4	12	15	20

The total number of Pythogoras Octagons possible

k=45

Fig. 3 One side length=15, Maximum limit=20



>> hexoct

Enter the length of any one of the sides of Pythogoras Octagon 49

Enter the maximum limit for the hypotenuse of the Pythogoras Octagon 50

1	4	4	4	5	5	49	50	
2	2	4	5	5	5	49	50	
2	3	3	4	5	6	49	50	
1	1	3	4	6	6	49	50	
1	1	4	4	4	7	49	50	
1	2	2	4	5	7	49	50	
2	2	3	3	3	8	49	50	
1	1	1	4	4	8	49	50	
1	1	2	2	5	8	49	50	

The total number of Pythogoras Octagons possible k=9

Fig. 4,5One side length=45, Maximum limit=50

IV.CONCLUSIONS

The process of finding side lengths using manually is very difficult. So by using MATLABfinding the Pythagoras octagon by knowing any one of the side length is becomes novel and easy process. Further we are planning to extend this to Pythagoras n-sided polygon.

REFERENCES

- S.N.R.G.Bharat Iragavarapu, B.saranya, "Extension of Pythagoras Theorem using Programming Language", Journal of Information Technology and Sciences, Volume 3 Issue 1, Pg 1-4, MAT Journals 2017.
- [2] S.N.R.G.Bharat Iragavarapu, Computational approach for finding Pythagoras quadrilateral (a, b, c, d), when a or b or c is given and hypotenuse d ≤ n, where n is a natural number'', International Journal of Advanced Research in management Engineering and Technology, ISSN: 2456 2998 (Online), Volume 2, Issue 3, March 2017, pg: 512-517.
- [3] S.N.R.G.Bharat Iragavarapu, B.saranya, Computational approach for finding Pythagoras pentagon (a, b, c, d, e), when a or b or c or d is given and hypotenuse $e \le n$, where n is a natural number", International Journal of Scientific Research and development, ISSN: 2321 0613 (Online), Volume 5, Issue 5, june, 2017.
- [4] S.N.R.G.Bharat Iragavarapu, k.kavya, Computational approach for finding Pythagoras hexagon (a, b, c, d, e, f), when a or b or c or d or e is given and hypotenuse f ≤ n, where n is a natural number", International Journal for Research in Applied Science & Engineering Technology (IJRASET), ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor:6.887, Volume 5 Issue VII, July 2017, pg: 2112-2116
- [5] S.N.R.G.Bharat Iragavarapu, k.kavya, Computational approach for finding Pythagoras heptagon (a, b, c, d, e, f, g), when a or b or c or d or e is given and hypotenuse $f \le n$, where n is a natural number", International Journal for Research in Applied Science & Engineering Technology (IJRASET), ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor:6.887, Volume 5 Issue VIII, August 2017, pg: 347-351



Author Profile: S.N.R.G.Bharat Iragavarapu (born April 25, 1990) is working as Assistant professor in the Department of Mathematics, GVP College of Engineering (A), Visakhapatnam, Andhra Pradesh, India. He completed his Masters degree in Mathematics from Andhra University in the year 2012 with 8.9 CGPA. (Secured university second). Worked as Junior research fellow in NBHM-DAE project during 2012-2015. Published 3 research papers in international journals. During this academic year 2016-17 there are 10 publications in national journals on the topic related to the present work.



Kota Sai Amulya is studying B.Tech, third year in the Department of Electrical and Electronics Engineering in GVP College of Engineering (A), Visakhapatnam, Andhra Pradesh, India.











45.98



IMPACT FACTOR: 7.129







INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 🕓 (24*7 Support on Whatsapp)