



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 2

Issue: X

Month of publication: October 2014

DOI:

www.ijraset.com

Call: ☎ 08813907089

E-mail ID: ijraset@gmail.com

An investigation study on cancer and its preventions

A. Rethinavelsubramanian¹ K.Suganya²¹Assistant Professor & Head, Department of Mechanical Engineering

K.S.K College of engineering & Technology, Kumbakonam

²Associate Professor, Department of Software Engineering & IT[PG]

A.V.C College of Engineering, Mayiladuthurai

Abstract: Cancers are usually named using -carcinoma, -sarcoma or -blastoma as a suffix, with the Latin or Greek word for the organ or tissue of origin as the root. For example, cancers of the liver parenchyma arising from malignant epithelial cells is called hepatocarcinoma, while a malignancy arising from primitive liver precursor cells is called a hepatoblastoma, and a cancer arising from fat cells is called a liposarcoma. For some common cancers, the English organ name is used. For example, the most common type of breast cancer is called ductal carcinoma of the breast

Keywords – HNPCC, DNA

I. INTRODUCTION

Cancers are a large family of diseases which involve abnormal cell growth with the potential to invade or spread to other parts of the body

Characteristics of Cancer:

- self-sufficiency in growth signalling
- insensitivity to anti-growth signals
- evasion of apoptosis
- enabling of a limitless replicative potential
- induction and sustainment of angiogenesis
- Activation of metastasis and invasion of tissue.

A. Symptoms of Cancer:

When cancer begins, it invariably produces no symptoms. Signs and symptoms only appear as the mass continues to grow or ulcerates. The findings that result depend on the type and location of the cancer. Few symptoms are specific, with many of them also frequently occurring in individuals who have other conditions. Cancer is the new "great imitator". Thus it is not uncommon for people diagnosed with cancer to have been treated for other diseases which were assumed to be causing their symptoms

B. History of Cancer

Cancer has existed for all of human history. The earliest written record regarding cancer is from circa 1600 BC in the Egyptian Edwin Smith Papyrus and describes cancer of the breast. Hippocrates (ca. 460 BC – ca. 370 BC) described several kinds of cancer, referring to them with the Greek word *karkinos* (crab or crayfish). This name comes from the appearance of the cut surface of a solid malignant tumor, with "the veins stretched on all sides as the animal the crab has its feet, whence it derives its name". Galen stated that "cancer of the breast is so called because of the fancied resemblance to a crab given by the lateral prolongations of the tumor and the adjacent distended veins"⁷³⁸ Celsus (ca. 25 BC – 50 AD) translated *karkinos* into the Latin *cancer*, also meaning crab and recommended surgery as treatment. Galen (2nd century AD) disagreed with the use of surgery and recommended purgatives instead. These recommendations largely stood for 1000 years.

In the 15th, 16th and 17th centuries, it became acceptable for doctors to dissect bodies to discover the cause of death. The German professor Wilhelm Fabry believed that breast cancer was caused by a milk clot in a mammary duct. The Dutch professor Francois de la Boe Sylvius, a follower of Descartes, believed that all disease was the outcome of chemical processes, and that acidic lymph fluid was the cause of cancer. His contemporary Nicolaes Tulp believed that cancer was a poison that slowly spreads, and concluded that it was contagious.

The physician John Hill described tobacco snuff as the cause of nose cancer in 1761. This was followed by the report in 1775 by British surgeon Percivall Pott that chimney sweeps' carcinoma, a cancer of the scrotum, was a common disease among chimney

International Journal for Research in Applied Science & Engineering Technology (IJRASET)

sweeps. With the widespread use of the microscope in the 18th century, it was discovered that the 'cancer poison' spread from the primary tumor through the lymph nodes to other sites ("metastasis").


C. Metastasis

Cancer can spread from its original site by local spread, lymphatic spread to regional lymph nodes or by blood (haematogenous spread) to distant sites, known as metastasis. When cancer spreads by haematogenous route, it usually spreads all over body. The symptoms of metastatic cancers depend on the location of the tumor, and can include enlarged lymph nodes (which can be felt or sometimes seen under the skin and are typically hard), enlarged liver or enlarged spleen, which can be felt in the abdomen, pain or fracture of affected bones, and neurological symptoms.

D. Causes

The great majority of cancers, some 90–95% of cases, are due to environmental factors. The remaining 5–10% are due to inherited genetics. Environmental, as used by cancer researchers, means any cause that is not inherited genetically, such as lifestyle, economic and behavioral factors, and not merely pollution. Common environmental factors that contribute to cancer death include tobacco (25–30%), diet and obesity (30–35%), infections (15–20%), radiation (both ionizing and non-ionizing, up to 10%), stress, lack of physical activity, and environmental pollutants.

E. Chemicals



The incidence of lung cancer is highly correlated with smoking.

Exposure to particular substances has been linked to specific types of cancer. These substances are called carcinogens.

Tobacco smoking causes 90% of lung cancer. It also causes cancer in the larynx, head, neck, stomach, bladder, kidney, esophagus and pancreas. Tobacco smoke contains over fifty known carcinogens, including nitrosamines and polycyclic aromatic hydrocarbons. Tobacco is responsible for about one in three of all cancer deaths in the developed world, and about one in five worldwide. Lung cancer death rates in the United States have mirrored smoking patterns, with increases in smoking followed by dramatic increases in lung cancer death rates and, more recently, decreases in smoking rates since the 1950s followed by decreases in lung cancer death rates in men since 1990.

In Western Europe 10% of cancers in males and 3% of all cancers in females are attributed to alcohol exposure, especially cancer of the liver and of the digestive tract.

F. Diet and exercise

Diet, physical inactivity, and obesity are related to up to 30–35% of cancer deaths. In the United States excess body weight is associated with the development of many types of cancer and is a factor in 14–20% of all cancer deaths. Correspondingly, a UK study including data on over 5 million people showed higher body mass index to be related to at least 10 types of cancer, and responsible for around 12,000 cases each year in that country. Physical inactivity is believed to contribute to cancer risk not only through its effect on body weight but also through negative effects on immune system and endocrine system. More than half of the effect from diet is due to overnutrition (eating too much), rather than from eating too few vegetables or other healthful foods.

International Journal for Research in Applied Science & Engineering Technology (IJRASET)

G. Infection

Worldwide approximately 18% of cancer deaths are related to infectious diseases. This proportion varies in different regions of the world from a high of 25% in Africa to less than 10% in the developed world. Viruses are the usual infectious agents that cause cancer but bacteria and parasites may also have an effect.

A virus that can cause cancer is called an oncovirus. These include human papillomavirus (cervical carcinoma), Epstein-Barr virus (B-cell lymphoproliferative disease and nasopharyngeal carcinoma), Kaposi's sarcoma herpesvirus (Kaposi's sarcoma and primary effusion lymphomas), hepatitis B and hepatitis C viruses (hepatocellular carcinoma), and Human T-cell leukemia virus-1 (T-cell leukemias).

H. Radiation

Up to 10% of invasive cancers are related to radiation exposure, including both ionizing radiation and non-ionizing ultraviolet radiation. Additionally, the vast majority of non-invasive cancers are non-melanoma skin cancers caused by non-ionizing ultraviolet radiation, mostly from sunlight. Sources of ionizing radiation include medical imaging and radon gas.

Ionizing radiation is not a particularly strong mutagen. Residential exposure to radon gas. Radiation is a more potent source of cancer when it is combined with other cancer-causing agents, such as radon gas exposure plus smoking tobacco. Radiation can cause cancer in most parts of the body, in all animals, and at any age. Children and adolescents are twice as likely to develop radiation-induced leukemia as adults; radiation exposure before birth has ten times the effect.

Ionizing radiation may be used to treat other cancers, but this may, in some cases, induce a second form of cancer. It is also used in some kinds of medical imaging.

Non-ionizing radio frequency radiation from mobile phones, electric power transmission, and other similar sources has been described as a possible carcinogen by the World Health Organization's International Agency for Research on Cancer.

I. Heredity

The vast majority of cancers are non-hereditary ("sporadic cancers"). Hereditary cancers are primarily caused by an inherited genetic defect. Less than 0.3% of the population are carriers of a genetic mutation which has a large effect on cancer risk and these cause less than 3–10% of all cancer. Some of these syndromes include: certain inherited mutations in the genes *BRCA1* and *BRCA2* with a more than 75% risk of breast cancer and ovarian cancer, and hereditary nonpolyposis colorectal cancer (HNPCC or Lynch syndrome) which is present in about 3% of people with colorectal cancer, among others

J. Hormones

Some hormones play a role in the development of cancer by promoting cell proliferation. Insulin-like growth factors and their binding proteins play a key role in cancer cell proliferation, differentiation and apoptosis, suggesting possible involvement in carcinogenesis.

Hormones are important agents in sex-related cancers such as cancer of the breast, endometrium, prostate, ovary, and testis, and also of thyroid cancer and bone cancer. For example, the daughters of women who have breast cancer have significantly higher levels of estrogen and progesterone than the daughters of women without breast cancer.. Some treatments and prevention approaches leverage this cause by artificially reducing hormone levels, and thus discouraging hormone-sensitive cancers.

K. Pathophysiology

Genetics




International Journal for Research in Applied Science & Engineering Technology (IJRASET)

Cancers are caused by a series of mutations. Each mutation alters the behavior of the cell somewhat

Cancer is fundamentally a disease of tissue growth regulation failure. In order for a normal cell to transform into a cancer cell, the genes which regulate cell growth and differentiation must be altered.

The affected genes are divided into two broad categories. Oncogenes are genes which promote cell growth and reproduction. Tumor suppressor genes are genes which inhibit cell division and survival.

L. Epigenetics



The central role of DNA damage and epigenetic defects in DNA repair genes in carcinogenesis

Epigenetic alterations refer to functionally relevant modifications to the genome that do not involve a change in the nucleotide sequence. Epigenetic alterations occur frequently in cancers. Cancers usually arise from an assemblage of mutations and epimutations that confer a selective advantage leading to clonal expansion

M. Diagnosis



Chest x-ray showing lung cancer in the left lung

International Journal for Research in Applied Science & Engineering Technology (IJRASET)

Most cancers are initially recognized either because of the appearance of signs or symptoms or through screening. Neither of these lead to a definitive diagnosis, which requires the examination of a tissue sample by a pathologist. People with suspected cancer are investigated with medical tests. These commonly include blood tests, X-rays, CT scans and endoscopy

II. TYPES OF CANCER

Cancers are classified by the type of cell that the tumor cells resemble and are therefore presumed to be the origin of the tumor. These types include:

- A. *Carcinoma*: Cancers derived from epithelial cells. This group includes many of the most common cancers, particularly in the aged, and includes nearly all those developing in the breast, prostate, lung, pancreas, and colon.
- B. *Sarcoma*: Cancers arising from connective tissue (i.e. bone, cartilage, fat, nerve), each of which develop from cells originating in mesenchymal cells outside the bone marrow.
- C. *Lymphoma and leukemia*: These two classes of cancer arise from hematopoietic (blood-forming) cells that leave the marrow and tend to mature in the lymph nodes and blood, respectively. Leukemia is the most common type of cancer in children accounting for about 30%.
- D. *Germ cell tumor*: Cancers derived from pluripotent cells, most often presenting in the testicle or the ovary (seminoma and dysgerminoma, respectively).
- E. *Blastoma*: Cancers derived from immature "precursor" cells or embryonic tissue. Blastomas are more common in children than in older adults.

III. PREVENTION FOR CANCER

A. Dietary

While many dietary recommendations have been proposed to reduce the risk of cancer, the evidence to support them is not definitive. The primary dietary factors that increase risk are obesity and alcohol consumption; with a diet low in fruits and vegetables and high in red meat being implicated but not confirmed. Consumption of coffee is associated with a reduced risk of liver cancer.

B. Medication

The concept that medications can be used to prevent cancer is attractive, and evidence supports their use in a few defined circumstances. Daily use of tamoxifen or raloxifene has been demonstrated to reduce the risk of developing breast cancer in high-risk women.

C. Vaccination

Vaccines have been developed that prevent infection by some carcinogenic viruses. Human papillomavirus vaccine (Gardasil and Cervarix) decreases the risk of developing cervical cancer. The hepatitis B vaccine prevents infection with hepatitis B virus and thus decreases the risk of liver cancer.

D. Genetic testing

Gene

BRCA1, BRCA2

HNPCC, MLH1, MSH2, MSH6, PMS1, PMS2

Cancer types

Breast, ovarian, pancreatic

Colon, uterine, small bowel, stomach, urinary tract

Genetic testing for individuals at high-risk of certain cancers is recommended. Carriers of these mutations may then undergo enhanced surveillance, chemoprevention, or preventative surgery to reduce their subsequent risk.

E. Chemotherapy

Chemotherapy is the treatment of cancer with one or more cytotoxic anti-neoplastic drugs (chemotherapeutic agents) as part of a standardized regimen. The efficacy of chemotherapy depends on the type of cancer and the stage. In combination with surgery,

International Journal for Research in Applied Science & Engineering Technology (IJRASET)

chemotherapy has proven useful in a number of different cancer types including: breast cancer, colorectal cancer, pancreatic cancer, osteogenic sarcoma, testicular cancer, ovarian cancer, and certain lung cancers

F. Radiation

Radiation therapy involves the use of ionizing radiation in an attempt to either cure or improve the symptoms of cancer. It works by damaging the DNA of cancerous tissue leading to cellular death. Radiation therapy is used in about half of all cases and the radiation can be from either internal sources in the form of brachytherapy or external sources.

G. Surgery

Surgery is the primary method of treatment of most isolated solid cancers and may play a role in palliation and prolongation of survival. It is typically an important part of making the definitive diagnosis and staging the tumor as biopsies are usually required

H. Palliative care

Palliative care refers to treatment which attempts to make the person feel better and may or may not be combined with an attempt to treat the cancer. Palliative care includes action to reduce the physical, emotional, spiritual, and psycho-social distress experienced by people with cancer.

IV. CONCLUSION

To all that, have love ones that have cancer; lend a helping hand, comfort them, show support, and give love. Ask God to heal and take care of them always know that God would not forsake you, for he is always with you. Give that person hope and confidence that they will get over this and be better. If you believe and support them, that will help keep their spirit up, and the will to live

REFERENCES

- [I] Jump up ^ *World Cancer Report 2014*. World Health Organization. 2014. pp. Chapter 6.7. ISBN 9283204298.
- [II] Jump up ^ "Cancer Glossary". *cancer.org*. American Cancer Society. Retrieved September 11, 2013.
- [III] Jump up ^ "What is cancer?". *cancer.gov*. National Cancer Institute. Retrieved September 11, 2013.
- [IV] ^ Jump up to: Hanahan, Douglas; Weinberg, Robert A. (January 7, 2000). "The hallmarks of cancer". *Cell* **100** (1): 57–70. doi:10.1016/S0092-8674(00)81683-9. PMID 10647931.
- [V] ^ Jump up to: Hanahan, Douglas; Weinberg, Robert A. (2011). "Hallmarks of Cancer: The Next Generation". *Cell* **144** (5): 646–74. doi:10.1016/j.cell.2011.02.013. PMID 21376230.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)