



# **iJRASET**

International Journal For Research in  
Applied Science and Engineering Technology



---

# **INTERNATIONAL JOURNAL FOR RESEARCH**

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

---

**Volume: 5      Issue: XII      Month of publication: December 2017**

**DOI:**

**[www.ijraset.com](http://www.ijraset.com)**

**Call: ☎ 08813907089**

**E-mail ID: [ijraset@gmail.com](mailto:ijraset@gmail.com)**

# Effect of Low Frequency Electromagnetic Field on Glycogen Content in Vital Organs of the Fresh Water Fish, *Rasboradaniconius* (Hamilton, 1822)

S. A. Shaikh<sup>1</sup>, K. R. Reddy<sup>2</sup>

<sup>1,2</sup> Department of Zoology, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad 431004, Maharashtra, India

**Abstract:** Freshwater fish *Rasboradaniconius* was exposed to low frequency electromagnetic field (LF-EMF) for 7 and 15 days to estimate the glycogen content in vital organs. In the present study, it is found that, the significant decline ( $P < 0.05$ ) in glycogen content for 7 days exposure it was in order Brain < Heart < Testis < Ovary < Gills < Intestine < Muscles < Kidney < Liver viz.  $07.5312^* \pm 0.1157 < 12.0820^* \pm 0.0337 < 18.8894^* \pm 0.1207 < 27.5116^* \pm 1.7155 < 32.3926^* \pm 0.5283 < 33.7180^* \pm 0.0667 < 40.7205^* \pm 0.1530 < 41.7205^* \pm 0.0578 < 48.1035^* \pm 0.1204$  mg/100mg and for 15 days exposure it was Brain < Heart < Testis < Ovary < Gills < Intestine < Kidney < Muscles < Liver viz.  $02.9995^* \pm 0.0333 < 6.0656^* \pm 0.1204 < 12.5255^* \pm 0.0883 < 12.6413^* \pm 0.0883 < 15.3217^* \pm 0.0333 < 18.0792^* \pm 0.0334 < 19.1398^* \pm 0.0883 < 20.8079^* \pm 12.013 < 21.6466^* \pm 0.0578$  mg/100mg. Since, the LF-EMF exposure affects glycogen metabolism in various tissues of *Rasboradaniconius*.

**Keywords:** Glycogen, *Rasboradaniconius*, low frequency electromagnetic field.

## I. INTRODUCTION

During the last few decades various behavioural, biochemical and physiological changes have been reported by several workers on the aquatic animals mostly in fishes to low frequency electromagnetic field. There are many underwater current cables passing under the sea and these electrical current induce static magnetic fields around the cables and its effects on many invertebrate and vertebrate species (Bochert and Zettler, 2004). The exposure to static magnetic field could influence the survival rate and fitness of flounder, *Plathichthys flesus* animals of the Baltic Sea. The physiological and biochemical alterations observed in an animal under any physiological stress can be correlated with the structural and functional changes of cellular proteins.

Since many low electromagnetic fields have directly influence on metabolic processes, it is possible that, they may only modify the existing signal transduction procession in cell membrane, thus producing both transduction and biochemical amplification of the effects of the field itself (Luben, 1991). The exposure of the animals to 50 Hz, 0.2 mT magnetic field resulted in the decrease of RBCs membrane elasticity and permeability and changes in the molecular structure of Hb (Aliet. al., 2003). Some studies also suggest that power frequency magnetic field of 0.6mT applied to humans has an effect on cognitive function as has been suggested by animal studies (Preeceet. al., 1998). Some reports show that, an increased risk of breast cancer in females is associated with 50 Hz magnetic field exposure (Forssen, et. al., 2000; Kliukieneet. al., 2004), brain cancer in adult and children (Gurney, and Van Wijngaarden, 1999). In spite of the fact that, the biochemical studies of the fish tissues has drawn the attention of several researchers but the information regarding the detection and estimation of proteins, lipids, carbohydrates and enzymes in the various digestive organs of fishes are limited.

Hence, it is essential to understand the significance alteration or changes in tissues exposure to Low frequency magnetic field on biochemical contents in freshwater fish, *Rasboradaniconius* are less studied, but may contribute to the understanding of the underlying basic mechanisms in the living cells. The study presented in this paper was focused on the biochemical changes induced by low frequency electromagnetic field exposure to freshwater fish, *Rasboradaniconius* for 7 days and 15 days durations.

## II. MATERIAL AND METHODS

### A. Experimental Setup

The freshwater fish, *Rasboradaniconius* were collected from local lake, near to the Aurangabad, Maharashtra and brought them to the laboratory in an oxygenated polythene bags. The healthy adult specimens of *Rasboradaniconius* ranging in length from 10 to 12 cm were selected randomly and acclimatized for the laboratory conditions for experimental purpose. Water was replaced regularly and the fishes were fed daily.

### B. Exposure to LF-EMF

An experimental aquarium was set up with a LF- EMF device made up of concentric coils connected to a solenoid plates. The solenoid plates were wound with 400 turns of 1.6 mm diameter copper wire. The aquarium was filled with 20 L de chlorinated water and the LF- EMF device was immersed in the aquarium. Then it was driven by the 50 Hz power line with a load of 200W bulb so as to establish the low frequency electromagnetic field. Another aquarium is maintained without LF- EMF device for control.



Figure- 1: Freshwater fish, *Rasboradaniconius* (Hamilton, 1822)




Figure- 2: Control and treatment set up of (LF-EMF) exposure to *Rasboradaniconius* under laboratory condition.

### C. Estimation of glycogen:

Wet tissues of the muscle, gills, brain, heart, gut, liver, kidney, testis and ovary of the freshwater fish *Rasboradaniconius* were taken for the estimation of glycogen. Glycogen was estimated by Anthrone reagent method as given by Dezwaan and Zandee (1972). The statistical analysis of the results were done by, One-way Analysis of Variance (ANOVA) using the software package, SPSS (16.0 version) to test the level of significance ( $P < 0.05$ ).



### III. RESULTS AND DISCUSSION

Glycogen is the secondary long term storage form of glucose in animals and energy reserve that can be quickly mobilized to meet a sudden need for glucose. When it is needed for energy, glycogen is broken down and converted again in to glucose. Glycogen phosphorylases is the primary enzyme of glycogen breakdown. For the next 8–12 hours, glucose derived from liver glycogen is the primary source of blood glucose used by the rest of the body as fuel. Liver convert glycogen in to glucose for use throughout the body including the central nervous system that's why glycogen is the essential biomolecule for animal. In present study the comparative data of the glycogen content of both that is control and experimental fishes have been summarized in the table 1.

Sr. No.	Tissues	LF-EMF field exposure period			
		7 days		15 days	
		Control	Experimental	Control	Experimental
01	Brain	16.2087 <sup>*</sup> ± 0.1530	07.5312 <sup>*</sup> ± 0.1157	04.0987 <sup>*</sup> ± 0.0883	02.9995 <sup>*</sup> ± 0.0333
02	Heart	16.9607 <sup>*</sup> ± 0.0578	12.0820 <sup>*</sup> ± 0.0337	12.2363 <sup>*</sup> ± 0.1767	6.0656 <sup>*</sup> ± 0.1204
03	Testis	27.8944 <sup>*</sup> ± 0.1735	18.8894 <sup>*</sup> ± 0.1207	15.3410 <sup>*</sup> ± 0.0578	12.5255 <sup>*</sup> ± 0.0883
04	Ovary	36.1864 <sup>*</sup> ± 0.0334	27.5116 <sup>*</sup> ± 1.7155	27.7595 <sup>*</sup> ± 0.0334	12.6413 <sup>*</sup> ± 0.0883
05	Gills	39.7924 <sup>*</sup> ± 0.1204	32.3926 <sup>*</sup> ± 0.5283	30.7677 <sup>*</sup> ± 0.1204	15.3217 <sup>*</sup> ± 0.0333
06	Gut	45.6353 <sup>*</sup> ± 0.1204	33.7180 <sup>*</sup> ± 0.0667	25.9565 <sup>*</sup> ± 0.2651	18.0792 <sup>*</sup> ± 0.0334
07	Muscles	48.0264 <sup>*</sup> ± 0.1735	40.7205 <sup>*</sup> ± 0.1530	30.5748 <sup>*</sup> ± 0.1669	20.8079 <sup>*</sup> ± 12.013
08	Kidney	51.7288 <sup>*</sup> ± 0.1735	41.7205 <sup>*</sup> ± 0.0578	44.0540 <sup>*</sup> ± 0.1456	19.1398 <sup>*</sup> ± 0.0883
09	Liver	59.6158 <sup>*</sup> ± 0.1859	48.1035 <sup>*</sup> ± 0.1204	55.2632 <sup>*</sup> ± 0.6134	21.6466 <sup>*</sup> ± 0.0578

Values are expressed in mg/100mg of wet tissue weight (Mean ± SD); (\* indicate P<0.05)

Table- 1: Glycogen content in different tissues of *Rasboradaniconius* exposed to low frequency electromagnetic field.

The result indicates that, there is a significant decrease in the glycogen content in all vital organs as the exposure duration increased. The freshwater fish *Rasboradaniconius* exposed to low frequency electromagnetic field showed a significant decreased (P<0.05) level of glycogen in all vital organs like muscle, gills, brain, heart, gut, liver, kidney, testis and ovary tissues when compared control with experimental. Decline in order of glycogen content in tissue for 7 days exposure was Brain< Heart< Testis< Ovary< Gills< Intestine< Muscles< Kidney< Liver viz. 07.5312<sup>\*</sup>±0.1157< 12.0820<sup>\*</sup>±0.0337< 18.8894<sup>\*</sup>±0.1207< 27.5116<sup>\*</sup>±1.7155< 32.3926<sup>\*</sup>±0.5283< 33.7180<sup>\*</sup>±0.0667< 40.7205<sup>\*</sup>±0.1530< 41.7205<sup>\*</sup>±0.0578< 48.1035<sup>\*</sup>±0.1204 mg/100mg and for 15 days exposure it was Brain< Heart< Testis< Ovary< Gills< Intestine< Kidney< Muscles< Liver viz. 02.9995<sup>\*</sup>±0.0333< 6.0656<sup>\*</sup>±0.1204< 12.5255<sup>\*</sup>±0.0883< 12.6413<sup>\*</sup>±0.0883< 15.3217<sup>\*</sup>±0.0333< 18.0792<sup>\*</sup>±0.0334< 19.1398<sup>\*</sup>±0.0883< 20.8079<sup>\*</sup>±12.013< 21.6466<sup>\*</sup>±0.0578 mg/100mg.




Figure-2: Glycogen content in different tissues of *Rasboradaniconius* after 7 days exposure to low frequency electromagnetic field




Figure- 3: Glycogen content in different tissues of *Rasbora daniconius* after 15 days exposure to low frequency electromagnetic field

In the present study it is observed that after exposure of the LF-EMF on *Rasbora daniconius* lead to decline in glycogen content. After exposure of *Rasbora daniconius* to LF-EMF for 7 and 15 days the decline glycogen content is observed in various tissues. After 7 days exposure the decline trend in glycogen content was Brain < Heart < Testis < Ovary < Gills < Intestine < Muscles < Kidney < Liver and for 15 days exposure it was Brain < Heart < Testis < Ovary < Gills < Intestine < Kidney < Muscles < Liver at significant level decreased  $P < 0.05$ . More scientific reports are focused on the stimulation of living organisms, including plants, in maize plants biochemical changes induced by extremely low frequency magnetic field during their early ontogeny stages (first 12 days of growth) Stimulatory effect on the photosynthesis was noticed to the magnetic energy dose increasing, suggesting putative stimulation of photosynthesis process, while the nucleic acid biosynthesis seems to be inhibited however their electromagnetic energy could trigger complex synergetic cellular mechanisms that finally can lead to the growth disturbing (Mihaelaraucuiet. al., 2006). The long term exposure of electromagnetic field on rat the affecting parameters like body weight, blood glucose and fatty acid metabolism. Exposure appears to modify blood glucose content in adult rat 12 and 72 weeks old the rat glycogen amount in hepatocytes cytoplasm was higher and diffusely distributed (Gabriele et. al., 2008). Study of microscopic sections of white rabbit liver, exposed to magnetic field with of 700 MilyGaus intensity and 100 Hz frequency, some cells were seen around lobular central vein having high amount of glycogen near central vein is more and it shows no glycogen discharging thus, the electromagnetic field decreases amount of glycogen in liver (Nafisaisaid, 2010). The electromagnetic field at 50 Hz may modify cell membrane morphology and interfere with initiation of the signal cascade pathway and cellular adhesion (Manniet. al., 2002). The EMF do not have disturbed effects in liver organ, but vaculation of the cytoplasm was reported in this work that because of glycolgenolysis inhibition by EMF (Zareet. al., 2007). Neveen, (2002) has stated that the ultrathin sections of mouse of liver showed highly affected hepatocytes. An opaque cytoplasmic matrix, severe reduction in cytoplasmic organelles, depletion of glycogen, few mitochondria and large electro translucent spaces were observed. Thus, the pattern of reduction in the total amount of glycogen due to exposure of LF EMF can be observed and this corroborates with the earlier findings. Pakhreet. al., (2016) was observed that the significant reduction in glycogen content in different tissues like, gill, liver, muscle and kidney of fresh water fish *Channagachua* in exposure of quinalphos. Gijareet al., (2011) observed the notable alteration in liver and intestine glycogen of *Ophiocephalus punctatus* exposed to sub-lethal concentration of cypermethrin. The glycogen content was analysed from the tissues of experimental and control fish the significant decrease of glycogen content in the gills, kidney and brain and the moderate decline was observed in gonads, muscles whereas slight change found in liver and intestine (Ganeshwadeet al., 2011).

#### IV. CONCLUSION

The present study was concluded that, the LF-EMF significant decline ( $P < 0.05$ ) glycogen content in vital tissues of fresh water fish, *Rasbora daniconius* and it was indicates that, the all vital organs used glycogen as energy source for metabolism; and it was

drastically affected due to exposure to LF-EMF. Among wet tissues, higher glycogen content was observed in liver in control and treated fish and is acceptable because liver is the major storage site of glycogen.

#### V. ACKNOWLEDGEMENT

The authors are thankful to Professor and Head, Department of Zoology, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad (M.S.), India, for providing laboratory facilities during experimentation. We are also thankful to University Grants Commission for financial assessment.

#### REFERENCES

- [1] Ali, F. M., Mohamed W. S and Mohamed, M. R. 2003. Effect of 50 Hz, 0.2 mT magnetic fields on RBC properties and heart functions of albino rats, *Bioelectromagnetics*, 24(8), 535.
- [2] Bocher, R and Zettler M.L. 2004. Long-term exposure of several marine benthic animals to static magnetic fields. *Bioelectromagnetics*, 25:498-502.
- [3] Dezwaan, A. & Zandee, D.I. 1972. Body distribution and seasonal changes in glycogen content of the common sea mussel, *Mytilus edulis*. *Comparative Biochemistry & Physiology* 43(A):53-56.
- [4] Forssen, U. M., Feychting, M., Rutqvist, L. E., Floderus, B. and Ahlbom, A. 2000. Occupational and residential magnetic field exposure and breast cancer in females. *Epidemiology*, 11(1), 24.
- [5] Gabriele, G., Nino, A. D., Prosdocimi, M., Ferrari, V., Barbaro, F., Mazzariol, S., Bernardini, D. and Talpo, G. 2008. Effect of electromagnetic fields of low frequency and low intensity on rat metabolism, *Biomegnetic research and technology*, pp. 6-8.
- [6] Ganeshwade, R.M., Pawar, S.M., Sonawane, S.R., Devararoo, K.D and Sutar, V.B. 2011. Effect of dimethoate on the level of glycogen in tissues of freshwater fish *Puntius ticto* (HAM). *The Bioscan* 6(1): 131-133.
- [7] Gijare, S.S., Raja, I.A., Tantarapale, V.T and Kulkarni, K.M. 2011. Modulation in glycogen of the freshwater fish *Ophiocephalus punctatus* exposed to cypermethrin. *The Bioscan*. 6 (1): 129-130.
- [8] Gurney, J. G. and Van Wijngaarden. E. 1999. Extremely low frequency electromagnetic fields (EMF) and brain cancer in adults and children: Review and comment, *Neuro-oncol*, 1(3), 212.
- [9] Kliukiene, J., Tynes, T. and Andersen, A. 2004. Residential and occupational exposures to 50 Hz magnetic fields and breast cancer in women: a population-based study, *American Journal of Epidemiology*, 159(9), 852.
- [10] Luben, R. A. 1991. Effects of low-energy electromagnetic fields (pulsed and DC) on membrane signal transduction processes in biological systems, *Health-Phys.*, 61, 15.
- [11] Manni V, Lisi A, Pozzi D, Rieti S, Serafino A, Giuliani L. and Grimaldi, S. 2002. Effects of extremely low frequency (50 Hz) magnetic field on morphological and biochemical properties of human keratinocytes. *Bioelectromagnetics*, May; 23(4): 298-305.
- [12] Neveen Hussein Mahmoud. 2002. Ultrastructure study on the effect of alternating magnetic field on mice liver. *The Egyptian journal of hospital medicine* vol., 19: 168-178.
- [13] Pakhre, N.B., Gaikwad, S. B. and Reddy, K. R. 2016. Acute toxicity of quinalphos (25% EC) on glycogen content in different tissues of the freshwater fish, *Channagachua* (Hamilton, 1822). *The Ecoscan*. Vol. ix: 783-787.
- [14] Preece, A. W., Wesnes, K. A and Iwi, G. R. 1998. The effect of a 50 Hz magnetic field on cognitive function in humans. *Int J Radiat Biol.* Oct; 74 (4): 463-70.
- [15] Racuciu, M. D. and Amaraitei, C. 2006. Biochemical changes induced by low frequency magnetic field exposure of vegetal organisms. *Rom. Journ. Phys.*, Vol. 52, Nos. 5-7, p. 645-651, Bucharest.
- [16] Saeid, N., Yusof, B., Rahim, H., Shamsadin, A. S., Fateme, H., Fatere Y. and Leili, M. 2010. Influence of electromagnetic fields of two phases square wave with low frequency on serum alt and ast levels and histochemistry of hepatocytes glycogen. *Global Veterinaria* 5 (4): 204-208.
- [17] Zare, S., Alivandi, S. and Ebadi, A. G. 2007. Histological studies of the low frequency electromagnetic fields effect on liver, testes and kidney in guinea pig. *World Applied Sciences Journal* 2 (5): 509-511.





10.22214/IJRASET



45.98



IMPACT FACTOR:  
7.129



IMPACT FACTOR:  
7.429



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24\*7 Support on Whatsapp)