



IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 6 Issue: III Month of publication: March 2018 DOI: http://doi.org/10.22214/ijraset.2018.3113

www.ijraset.com

Call: 🕥 08813907089 🔰 E-mail ID: ijraset@gmail.com



Self-Weak Complementary Fuzzy Soft Graph

Dr. N. Sarala¹ R. Deepa²

¹Department of Mathematics, A. D.M College for Women Nagapattinam, Tamilnadu,India ²Department of Mathematics, E.G. S. Pillay Engineering College Nagapattinam, Tamilnadu,India

Abstract: In this paper, we discussed the notion of fuzzy soft graph and concepts of homomorphism, isomorphism, weak isomorphism of fuzzy soft graphs. Also some properties of isomorphism on fuzzy soft graphs, self-complementary and self-weak complementary fuzzy soft graphs are discussed.

Keywords: Weak isomorphism, self-complementary fuzzy soft graphs, self-weak complementary fuzzy soft graphs.

I. INTRODUCTION

The concept of fuzzy set theory was introduced by zadeh. A [1]. This concept have potential application in various fields such as the smoothness of functions, medical and life sciences, social sciences, Engineering graph theory, artificial intelligence, robotics, computer, networks, decision making and automata theory. The concept of fuzzy soft sets with is a new mathematical tool was firstly introduced by Maji et al [2]. They presented the definition of fuzzy soft sets and investigated some properties at this notion. Therefore many researchers have applied this concept on different branches. In 1975, Rosenfeld [8] introduced the concept of fuzzy sets. P. Bhattacharya in [3] showed that a fuzzy graph can be associated with a fuzzy group in a natural way as an automorphism group. K.R. Bhutani in [4] introduced the concept of weak isomorphism and isomorphism between fuzzy graphs. In this paper we define the concept of homomorphism, isomorphism, weak isomorphism, self weak isomorphism of fuzzy soft graph. We also study some of their important properties.

II. PRELIMINARIES

In this section ,We recall some basic notion of graph ,fuzzy graph, soft graph and fuzzy soft graphs

A. Definition:2.1

A graph G = (V, E) consists of a non-empty set of objects V called vertices and a set E of two element subset of V called edges.

B. Definition:2.2[5]

Let V be a non-empty finite set $\mu: V \to [0, 1]$ and $v = VxV \to [0, 1]$. If $v(x, y) \le \mu(x) \land \mu(y)$ for all $x, y \in V$. Then the pair $G = (\mu, v)$ is called a *fuzzy graph* over the set V. Here μ and v are called *fuzzy vertex* and *fuzzy edge* of the fuzzy graph (μ, v) .

C. Definition:2.3[6]

Let (F, A) be a soft set over V. Then (F, A) is said to be a *soft graph* of G if the subgraph induced by F(x) in G. F(x) is a connected subgraph of G for all $x \in A$.

D. Definition:2.4

A fuzzy soft graph $\tilde{G} = (G^*, \tilde{F}, \tilde{K}, A)$ is a 4 tuple such that

(i) $G^* = (V, E)$ is a simple graph

(ii) A is a nonempty set of parameters

(iii) (\tilde{F}, A) is a fuzzy soft set over V.

(iv) (\widetilde{K}, A) is a fuzzy soft set over E.

(v) $(\tilde{F}(a), \tilde{K}(a))$ is a fuzzy (sub) graph of G*, for all $a \in A$.



That is \widetilde{K} (a) (xy) $\leq \min \{\widetilde{F}(a)(x), \widetilde{F}(a)(y)\}\$ for all $a \in A$, x, $y \in V$. The fuzzy graph ($\widetilde{F}(a), \widetilde{K}(a)$) is denoted by \widetilde{H} (a) for convenience.

On the other hand, a fuzzy soft graph is a parameterized family of fuzzy graphs. The class of all fuzzy soft graphs of G* is denoted by F(G*).

E. Definition:2.5

Let $\widetilde{G_1} = (G^*, \widetilde{F_1}, \widetilde{K_1}, A)$ and $\widetilde{G_2} = (G^*, \widetilde{F_2}, \widetilde{K_2}, A)$ be two fuzzy soft graphs. A homomorphism $f: \widetilde{G_1} \to \widetilde{G_2}$ is a mapping $f: V_1 \to V_2$ which satisfies the following conditions.

- (i)
- $$\begin{split} \widetilde{F_1} & (a) \ (x) \leq \widetilde{F_2} \ (a)(f(x)) \\ \widetilde{K_1} & (a)(xy) \leq \widetilde{K_2} \ (a) \ (f(x)f(y)) \quad \text{for all } a \in A, \quad x, y \in V_1, \ x \ y \in E. \end{split}$$
 (ii)
- F. Definition:2.6

Let $\widetilde{G_1} = (G^*, \widetilde{F_1}, \widetilde{K_1}, A)$ and $\widetilde{G_2} = (G^*, \widetilde{F_2}, \widetilde{K_2}, A)$ be two fuzzy soft graphs. An *isomorphism*

 $f: \widetilde{G_1} \to \widetilde{G_2}$ is a Bijective mapping $f: V_1 \to V_2$ which satisfies the following conditions.

- $F_1(a)(x) = F_2(a(f(x)))$ (i)
- $K_1(a)(xy) = K_2(a)(f(x)f(y))$ for all $a \in A$, $x, y \in V_1$, $xy \in E$ (ii)
- G. Definition:2.7

Let $\widetilde{G_1} = (G^*, \widetilde{F_1}, \widetilde{K_1}, A)$ and $\widetilde{G_2} = (G^*, \widetilde{F_2}, \widetilde{K_2}, A)$ be two fuzzy soft graphs. Then a *weak isomorphism*

 $f: \widetilde{G_1} \to \widetilde{G_2}$ is a bijective mapping $f: V_1 \to V_2$ which satisfies the following conditions.

- (i) f is homomorphism $F_1(a)(x) = F_2(a)(f(x))$ for all $a \in A$, $x \in V_1$ (ii)
- H. Definition:2.8

Let $\widetilde{G_1} = (G^*, \widetilde{F_1}, \widetilde{K_1}, A)$ and $\widetilde{G_2} = (G^*, \widetilde{F_2}, \widetilde{K_2}, A)$ be two fuzzy soft graphs. A Self Weak Isomorphism

 $f: \widetilde{G_1} \to \widetilde{G_2}$ is a bijective mapping $f: V_1 \to V_2$ which satisfies the following conditions.

- f is homomorphism (i)
- $\widetilde{K_1}$ (a)(xy) = $\widetilde{K_2}$ (a) (f(x)f(y)) for all a $\in A$, x, y $\in V_1$, xy $\in E$. (ii)

III. SELF COMPLEMENTARY FUZZY SOFT GRAPHS

A. Definition:3.1

A fuzzy soft graph G is self complementary if $G \cong G'$ where G' is the complement of fuzzy soft graph G.

B. Theorem :3.1

If $\tilde{G} = (G^*, \tilde{F}, \tilde{K}, A)$ be a self complementary fuzzy soft graph then

$$\sum_{x=y}^{\Sigma} \widetilde{K}(a) (xy) \le \frac{1}{2} \sum_{x=y}^{\Sigma} \min[(\widetilde{F}(a) (x), \widetilde{F}(a) (y))]$$

Proof:



International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 6.887 Volume 6 Issue III, March 2018- Available at www.ijraset.com

Let $\tilde{G} = (G^*, \tilde{F}, \tilde{K}, A)$ be a self complementary fuzzy soft graph. Where $\tilde{G_1}$ is a finite set. Then there exists an isomorphism $f: \tilde{G_1} \to \tilde{G_1}$ is a mapping $f: V_1 \to V_1$

$$\widetilde{F}_{1}^{C}(a)(x) = \widetilde{F}_{1}(a)(x) = \widetilde{F}_{1}(a)(f(x))$$
 ------(3.1.1)

 $\widetilde{K_1}^{C}(a)(xy) = \widetilde{K_1}(a)(xy) = \widetilde{K_1}(a)(f(x) f(y))$ ------(3.1.2)

By definition of complement of fuzzy soft graph,

$$\widetilde{K_{1}}^{C}(\mathbf{a})\,(\mathbf{x}\mathbf{y}) \ = \widetilde{F_{1}}^{C}(\mathbf{a})\,(\mathbf{x}) \wedge \widetilde{F_{1}}^{c}(\mathbf{a})\,(\mathbf{y}) - \widetilde{K_{1}}^{c}(\mathbf{a})\,(\mathbf{f}(\mathbf{x})\,\mathbf{f}(\mathbf{y}))$$

By using (3.1.1) and (3.1.2) we get,

$$\begin{split} &\widetilde{K_1} (a) (xy) = \widetilde{F_1} (a) (x) \wedge \widetilde{F_1} (a) (y) - \widetilde{K_1} (a) (f(x) f(y)) \\ &\widetilde{K_1} (a) (xy) + \widetilde{K_1} (a) (f(x) f(y)) = \widetilde{F_1} (a) (x) \wedge \widetilde{F_1} (a) (y) \end{split}$$

again using (3.1.1) and (3.1.2) we get,

$$\widetilde{K_1} (a) (xy) + \widetilde{K_1} (a) (xy) = \widetilde{F_1} (a) (x) \wedge \widetilde{F_1} (a) (y)$$

$$\Rightarrow 2\widetilde{K_1} (a) (xy) = \widetilde{F_1} (a) (x) \wedge \widetilde{F_1} (a) (y)$$

$$\Rightarrow \widetilde{K_1} (a) (xy) = \frac{1}{2} \widetilde{F_1} (a) (x) \wedge \widetilde{F_1} (a) (y)$$

similarly, we have

$$\sum \widetilde{K_1} (a) (xy) = \frac{1}{2} \sum_{x=y}^{\Sigma} \min[\widetilde{F_1}(a) (x), \widetilde{F_1} (a) (y)]$$

In general,

 $\sum_{x=y}^{\Sigma} \widetilde{K}(a) (xy) \leq \frac{1}{2} \sum_{x=y}^{\Sigma} \min[(\widetilde{F}(a) (x), \widetilde{F}(a) (y))].$ Hence \widetilde{G} be a self complementary fuzzy soft graph.

C. Definition: 3.2

A fuzzy soft graph \tilde{G} is self weak complementary if \tilde{G} is weak isomorphic with \tilde{G}^{C} , where \tilde{G}^{C} is the complement of fuzzy soft graph \tilde{G} .

 $\widetilde{F}_{1} (a) (x) = \widetilde{F}_{1} (a) (f(x))$ (3.2.1) $\widetilde{K}_{1} (a) (xy) = \widetilde{K}_{1}^{C} (a) (f(x) f(y))$ (3.2.2)

D. Theorem 3.2.1 Let $\tilde{G} = (G^*, \tilde{F}, \tilde{K}, A)$ be a self weak complementary fuzzy soft graph then

$$\sum_{x=y}^{\Sigma} \widetilde{K_{1}}(a) (xy) \leq \frac{1}{2} \sum_{x=y} [\widetilde{F_{1}}(a) (x) \wedge \widetilde{F_{1}}(a) (y)]$$

1) Proof: Let $\tilde{G} = (G^*, \tilde{F}, \tilde{K}, A)$ be a self weak complementary fuzzy soft graph. Then there exists a weak isomorphism $f: V_1 \rightarrow V_1$ such that f(x) = x, and $\tilde{K}_1(a)(xy) \leq \tilde{K}_1^C(a)(f(x) f(y))$

By definition of complement of fuzzy soft graph

 $\widetilde{K_{1}}^{C}(a) (f(x), f(y)) = \widetilde{F_{1}}(a) (f(x)) \wedge \widetilde{F_{1}}(a) (f(y)) - \widetilde{K_{1}}(a) (f(x), f(y)) - \cdots - (3.2.3)$



By using (3.2.2) in (3.2.3), we have,

$$\begin{split} &\widetilde{K_1}(a) \ (xy) \leq \widetilde{F_1}(a)(f(x)) \ \land \widetilde{F_1}(a) \ (f(y)) \cdot \widetilde{K_1}(a) \ (f(x) \ f(y)) \\ & \text{By using (3.2.1) in (3.2.3) we have,} \\ & \widetilde{K_1}(a) \ (xy) \leq \widetilde{F_1}(a)(x) \ \land \widetilde{F_1}(a) \ (y) \ - \widetilde{K_1}(a) \ [f(x) \ f(y)] \\ & \widetilde{K_1}(a) \ (xy) + \widetilde{K_1}(a)(f(x) \ f(y) \leq \widetilde{F_1}(a) \ (x) \ \land \widetilde{F_1}(a) \ (y) \\ & \text{Similarly, we have} \\ & x = \frac{v}{y} \widetilde{K_1}(a) \ (xy) + x = \frac{v}{y} \widetilde{K_1}(a) \ (f(x) \ f(y)) \leq \sum_{x = y} [\widetilde{F_1}(a) \ (x) \ \land \widetilde{F_1}(a \ (y)] \\ & \text{By using (3.2.2) in (3.2.4), we have,} \\ & x = \frac{v}{y} \widetilde{K_1}(a) \ (xy) + x = \frac{v}{y} \widetilde{K_1}(a) \ (xy) \leq \sum_{x = y} [\widetilde{F_1}(a)(x) \ \land \widetilde{F_1}(a) \ (y)] \\ & 2 \ x = \frac{v}{y} \widetilde{K_1}(a) \ (xy) \leq x = \frac{v}{y} [\widetilde{F_1}(a) \ (x) \ \land \widetilde{F_1}(a) \ (y)] \\ & x = \frac{v}{y} \widetilde{K_1}(a) \ (xy) \leq \frac{1}{2} \ x = \frac{v}{y} [\widetilde{F_1}(a) \ (x), \ \land \widetilde{F_1}(a) \ (y)] \\ & \text{Hence } \widetilde{G} \ \text{is a self weak complementary fuzzy soft graph.} \\ & \text{by equation (3.2.5))} \\ & \widetilde{K_1}^{\widetilde{c}}(a) \ (f(x), f(y)) \qquad \geq \frac{1}{2} \ \Sigma \ (F_1(a) \ (x) \ \widetilde{F_1}(a) \ (y)] \end{aligned}$$

 $\geq \sum_{x=y}^{\Sigma} \widetilde{K_1} (a) (xy) \qquad \forall xy \in \text{ Identity map.}$ i.e., $\sum_{x=y}^{\Sigma} \widetilde{K_1} (a) (xy) \qquad \leq \widetilde{K_1} (a) (f(x), f(y)) \qquad -----(3.2.8)$

From (3.2.6) and (3.2.8) \tilde{G} is weak isomorphic with \tilde{G}^{c} therefore G is a self weak complementary fuzzy soft graph.

IV. CONCLUSION

Fuzzy soft graph theory is an extremely useful tool in solving the combinational problems in different areas. In this paper we gave some new concepts such as weak isomorphism, self weak complementary fuzzy soft graph and studied some at their properties.

REFERENCES

- A. Journal Papers
- [1]Zadeh,A. Information and control (1965)
- [2] Maji .P.K, Roy A.R.and Biswas.R Journal at Fuzzy mathematics (2001).
- [3] Bhattacharya.P.Some Remarks on fuzzy graphs,Pattern Recognition Letter 6:297-302,(1987).
- [4] Bhutani.K.R. On Automorphism of Fuzzy graphs, Pattern Recognition Letter 9: 159-162, (1989).
- [5] SunithaM.S,andVijayakumar.A, Complement of fuzzy graph.IndianJ.,Pure appl,math,33(9),1451-1464 september 2002
- [6] Akram. M , Information sciences. 181,5548-5564 (2011).
- [7] Akram. M and Dudek W.A, Computers Mathematics. Appl. 61, 289-299 (2011).

B. Books

[1] Rosenfeld, A, Fuzzy sets and their applications (Academic Press, New York, 1975)











45.98



IMPACT FACTOR: 7.129







INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 🕓 (24*7 Support on Whatsapp)