



# **iJRASET**

International Journal For Research in  
Applied Science and Engineering Technology



---

# **INTERNATIONAL JOURNAL FOR RESEARCH**

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

---

**Volume: 6      Issue: IV      Month of publication: April 2018**

**DOI: <http://doi.org/10.22214/ijraset.2018.4711>**

**[www.ijraset.com](http://www.ijraset.com)**

**Call: ☎ 08813907089**

**E-mail ID: [ijraset@gmail.com](mailto:ijraset@gmail.com)**

# A Model Study of Lower Ionospheric Composition during Solar Flares: I- Negative Ions and Electron Concentration

Yogendra Kumar Verma<sup>1</sup>, Sanjeev Srivastava<sup>2</sup>

<sup>1</sup>Department of Physics, Guru Hari Kishan Degree College Gola-Kheri,

<sup>2</sup>Principal, Govt. Degree college, Hempur, Bisalpur(Pilibhit)

**Abstract:** An ion chemical model study of seasonal and latitudinal variation of the composition of lower terrestrial ionosphere under M, C and X class x-ray flares is made under varying solar and geomagnetic activity. Computed concentrations are seen to vary from  $10^2$  to  $10^4$  for dominant ions for class C to class X flare for all latitudes. The seasonal variation of flare time concentrations is however not very well defined. Effectiveness of a flare is seen to depend on its time of occurrence.

**Keywords:** Lower Terrestrial Ionosphere, Solar Flares, Geomagnetic Activity, Negative Ions, Electron Concentration

## I. INTRODUCTION

The lower part of the ionosphere, which comprises of the D-region and lower E-region covering the altitude range from nearly 60 km to 110 km has practical importance because, it influences radio propagation to distant places on the Earth K. Rawer [1]. Solar flares affect this region due to excess ionization caused by the enhanced solar x-rays flux, below  $10\text{\AA}$ . This increase in x-ray radiation could be of several orders of magnitude for very strong flares. The excess ionization is responsible for enhanced radio wave absorption.

The solar x-ray flares which occurred on 19.01.2005, 04.11.2004 and 23.07.2002 and belonging to Class C7.7, M5.4 and X4.8 respectively have been adopted as representative flares for C, M and X classes, are used to study seasonal and latitudinal variation of negative ions and electron concentrations under Solar Maximum ( $F_{10.7} > 250$ ) and Solar Minimum ( $F_{10.7} < 70$ ) conditions for geomagnetically quiet ( $A_p < 20$ ) and disturbed days ( $A_p > 50$ ).

In order to do that, for taking the inputs from various models, we have chosen the dates 24.2.2008, 18.8.2008 and 25.10.2007 as representative of winter, summer and equinox respectively under solar minimum and geomagnetically quiet conditions; 15.12.2006, 15.5.2005 and 11.9.2005 as representative of winter, summer and equinox respectively under solar minimum and geomagnetically disturbed conditions; 26.12.2001, 19.7.2000 and 24.9.2001 as representative of winter, summer and equinox respectively under solar maximum and geomagnetically quiet conditions; 06.11.2001, 15.7.2000 and 31.3.2001 as representative of winter, summer and equinox respectively under solar maximum and geomagnetically disturbed conditions as these days were found to have the appropriate levels of solar and geomagnetic activity.

## II. ION CHEMICAL MODEL

### A. Reaction schemes

In order to obtain the densities of major negative ions and electrons, in this work we have employed detailed ion-chemical model is shown in fig 1, which present the negative ion flow chart. This scheme takes into 36 negative and electron is essentially similar to the reaction scheme given by J. Kazil [2]. Details of the reactions and their kinetic rates are given elsewhere Y.K.Verma [3].

### B. Neutral atmosphere

Neutral atmospheric parameters namely the number densities of He, O, N<sub>2</sub>, O<sub>2</sub>, Ar, H, and N, total mass density; neutral temperature and exospheric temperature, are adopted from the MSISE model A.E. Hedin[4],[5], which describes the neutral temperature and densities in the Earth's atmosphere from ground to thermospheric heights.

### C. Ionizing fluxes




Fig.1 Negative ion flow chat for 37 ion scheme

The significant contributors to the lower ionospheric ionization are solar Lyman-alpha ( $1216\text{\AA}$ ), solar x-rays in the range  $1-10\text{\AA}$  and galactic cosmic rays below  $65\text{ km}$ . During a flare event, the Lyman-alpha increases by relatively small amount compared to the huge normal solar Lyman-alpha emission. By contrast, observations of solar x-ray from GOES satellites show huge increase during flares. The solar flux above the atmosphere  $I_{\infty}(\lambda)$  is estimated by the SOLAR 2000 model, which provides various daily 1 AU adjusted (or observed) solar irradiance products and historical data W.L.Tobiska et al.[6]. This spectral range of solar radiation is responsible for the main photo ionisation and photo dissociation processes in the upper stratosphere, mesosphere, and lower thermosphere Brasseur and Solomon [7]. A spectrum of the power law form-

$$I(\lambda) = A \lambda^B \dots\dots\dots(1)$$

has been found to be adequate by many workers Rowe et al.[8]; P. A. J. Ratnasiri and C. F.Sechrist[9]; J. W. Parker and S. A. Bowhill[10] and in present work, the same form has been employed. Where  $I$  is the flux of wavelength in  $\text{Wm}^{-2}$ ,  $\lambda$  the wavelength in  $\text{\AA}$  and  $A$  and  $B$  are constants depending on GOES x-ray detector chamber responses for  $0.5-4\text{\AA}$  and  $1-8\text{\AA}$ . typical spectrum is shown in figs 2-4

### III. COMPUTATIONS

#### A. Production Rates

The ion production rates have been computed using the well known Chapman production function Rishbeth and Garriot, [11] using the fluxes described in the preceding sections and the ionization and absorption cross sections compiled from various sources. The photoionization of metastable  $\text{O}_2(^1\Delta_g)$  by ultra violet radiations in the range  $1027-1118\text{\AA}$  is computed from the expression given by Paulsion et al. [12] and the cosmic rays ionization has been found by the expression given by Brasseur and soloman, 2005[7]

$$q(j, \lambda, h) = F(\lambda, \infty) I(j, \lambda) n(j, h) \exp \left[ - \sum_j A(j, \lambda) \int_h^\infty n(j, z) dz \right] \dots\dots\dots(2)$$




Fig. 2 - Typical X4.8 Class flare





Fig. 3 Typical C7.7 Class Flare

Fig. 4 - Typical M5.4 Class flare

### B. Ion Composition

The reaction schemes are used to construct the rate equations for computing the densities of various ions. The some typical computed height profiles of negative ions under C, M and X class flare and under Solar Maximum ( $F_{10.7} > 250$ ) and Solar Minimum ( $F_{10.7} < 70$ ) conditions for geomagnetically quiet ( $A_p < 20$ ) and disturbed days ( $A_p > 50$ ).





#### IV. RESULTS AND DISCUSSION

##### A. Production Rates

computed results are obtained for the effect on the ion production rate for latitude  $0^{\circ}\text{N}$  to  $90^{\circ}\text{N}$  and  $0^{\circ}\text{S}$  to  $90^{\circ}\text{S}$  for solar minimum and solar maximum conditions for geomagnetically quiet and disturbed days. Fig 6-11 show typical plots of ion production rate at 70 Km. Under varying solar and geomagnetic activity levels, for all latitudes, the galactic cosmic ray contribution is less compared to the ionization due to Lyman alpha which in turn is less than the contribution of metastable oxygen. The total non flare value of ion production is found to be very close to the total ion production rate for C-class flare typically of the order of  $10^6 \text{ cm}^{-3}$ . However, this value is surpassed for M-class flare ( $10^7 \text{ cm}^{-3}$ ) and X-class flare ( $10^8 \text{ cm}^{-3}$ ), with hard X-rays under flare conditions contributing significantly to the ion production. Yet another feature quite discernible from our model computations is that  $(\text{N}_2^+ + \text{O}_2^+)$  usually much below the  $\text{NO}^+$ , quickly overtakes the production rate of  $\text{NO}^+$  under flare conditions. These results are in conformity with earlier studies carried out for specific flares based on experimental observations A.P. Mitra[13], Srivastava et al.[14]; Verronen et al.[15]. The flare induced ionization is found to vary with solar zenith angle, i.e. a solar flare of same class occurring at different times of the day induces varying degree of ionization at different latitudes. During flare, the ion production rates are found to reach a maximum between 85 km and 100 Km and show a decrease thereafter, a feature which can be very well understood in the light of the effectiveness of hard x-rays in producing ionization at lower altitudes.




Fig. 5- Computed ion production Rates for winter

##### B. Negative Ion and Electron Concentrations

The model results obtained for all latitudes under varying level of solar and geomagnetic activity show the concentrations to vary from  $10^2$  to  $10^4$  for the dominant negative ions for C class flare to X class flare. Similar effects have been obtained by other workers. The seasonal variation for flare time concentrations is however again seen to be not very well defined. Effectiveness of a flare is seen to depend on the time of occurrence of the flare. The hydrated cluster ions are seen to be present only up to a height of about 70 Km and their concentrations fall off very rapidly above this height. This trend is observed for all flare types. The radio wave absorption has also been computed for a radio wave of 2.2 MHz using Sen-Wyller magnetoionic theory in conjunction with appropriate collision frequency profiles and electron density obtained from our model. Typical values of 42 dB for the C-class flare, 63dB for M- class flare and 96 dB for X-class flare are obtained.

#### V. ACKNOWLEDGEMENTS

The authors are thankful to Dr. Raj Kumar Tyagi for help in plotting graphs And Dr. N K Batra for completing this work.

## REFERENCES

- [1] K Rawer, "Wave Propagation in the Ionosphere", Kluwer Acad. Publ. Dordrecht., 1993
- [2] J Kazil, "The University of Bern Atmospheric Ion model: Time dependent ion modelling in the stratosphere", mesosphere and lower thermosphere", Ph.D. thesis, University of Bern, 2002
- [3] Y K Verma, "A Study of Solar Flare Effects and Related Phenomena in Lower Terrestrial Ionosphere", Ph.D. Thesis, MJP Rohilkhand University, Bareilly(U.P.), 2014
- [4] A E Hedin, "Extension of the MSIS thermosphere model to the middle and lower atmosphere", Journal of Geophysical Research, 96, 1159-1172, 1991
- [5] A E Hedin, "MSIS-86 Thermospheric Mode" I, J. Geophys. Res., 92, 4649, 1987
- [6] W K Tobiska, T Woods, F Eparvier, R Viereck, L Floyd, D Bouwer, G Rottman, O R White, "The SOLAR2000 empirical solar irradiance model and forecast tool", JASTP, 62, 1233-1250, 2000
- [7] G Brasseur and S Solomon, "Aeronomy of middle atmosphere" Springer, Dordrecht, 3<sup>rd</sup> revised and enlarged edn., 2005
- [8] J N Rowe, A J Ferraro, H S Lee, R W Kreplin and A P Mitra, "Observations of electron density during a solar flare" J Atmos. Terr. Phys., 32, 1609, 1970
- [9] P A J Ratnasiri and C F Sechrist, "An investigation of the solar zenith angle variation of D region ionization", Aeronomy Rep. No. 67, Elec. Eng. Dep, University of Illinois, Urbana-Champaign, 1975
- [10] J W Parker and S A Bowhill, "Solar flare ionization in the mesosphere observed by Coherent Scatter Radar", Aeronomy Rep No.118, Elec. Eng. Dep. University of Illinois, Urbana-Champaign, 1986
- [11] H Rishbeth and O K Garriot, "Introduction to ionospheric Physics", Academic Press Inc, New York, 1969
- [12] D E Paulson, R E Huffman and J C Larrabee, "Radio Science" USA, 7, 51, 1972
- [13] A P Mitra and J N Rowe, "Ionospheric effects of solar flares-VI: Changes in D region ion chemistry during solar flares", J. Atmos. Terr. Phys., 34, 795, 1972
- [14] S Srivastava, N Nath, C S G K Setty, "A model study of lower ionosphere during solar flare events and ionospheric absorption at Delhi", Indian J. Radio and Space Physics, 23, 5, 347-358, 1994
- [15] P T Verronen, E Turunen, T Ulich and E Kyrola, "Modelling the effect of the October 1989 solar proton event on mesospheric odd nitrogen using a detailed and neutral chemistry model", Ann. Geophysicae, 1967-1976, 2002




Fig. 6 -Computed Negative ions and Electrons height profiles for winter in Solar Minimum and Geomagnetically disturbed




Fig. 7 -Computed Negative ions and Electrons height profiles for winter in Solar Maximum and Geomagnetically disturbed




Fig. 8 -Computed Negative ions and Electrons height profiles for Equinox in Solar Minimum and Geomagnetically disturbed




Fig. 9 -Computed Negative ions and Electrons height profiles for Equinox in Solar Minimum and Geomagnetically disturbed




Fig. 10-Computed Negative ions and Electrons height profiles for winter in Solar Maximum and Geomagnetically disturbed







Fig. 11 -Computed Negative ions and Electrons height profiles for winter in Solar Maximum and Geomagnetically disturbed



10.22214/IJRASET



45.98



IMPACT FACTOR:  
7.129



IMPACT FACTOR:  
7.429



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24\*7 Support on Whatsapp)