



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 7 Issue: V Month of publication: May 2019

DOI: <https://doi.org/10.22214/ijraset.2019.5466>

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

Equivalence of Two Summability Methods for Improper Integral

Dr. Banitamani Mallik, Mr Radhamadhab Dash, Dr. Umakanta Misra

¹Centurion University of Technology and Management, Paralakhemundi, Odisha, India

²Delhi Public School, Nalco Nagar, Angul, Odisha, India

³National Institute of Science and Technology, Ganjam, Odisha, India

Abstract: In this paper, we introduce the concept of $|N, p; \delta, \mu|_k, k \geq 1$ summability of improper integrals and by this definition we prove a theorem that generalizes theorems of Orhan, Ozgen and Acharya.

2010 Mathematics Subject Classification: 26D15; 40F05; 40D25; 40A10; 35A23

Keywords: Cesaro and Riesz method, summability of improper integrals

I. INTRODUCTION

Let f be a real valued function which is continuous on $[0, \infty)$. Let $s(x) = \int_0^x f(t) dt$. We denote the Cesaro mean of $s(x)$ by

$\sigma(x)$. The improper integral $\int_0^\infty f(t) dt$ is said to be integrable $|C, 1|_k, k \geq 1$, in the sense of Flett[3] if

$$(1.1) \quad \int_0^\infty x^{k-1} |\sigma'(x)|^k dx = \int_0^\infty \frac{|v(x)|^k}{x} dx$$

is convergent, where $v(x) = \frac{1}{x} \int_0^\infty t f(t) dt$ usually called a generator of the integral $\int_0^\infty f(t) dt$.

Let $p(x)$ be a non-decreasing real valued function on $[0, \infty)$. We define

$$P(x) = \int_0^x p(t) dt, p(x) \neq 0, p(0) = 0.$$

Then we define the Norlund mean of $s(x)$ by

$$\sigma_p(x) = \frac{1}{P(x)} \int_0^x p(t) s(t) dt.$$

We say that the improper integral $\int_0^\infty f(t) dt$ is summable $|N, p|_k, k \geq 1$, if

$$(1.2) \quad \int_0^\infty x^{k-1} |\sigma'_p(x)|^k dx$$

is convergent. In the special case if we take $p(x) = 1$ for all values of x , then $|N, p|_k$

summability reduces to $|C, 1|_k$ summability of improper integrals.

For any two functions f and g , it is customary to write $g(x) = O(f(x))$, if there exist K and N , for every

$x > N$, $\left| \frac{g(x)}{f(x)} \right| \leq K$. Clearly the difference between $s(x)$ and its n th weighted mean $\sigma_p(x)$, which is called the weighted

Kronecker identity, is given by the identity

$$(1.3) \quad s(x) - \sigma_p(x) = v_p(x),$$

where

$$v_p(x) = \frac{1}{P(x)} \int_0^\infty p(u) f(u) du.$$

We note that if we take $p(x) = 1$, for all values of x , then we have the following identity (see [2])

$$s(x) - \sigma_p(x) = v(x).$$

Since

$$\sigma'_p(x) = \frac{p(x)}{P(x)} v_p(x),$$

condition (1.3) can be written as

$$(1.4) \quad s(x) = v_p(x) + \int_0^x \frac{p(u)}{P(u)} v_p(u) du.$$

In view of the identity (1.4), the function $v_p(x)$ is called the generator function of $s(x)$.

Condition (1.1) can also be written as

$$(1.5) \quad \int_0^\infty x^{k-1} \left(\frac{p(x)}{P(x)} \right)^k |v_p(x)|^k dx$$

is convergent. It may be noted that for infinite series, an analogous definition was introduced by Orhan [4]. The improper integral

$\int_0^\infty f(t) dt$ is summable $|N, p; \delta|_k, k \geq 1, 0 \leq \delta k < 1$, if

$$(1.6) \quad \int_0^\infty x^{\delta k + k - 1} \left(\frac{p(x)}{P(x)} \right)^k |v_p(x)|^k dx$$

and summable $|N, p; \delta, \mu|_k, k \geq 1, 0 \leq \delta k < 1, \mu \geq 1$, if

$$(1.7) \quad \int_0^\infty x^{\mu(\delta k + k - 1)} \left(\frac{p(x)}{P(x)} \right)^k |v_p(x)|^k dx.$$

II. KNOWN RESULTS

Dealing with $|R, p_n|_k$ and $|R, q_n|_k$ summability methods, Orhan [4] proved the following theorem:

1) *Theorem 2.1.* The $|R, p_n|_k, (k \geq 1)$ summability implies the $|R, q_n|_k, (k \geq 1)$ summability provided that

$$a) \quad nq_n = O(Q_n),$$

$$b) \quad P_n = O(np_n) \quad \text{and}$$

$$c) \quad Q_n = O(nq_n).$$

Subsequently, dealing with summability of improper integrals, Ozgen has established the following theorem:

2) *Theorem 2.2.* [5] Let $p(x)$ and $q(x)$ be non-decreasing real valued functions on $[0, \infty)$ such that as $x \rightarrow \infty$

a) $xq(x) = O(Q(x))$,

b) $P(x) = O(xp(x))$,

and

c) $Q(x) = O(xq(x))$

If $\int_0^\infty f(t)dt$ is summable $|R, p|_k$, then it is also summable $|R, q|_k$, ($k \geq 1$).

Extending the result of Ozgen, Acharya[1] has established the following result.

3) *Theorem 2.3.* Let $p(x)$ and $q(x)$ be non-decreasing real valued functions on $[0, \infty)$ such that as $x \rightarrow \infty$

a) $xq(x) = O(Q(x))$,

b) $P(x) = O(xp(x))$,

c) $Q(x) = O(xq(x))$,

d) $\int_t^m \frac{x^{\delta k} q(x)}{Q^2(x)} dx = O\left(\frac{t^{\delta k}}{Q(t)}\right)$

and

e) $\int_0^m t^{\delta k - 1} |v_p(t)|^k dt = O(1)$.

If $\int_0^\infty f(t)dt$ is summable $|N, p; \delta|_k$, then it is also summable $|N, q, \delta|_k$, ($k \geq 1$).

However, extending the above result, in the present paper we establish the following theorem :

III. MAIN RESULT

1) *Theorem 3.1.* Let $p(x)$ and $q(x)$ be non-decreasing real valued functions on $[0, \infty)$ such that as $x \rightarrow \infty$

a) $xq(x) = O(Q(x))$,

b) $P(x) = O(xp(x))$,

c) $Q(x) = O(xq(x))$,

d) $\int_t^m \frac{x^{\mu(\delta k + k - 1) - k + 1} q(x)}{Q^2(x)} dx = O\left(\frac{t^{\mu(\delta k + k - 1) - k + 1}}{Q(t)}\right)$

and

e) $\int_0^m t^{\mu(\delta k + k - 1) - k} |v_p(t)|^k dt = O(1)$.

If $\int_0^\infty f(t)dt$ is summable $|N, p; \delta, \mu|_k$, then it is also summable $|N, q, \delta, \mu|_k$, ($k \geq 1$).

IV. PROOF OF THE THEOREM

Let $\sigma_p(x)$ and $\sigma_q(x)$ be the functions of (N, p) and (N, q) means of the integral $\int_0^\infty f(t)dt$. Since $\int_0^\infty f(t)dt$ is summable

$|N, p; \delta, \mu|_k$, $k \geq 1$, $0 \leq \delta k < 1$, $\mu \geq 1$, we can write

$$\int_0^\infty x^{\mu(\delta k + k - 1)} \left(\frac{p(x)}{P(x)} \right)^k |v_p(x)|^k dx$$

is convergent. Differentiating the equation (1.4), we have

$$f(x) = v_p'(x) + \frac{p(x)}{P(x)} v_p(x) .$$

By definition, we obtain

$$\sigma_q(x) = \frac{1}{Q(x)} \int_0^x q(t) \mathfrak{P}(t) dt = \frac{1}{Q(x)} \int_0^x (Q(x) - Q(t)) f(t) dt$$

and

$$\begin{aligned} \sigma_q'(x) &= \frac{q(x)}{Q^2(x)} \int_0^x Q(t) f(t) dt \\ &= \frac{q(x)}{Q^2(x)} \int_0^x Q(t) \left[v_p'(t) + \frac{p(t)}{P(t)} v_p(t) \right] dt \\ &= \frac{q(x)}{Q^2(x)} \int_0^x Q(t) v_p'(t) dt + \frac{q(x)}{Q^2(x)} \int_0^x Q(t) \frac{p(t)}{P(t)} v_p(t) dt \end{aligned}$$

Integrating by parts of the first statement, we have

$$\begin{aligned} \sigma_q'(x) &= \frac{q(x)}{Q^2(x)} \left[Q(x) v_p(x) - \int_0^x q(t) v_p(t) dt \right] + \frac{q(x)}{Q^2(x)} \int_0^x Q(t) \frac{p(t)}{P(t)} v_p(t) dt \\ &= \frac{q(x)}{Q^2(x)} v_p(x) + \frac{q(x)}{Q^2(x)} \int_0^x Q(t) \frac{p(t)}{P(t)} v_p(t) dt - \frac{q(x)}{Q^2(x)} \int_0^x q(t) v_p(t) dt \\ &= \sigma_{q,1}(x) + \sigma_{q,2}(x) + \sigma_{q,3}(x), \text{ say.} \end{aligned}$$

In order to complete the proof of the theorem, it is sufficient to show that

$$\int_0^m x^{\mu(\delta k + k - 1)} |\sigma_{q,r}(x)|^k dx = O(1) \text{ as } m \rightarrow \infty, \text{ for } r = 1, 2, 3 .$$

Using conditions (7.3.1.1) and (7.3.1.2), we have

$$\begin{aligned} \int_0^m x^{\mu(\delta k + k - 1)} |\sigma_{q,1}(x)|^k dx &= \int_0^m x^{\mu(\delta k + k - 1)} \left| \frac{q(x)}{Q(x)} v_p(x) \right|^k dx \\ &= \int_0^m x^{\mu(\delta k + k - 1)} \left(\frac{q(x)}{Q(x)} \right)^k |v_p(x)|^k dx \\ &= O(1) \int_0^m x^{\mu(\delta k + k - 1)} \left(\frac{p(x)}{P(x)} \right)^k |v_p(x)|^k dx \\ &= O(1) \int_0^m x^{\mu(\delta k + k - 1)} |\sigma_p'(x)|^k dx \\ &= O(1) , \text{ as } m \rightarrow \infty , \end{aligned}$$

by virtue of the hypotheses of theorem 3.1.

Applying Holder's inequality with $k > 1$, we get

$$\int_0^m x^{\mu(\delta k + k - 1)} |\sigma_{q,2}(x)|^k dx = O(1) \int_0^m x^{\mu(\delta k + k - 1)} \left(\frac{q(x)}{Q^2(x)} \right)^k \left(\int_0^x \frac{Q(t) p(t)}{P(t)} |v_p(t)| dt \right)^k dx$$

$$\begin{aligned}
 &= O(1) \int_0^m \frac{x^{\mu(\delta k+k-1)-k+1} q(x)}{Q^{k+1}(x)} \left(\int_0^x \frac{Q(t)p(t)}{P(t)} |v_p(t)| dt \right)^k dx \\
 &= O(1) \int_0^m \frac{x^{\mu(\delta k+k-1)-k+1} q(x)}{Q^2(x)} \left(\int_0^x \left(\frac{Q(t)}{q(t)} \right)^k q(t) \left(\frac{p(t)}{P(t)} \right)^k |v_p(t)|^k dt \right) x \left(\frac{1}{Q(x)} \int_0^x q(t) dt \right)^{k-1} dx \\
 &= O(1) \int_0^m t^k q(t) \left(\frac{p(t)}{P(t)} \right)^k |v_p(t)|^k dt \int_t^m \frac{x^{\mu(\delta k+k-1)-k+1} q(x)}{Q^2(x)} dx \\
 &= O(1) \int_0^m t^{\mu(\delta k+k-1)+1} \frac{q(t)}{Q(t)} \left(\frac{p(t)}{P(t)} \right)^k |v_p(t)|^k dt \\
 &= O(1) \int_0^m t^{\mu(\delta k+k-1)} \left(\frac{p(t)}{P(t)} \right)^k |v_p(t)|^k dt \\
 &= O(1) \text{ as } m \rightarrow \infty,
 \end{aligned}$$

by virtue of the hypotheses of theorem 7.3.1.

Finally, again by Holder's inequality with $k > 1$, we have

$$\begin{aligned}
 \int_0^m x^{\mu(\delta k+k-1)} |\sigma_{q,3}(x)|^k dx &= O(1) \int_0^m x^{\mu(\delta k+k-1)} \left(\frac{q(x)}{Q^2(x)} \right)^k \left(\int_0^x q(t) |v_p(t)|^k dt \right)^k dx \\
 &= O(1) \int_0^m \frac{x^{\mu(\delta k+k-1)-k+1} q(x)}{Q^2(x)} \left(\int_0^x q(t) |v_p(t)|^k dt \right) x \left(\frac{1}{Q(x)} \int_0^x q(t) dt \right)^{k-1} dx \\
 &= O(1) \int_0^m q(t) |v_p(t)|^k dt \int_t^m \frac{x^{\mu(\delta k+k-1)-k+1} q(x)}{Q^2(x)} dx \\
 &= O(1) \int_0^m \frac{t^{\mu(\delta k+k-1)-k+1} q(t)}{Q(t)} |v_p(t)|^k dt \\
 &= O(1) \int_0^m t^{\mu(\delta k+k-1)-k} |v_p(t)|^k dt \\
 &= O(1) \text{ as } m \rightarrow \infty,
 \end{aligned}$$

by virtue of the hypotheses of theorem 3.1.

This completes the proof of the theorem.

REFERENCES

- [1] D. Acharya, Ph.D. thesis submitted to Fakir Mohan University, Odisha, (2019)
- [2] Canak and U. Totur, "A tauberian theorem for Cesaro summability factors of integrals," Appl. Math. Lett. vol. 24, no. 3, pp. 234-245, 2011
- [3] T.M. Flett, "On an extension of absolute summability and some theorems of littlewood and paley," Proc. London Math. Soc, vol.7, pp.113-141, 1957
- [4] C. Orhan, "On equivalence of summability methods, Math.Slovaca" vol.40, no.2, pp.171-175, 1990.
- [5] H.N. Ozgen, "On equivalence of two integrability methods", Miskole Mathematical Notes, Vol.18 (1), pp.391-396, (2017)



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)