



IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 7 Issue: VI Month of publication: June 2019

DOI: http://doi.org/10.22214/ijraset.2019.6121

www.ijraset.com

Call: 🕥 08813907089 🔰 E-mail ID: ijraset@gmail.com



Avifaunal Diversity of Bisalpur Dam and nearby Area of Tehsil Deoli, District Tonk (Rajasthan), India

Dinesh Meena¹, Priti Kaushik¹, Vivek Sharma², Divaker Yadav², Anita Gajraj³, Jai Bahadur Singh Kachhawa⁴

¹Department of Zoology, School of Basic and Applied Science, Poornima University, Jaipur, Rajasthan, India ²Biodiversity Research Laboratory, Department of Zoology, Maharashi Dayanand Saraswati University, Ajmer, Rajasthan, India ³Department of Zoology, Kanoria P.G. Mahila Mahavidyalaya, Jaipur, Rajasthan, India ⁴Centre for Advanced Research and Development, Jaipur, Rajasthan, India.

Abstract: Avifauna is the term used for the birds of a particular region, habitat, or geological period. Study area covers foothills of beautiful Aravali ranges. It is one of the world's oldest mountain ranges. Bisalpur Dam is avifaunal diversity rich area situated in district Tonk, Rajasthan. Deoli is 165 Kms away from Jaipur, 85 Kms from Kota and 145 Kms from Ajmer. The dam was constructed between two mountains on Banas river for the purpose of drinking water and irrigation. This dam attracts human beings as well as birds also. Nesting spots are easy to find along the bank of Banas river. Visual encounter method and different methods were used for the study purpose. Identification was done with the help of field guide books. Total 176 species of birds belonging to different family were spotted in the study area. Keyworss: Avifauna, bisalpur, diversity, terrestrial and water birds

I. INTRODUCTION

Birds are excellent indicators of water quality and measures of biodiversity. The present study was conducted to understand the diversity measures of avifauna in and nearby places of Bisalpura dam.

A. Study Area

Study area is Bisalpur Dam and nearby area. Bisalpur dam is constructed on Banas river near tehsil Deoli, district Tonk, Rajasthan, India (Figure 1-4)). The coordinates are 25°55'20.56" N & 75°27' 20.51" E. with an elevation of 1081 feet. The height of dam is 130 feet and length is 1883 feet; gross storage capacity is 1.10 KM³. Reservoir area is 218.36 KM². Effective storage capacity is 1.04 KM³. This Dam was completed in 1999. It supplies irrigation and drinking water for many towns of Rajasthan. Bisalpur dam and nearby area is rich in Avifaunal Diversity.




Figure 1: Study area in map



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.177 Volume 7 Issue VI, June 2019- Available at www.ijraset.com




Figure 2: Study area in map (Enlarge)



Figure 3: Broad view of Bisalpur Dam



Figure 4: Flora at study area

II. MATERIALS & METHODOLOGY:

A. Materials

Different apparatus i.e. digital camera (Canon SX10, Canon 1200D), binoculars, field guide and observation sheets were used for the present study.

B. Survey

Regular surveys were made at different time interval between 05:00 to 22:00 Hrs.

The study was based on the survey of the area. Quantitative avifauna data was collected in the point count survey and combined with distance estimation with the help of binocular and camera. Bisalpur Dam and nearby area was selected as site for observation of avifauna. These areas are rich in avifaunal diversity. Periodical studies were made to know about the status of birds. Direct count methods were adopted from *Bibbey et al.* [1] and Javad and Kaul [2] for recording and analysis. Birds were identified with the help of field guide [3,4,5,6,7,8]. Visual encounter method was also used. Most of the birds species were spotted during the winter season. Regular surveys were made near the breeding grounds. Representative cases were photographed from different angles in the nature. Morphometric observations were also made for the taxonomic identification and compared with the standard diagnostic keys. Birds were divided in three categories MC-most common, C-common, R-rare according to the number of individuals. We also described the IUCN status of the birds.



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.177 Volume 7 Issue VI, June 2019- Available at www.ijraset.com

III. RESULT & DISCUSSION

Total 176 species of birds belonging to different family were present in the study area. Avifaunal diversity was divided into two categories one is water birds (Table 1) and another is terrestrial birds (Table 2) found near water bodies.

| S. No. | Scientific name | Common name | IUCN status | Migratory status | Population trend |
|--------|--|---|--|------------------|---------------------|
| | Anatidae | | | | |
| 1 | Anser indicus, (Latham, 1790) | Bar-headed goose | Least Concern | Winter Visitor | Decreasing |
| 2 | Anas acuta, (Linnaeus, 1758) | Northern Pintail | Least Concern | Winter Visitor | Decreasing |
| 3 | Anas clypeata, (Linnaeus, 1758) | Northern shoveler | Least Concern | Winter Visitor | Decreasing |
| 4 | Anas crecca, (Linnaeus, 1758) | Eurasian teal/ Common Teal | Least Concern | Winter Visitor | Decreasing |
| 5 | Anas penelope, (Linnaeus, 1758) | Eurasian wigeon | Least Concern | Winter Visitor | Decreasing |
| 6 | Anas poecilorhyncha, (Forster, 1781) | Indian Spot-billed duck | Least Concern | Resident | Decreasing |
| 7 | Anas querquedula (Linnaeus, 1758) | Garganey | Least Concern | Winter Visitor | Decreasing |
| 8 | Anas strepera (Linnaeus, 1758) | Gadwall | Least Concern | Winter Visitor | Decreasing |
| 9 | Aythya ferina (Linnaeus, 1758) | Common Pochard | Vulnerable | Winter Visitor | Decreasing |
| 10 | Aythya fuligula, (Linnaeus, 1758) | Tufted duck | Least Concern | Winter Visitor | Stable |
| 11 | Aythya nyroca, (Güldenstädt, 1770) | Ferruginous pochard/ Ferruginous Duck | Near Threatened | Winter Visitor | Decreasing |
| 12 | Sarkidiornis melanotos, (Pennant, 1769) | Comb duck/ knob-billed duck | Least Concern | Resident | Decreasing |
| 13 | Tadorna ferruginea, (Pallas, 1764) | Ruddy shelduck | Least Concern | Winter Visitor | Stable |
| | Ardeidae | | | | |
| 14 | Ardea alba, (Linnaeus, 1758) | Great Egret/ Great white Egret | Least Concern | Resident | stable |
| 15 | Ardea cinerea, (Linnaeus, 1758) | Grey heron | Least Concern | Resident | Stable |
| 16 | Ardea purpurea, (Linnaeus, 1766) | Purple heron | Least Concern | Resident | Decreasing |
| 17 | Ardeola grayii, (Sykes, 1832) | Indian Pond Heron | Least Concern | Resident | Increasing |
| 18 | Bubulcus ibis (Linnaeus, 1758) | Cattle Egret | Least Concern | Resident | Increasing |
| 19 | Butorides Striatus (Linnaeus, 1758) | Striated Heron | This taxon has not yet been assessed for the IUCN Red List, but is in the Catalogue of Life | Winter Visitor | Unknown |
| 20 | Egretta garzetta, (Linnaeus, 1766) | Little Egret | Least Concern | Resident | Increasing |

Table 1: List of water birds spotted in study area



| | | 1 | | | |
|----|---|--------------------------------------|--|-----------------|------------|
| 21 | Mesophoyx intermedia (Wagler, 1827) | Intermediate Egret | This taxon has not yet been assessed for the IUCN Red List, but is in the Catalogue of Life | Resident | Unknown |
| 22 | Nycticorax nycticorax, (Linnaeus, 1758) | Black-crowned night heron | Least Concern | Resident | Decreasing |
| | Cerylidae | | | | |
| 23 | Ceryle rudis (Linnaeus 1758) | Pied Kingfisher | Least Concern | Winter Visitor | Decreasing |
| | Charadriidae | | | | |
| 24 | Charadrius dubius (Scopoli, 1786) | Little ringed plover | Least Concern | Resident | Stable |
| 25 | Vanellus indicus (Boddaert, 1783) | Red-wattled lapwing | Least Concern | Resident | Increasing |
| 26 | Vanellus malabaricus (Boddaert, 1783) | Yellow-wattled lapwing | Least Concern | Resident | Stable |
| | Ciconiidae | | | | |
| 27 | Anastomus oscitans (Boddaert, 1783) | Asian Open bill | Least Concern | Winter Visitor | Stable |
| 28 | Mycteria leucocephala (Pennant, 1769) | Painted stork | Near Threatened | winter Visitor | Decreasing |
| | Jacanidae | | | | |
| 29 | Hydrophasianus chirurgus (Scopoli, 1786) | Pheasant-tailed jacana | Least Concern | Resident | Decreasing |
| | Laridae | | | | |
| 30 | Chroicocephalus brunnicephalus (Jerdon, 1840) | Brown Headed Gull | This taxon has not yet been assessed for the IUCN Red List, but is in the Catalogue of Life | Winter Visitor | Unknown |
| 31 | Chroicocephalus ridibundus (Linnaeus,1766) | Black Headed gull | Least Concern | Winter Visitor | Decreasing |
| | Pelecanidae | | | | |
| 32 | Pelecanus onocrotalus (Linnaeus, 1758) | Great white pelican/ Rosy pelican | Least Concern | Winter Visitor | Stable |
| 33 | Pelecanus crispus (Bruch, 1832) | Dalmatian pelican | Vulnerable | Winter Visitor | Decreasing |
| | Phalacrororacidae | | | | |
| 34 | Phalacrocorax carbo (Linnaeus, 1758) | Great Cormorant | Least Concern | Resident | Increasing |
| 35 | Phalacrocorax fuscicollis (Stephens, 1826) | Indian Cormorant | Least Concern | Resident | Stable |
| 36 | Phalacrocorax niger (Vieillot, 1817) | Little Cormorant | Least Concern | Resident | Stable |
| | Phoenicopteridae | | | | |
| 37 | Phoenicopterus roseus (Linnaeus, 1758) | Greater Flamingo | Least Concern | Passage Visitor | Decreasing |
| | Podicipedidae | | | | |
| 38 | Tachybaptus ruficollis (Pallas, 1764) | Little grebe | Least Concern | Resident | Decreasing |
| | Rallidae | | | | |



| | Amaurornis | White-breasted | | | |
|------------|-------------------------|--------------------------------|------------------------------|-------------------|------------|
| 39 | phoenicurus | waterhen | Least Concern | Resident | Stable |
| | (Pennant, 1769) | | | | |
| 40 | Fulica atra | Common Coot / | Least Concern | Resident | Decreasing |
| | (Linnaeus, 1758) | Eurasian coot | | | |
| 41 | Gallinula chloropus | Common moorhen | Common moorhen Least Concern | | Stable |
| | (Linnaeus, 1758) | | | Resident | |
| 42 | Porphyrio Porphyrio | Purple swamp hen Least Concern | | Resident | Increasing |
| | (Linnaeus, 1758) | | | | 6 |
| | Recurvirostridae | | | | |
| 10 | Himantopus | | | D 11 | - · |
| 43 | himantopus | Black-winged Stilt | Least Concern | Resident | Increasing |
| | (Linnaeus, 1758) | | | | |
| 44 | Recurvirostra avosetta | Pied avocet | Least Concern | Winter Visitor | Stable |
| | (Linnaeus, 1758) | | | | |
| | Rostratulidae | | | | |
| 45 | Rostratula benghalensis | Greater Painted- | Least Concern | Winter Visitor | Decreasing |
| | (Linnaeus, 1758) | snipe | | | |
| | Scolopacidae | | | | |
| 46 | Actitis hypoleucos | Common sandpiper | Least Concern | Winter Visitor | Decreasing |
| - | (Linnaeus, 1758) | | | | Decreasing |
| 47 | Calidris minuta | Little Stint | Least Concern | Winter Visitor | Decreasing |
| | (Leisler, 1812) | | | | |
| 48 | 48 Calidris temminckii | Temminck's Stint | Least Concern | Winter Visitor | Unknown |
| | (Leisler, 1812) | | | | |
| 49 | Limosa limosa | Black-tailed godwit | Near Threatened | Winter Visitor | Decreasing |
| | (Linnaeus, 1758) | | | | |
| 50 | Philomachus pugnax | Ruff | Least Concern | Winter Visitor | Decreasing |
| | (Linnaeus, 1758) | | | | 6 |
| 51 | Tringa erythropus | Spotted redshank | Least Concern | Winter Visitor | Stable |
| | (Pallas, 1764) | • | | | |
| 52 | Tringa glareola | Wood sandpiper | Least Concern | Winter Visitor | Stable |
| | (Linnaeus, 1758) | | | | |
| 53 | Tringa nebularia | Common | Least Concern | Winter Visitor | Stable |
| | (Gunnerus, 1767) | greenshank | | | |
| 54 | Tringa ochropus | Green sandpiper | Least Concern | Winter Visitor | Stable |
| | (Linnaeus, 1758) | | | | |
| 55 | Tringa tetanus | Common redshank | Least Concern | Winter Visitor | Stable |
| | (Linnaeus, 1758) | | | | |
| | Sternidae | | | | |
| 56 | Chlidonias hybridus, | Whiskered tern | Least Concern | Winter Visitor | Stable |
| | (Pallas, 1811) | Lation D' (| | | |
| 57 | Sterna aurantia | Indian River tern/ | Near Threatened | Resident | Decreasing |
| | (Gray, JE, 1831) | River tern | | | |
| | Threskiornithidae | E | | | |
| 58 | Platalea leucorodia | Eurasian spoonbill/ | Least Concern | Winter Visitor | Stable |
| | (Linnaeus, 1758) | Common spoonbill | | | |
| 59 | Pseudibis papillosa | Red-naped Ibis/ | Least Concern | Winter Visitor | Decreasing |
| | (Temminck, 1824) | Black Ibis | | | |
| <i>c</i> 0 | Threskiornis | Black-headed ibis/ | | D 1 | D |
| 60 | melanocephalus | Oriental white ibis | Near Threatened | Resident | Decreasing |
| | (Latham, 1790) | | | | |



International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.177

Volume 7 Issue VI, June 2019- Available at www.ijraset.com

| Table 2: List of terrestrial birds spotted in study area | | | | | | |
|--|---|--|---------------|------------------|------------------|--|
| S. No. | Scientific name | Common name | IUCN status | Migratory status | Population trend | |
| | Accipitridae | | | | | |
| 1 | Accipiter badius (Gmelin, 1788) | Shikra | Least Concern | Resident | Decreasing | |
| 2 | Aquila nipalensis (Hadgson, 1833) | Steppe Eagle | Endangered | Resident | Decreasing | |
| 3 | Buteo Rufinus (Cretzschmar, 1829) | Long Legged Buzzard | Least Concern | Resident | Stable | |
| 4 | Circaetus gallicus (Gmelin, 1788) | Short-toed Snake- Eagle | Least Concern | Resident | Stable | |
| 5 | Circus aeruginosus (Linnaeus, 1758) | Eurasian Marsh Harrier | Least Concern | Winter visitor | Increasing | |
| 6 | Circus pygargus (Linnaeus, 1758) | Montagu's Harrier | Least Concern | Winter Visitor | Decreasing | |
| 7 | Elanus caeruleus (Desfontaines, 1789) | Black-Winged Kite | Least Concern | Resident | Stable | |
| 8 | Milvus migrans (Boddaert, 1783) | Black Kite | Least Concern | Resident | Stable | |
| 9 | Neophron percnopterus (Linnaeus, 1758) | Egyptian Vulture | Endangered | Resident | Decreasing | |
| 10 | Spilornis cheela (Latham, 1790) | Crested Serpent Eagle | Least Concern | Resident | Stable | |
| | Aegithinidae | | | | | |
| 11 | Aegithina tiphia (Linnaeus, 1758) | Common Iora | Least Concern | Resident | Unknown | |
| | Alaudidae | | | | | |
| 12 | Eremopterix grisea (Scopoli, 1786) | Ashy-crowned Sparrow-Lark | Least concern | Resident | Stable | |
| 13 | Galerida cristata (Linnaeus, 1758) | Crested Lark | Least concern | Resident | Stable | |
| 14 | Mirafra cantillans (Blyth, 1845) | Singing Bush-Lark | Least concern | Resident | Decreasing | |
| 15 | Mirafra erythroptera (Blyth, 1845) | Indian Bush-Lark | Least concern | Resident | Stable | |
| | Alcedinidae | | | | | |
| 16 | Alcedo atthis (Linnaeus, 1758) | Common Kingfisher | Least Concern | Resident | Decreasing | |
| | Apodidae | | | | | |
| 17 | Apus affinis (John Edward Gray, 1830) | Little Swift / House Swift | Least concern | Resident | Stable | |
| | Bucerotidae | | | | | |
| 18 | Ocyceros birostris (Scopoli, 1786) | Indian Grey Hornbill | Least concern | Winter visitor | Decreasing | |
| | Burhinidae | | | | | |
| 19 | Burhinus oedicnemus (Linnaeus, 1758) | Eurasian Stone- Curlew / Eurasian thicknee | Least concern | Winter visitor | Decreasing | |
| | Caprimulgidae | | | | | |
| 20 | Caprimulgus affinis (Horsfield, 1821) | Savanna Nightjar | Least concern | Resident | Decreasing | |
| 21 | Caprimulgus asiaticus (Latham, 1790) | Indian Nightjar | Least concern | Resident | Decreasing | |
| | | 1 | | | | |

Table 2: List of terrestrial birds spotted in study area

Cisticolidae



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.177

Volume 7 Issue VI, June 2019- Available at www.ijraset.com

| 22 | Orthotomus sutorius (Pennant, 1769) | Common Tailorbird | Least concern | Resident | Decreasing |
|----|---|--|---------------|-----------------|------------|
| 23 | Prinia hodgsonii (Blyth, 1844) | Grey Breasted Prinia | Least concern | Resident | Decreasing |
| 24 | Prinia inornata (Sykes,1832) | Plain Prinia/ White browed Wren-warbler | Least Concern | Resident | Stable |
| 25 | Prinia socialis (Sykes, 1832) | Ashy Prinia / Ashy wren-warbler | Least concern | Resident | Decreasing |
| | Columbidae | | | | |
| 26 | Columba livia (Gmelin, 1789) | Rock Pigeon | Least concern | Resident | Stable |
| 27 | Streptopelia decaocto (Frivaldszky, 1838) | Eurasian Collared- Dove | Least concern | Resident | Stable |
| 28 | Streptopelia senegalensis (Linnaeus, 1766) | Laughing Dove | Least concern | Resident | Stable |
| 29 | Streptopelia tranquebarica (Hermann, 1804) | Red Collared-Dove | Least concern | Resident | Stable |
| 30 | Treron phoenicoptera (Latham, 1790) | Yellow-Footed Green-Pigeon | Least concern | Winter visitor | Decreasing |
| | Coraciidae | | | | |
| 31 | Coracias benghalensis (Linnaeus, 1758) | Indian Roller | Least concern | Resident | Stable |
| 32 | Coracias garrulous (Linnaeus, 1758) | European Roller | Least Concern | Passage Visitor | Decreases |
| | Corvidae | | | | |
| 33 | Corvus macrorhynchos (Wagler, 1827) | Jungle Crow | Least concern | Resident | Decreasing |
| 34 | Corvus splendens (Vieillot, 1817) | House Crow | Least concern | Resident | Increasing |
| 35 | Dendrocitta vagabunda (Latham, 1790) | Rufous Treepie | Least concern | Resident | Stable |
| | Cuculidae | | | | |
| 36 | Centropus sinensis (Stephens, 1815) | Greater Coucal | Least concern | Resident | Stable |
| 37 | Clamator jacobinus (Boddaert,1783) | Jacobin cuckoo/ Pied cuckoo/ Pied crested cuckoo | Least Concern | Summer Visitor | Stable |
| 38 | Eudynamys scolopacea (Linnaeus, 1758) | Asian Koel | Least concern | Resident | Stable |
| 39 | Hierococcyx varius (Vahl, 1797) | Common hawk- cuckoo/ Brainfever Bird | Least concern | Resident | stable |
| 40 | Phaenicophaeus leschenaultia (Lesson, 1830) | Sirkeer Malkoha | Least concern | Resident | Stable |
| | Dicruridae | | | | |
| 41 | Dicrurus caerulescens (Linnaeus, 1758) | White bellied drongo | Least Concern | Resident | Unknown |
| 42 | Dicrurus macrocercus (Vieillot, 1817) | Black Drongo | Least concern | Resident | Increasing |
| | Emberizidae | | | | |
| 43 | Melophus lathami (Gray, 1831) | Crested Bunting | Least concern | Resident | Stable |
| 44 | Emberiza bruniceps (Brandt, 1841) | Red headed bunting | Least Concern | Winter Visitor | Stable |
| | Estrildidae | | | | |



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.177

Volume 7 Issue VI, June 2019- Available at www.ijraset.com

| 45 | Lonchura malabarica (Linnaeus, 1758) | Indian Silver bill | Least concern | Resident | stable |
|----|--|---|--|----------------|------------|
| | Falconidae | | | | |
| 46 | Falco tinnunculus (Linnaeus, 1758) | Common Kestrel | Least Concern | Winter visitor | Decreasing |
| | Glareolidae | | | | |
| 47 | Cursorius coromandelicus (Gmelin, 1789) | Indian Courser | Least concern | Resident | Decreasing |
| 48 | Glareola lacteal (Temminck,1820) | Little Pratincole/Small Indian Pratincole | Least Concern | Winter Visitor | unknown |
| | Halcyonidae | | | | |
| 49 | Halcyon smyrnensis (Olitari,1758) | white-throated kingfisher | Least Concern | Resident | Decreasing |
| | Hirudinidae | | | | |
| 50 | Cecropis daurica (Laxmann, 1769) | Red Rumped swallow | This taxon has not yet been assessed for the IUCN Red List, but is in the Catalogue of Life | Resident | Unknown |
| 51 | Hirundo rustica (Linnaeus, 1758) | Barn Swallow | Least Concern | Resident | Decreasing |
| 52 | Hirundo smithii (Leach, 1818) | Wire-tailed Swallow | Least Concern | Resident | Increasing |
| 53 | Petrochelidon fluvicola (Blyth, 1855) | Streak-throated- swallow | This taxon has not yet been assessed for the IUCN Red List, but is in the Catalogue of Life | Resident | Unknown |
| 54 | Ptyonoprogne concolor (Sykes, 1832) | Dusky crag martin | This taxon has not yet been assessed for the IUCN Red List, but is in the Catalogue of Life | Resident | Unknown |
| | Laniidae | | | | |
| 55 | Lanius meridionalis (Temminch, 1820) | Southern Grey Shrike | *This taxon has not yet been assessed for the IUCN Red List, but is in the Catalogue of Life | Resident | Stable |
| 56 | Lanius schach (Linnaeus, 1758) | Long tailed shrike /Rufous-backed Shrike | Least concern | Resident | Stable |
| 57 | Lanius vittatus (Valenciennes, 1826) | Bay-backed Shrike | Least concern | Resident | stable |
| | Leiothrichidae | | | | |
| 58 | Turdoides caudatus (Dumont, 1823) | Common Babbler | Least concern | Resident | Stable |
| 59 | Turdoides malcolmi (Sykes, 1832) | Large Grey Babbler | Least concern | Resident | stable |
| 60 | Turdoides striatus (Dumont, 1823) | Jungle Babbler | Least concern | Resident | Stable |
| | Megalaimidae | | | | |
| 61 | Megalaima haemacephala (Philipp Ludwig Statius | Coppersmith Barbet | Least concern | Resident | Decreasing |



| | Müller,1776) | | | | |
|----|--|--|---|------------------------------|------------|
| | Meropidae | | | | |
| 62 | Merops orientalis (Latham, 1801) | Green Bee-eater | Least concern | Resident | Increasing |
| 63 | Merops persicus (Pallas, 1773) | Blue-Cheeked Bee- Eater | Least concern | Least concern Summer Visitor | |
| 64 | Merops philippinus (Linnaeus, 1766) | Blue Tailed-bee-eater | Least Concern | Resident | Stable |
| | Monarchidae | | | | |
| 65 | Terpsiphone paradise (Linnaeus, 1758) | Indian Paradise flycatcher/ Asian Paradise flycatcher | Least Concern | Summer Visitor | Stable |
| | Motacillidae | | | | |
| 66 | Anthus rufulus (Vieillot, 1818) | Paddyfield pipit/ Oriental pipit | Least Concern | Resident | Stable |
| 67 | Motacilla alba (Linnaeus, 1758) | White Wagtail | Least Concern | Resident | Stable |
| 68 | Motacilla cinerea (Tunstall, 1771) | Grey wagtail | Least concern | Resident | Stable |
| 69 | Motacilla Citreola (Pallas, 1776) | Citrine wagtail | Least concern | Resident | Stable |
| 70 | Motacilla flava (Linnaeus, 1758) | Yellow wagtail | Least concern | Resident | Decreasing |
| 71 | Motacilla maderaspatensis (Gmelin, 1789) | White-browed wagtail | This taxon has not yet been assessed for the IUCN Red List, but is in the Catalogue of Life | Resident | Unknown |
| | Muscicapidae | | - | | |
| 72 | Cercomela fusca (Blyth, 1851) | Brown rock Chat | Least concern | Resident | Stable |
| 73 | Copsychus saularis (Linnaeus, 1758) | Oriental Magpie- Robin | Least concern | Resident | Stable |
| 74 | Ficedula parva (Bechstein, 1792) | Red-breasted flycatcher | Least Concern | Winter Visitor | Stable |
| 75 | Luscinia svecica (Linnaeus, 1758) | Blue throat | Least concern | Winter visitor | Stable |
| 76 | Monticola solitarius (Linnaeus, 1758) | Blue Rock-Thrush | least concern | Winter visitor | Stable |
| 77 | Oenanthe deserti (Temminck, 1829) | Desert wheatear | Least Concern | Winter Visitor | Stable |
| 78 | Oenanthe picata (Blyth, 1847) | Variable Wheatear | Least concern | Winter visitor | Stable |
| 79 | Phoenicurus ochruros (S.G.Gmelin, 1774) | Black redstart | Least Concern | Winter Visitor | Increasing |
| 80 | Saxicola caprata (Linnaeus, 1766) | Pied Bushchat | Least concern | Winter visitor | Stable |
| 81 | Saxicola torquata (Linnaeus, 1766) | Common Stonechat | Least concern | Winter visitor | Stable |
| 82 | Saxicoloides fulicata (Linnaeus, 1776) | Indian Robin | Least concern | Resident | Stable |
| | Nectariniidae | | | | |
| 83 | Nectarinia asiatica (Latham, 1790) | Purple Sunbird | Least concern | Resident | Stable |
| | Oriolidae Oriolus oriolus | Eurasian Golden | | | |
| 84 | (Linnaeus, 1758) | Oriole | Least concern | Summer Visitor | Stable |



| | Passeridae | | | | |
|-----|--|----------------------------|-----------------|----------------|--------------|
| | Passer domesticus | | | | |
| 85 | (Linnaeus, 1758) | House Sparrow | Least concern | Resident | Increasing |
| | Phasianidae | | | | |
| | Francolinus | | | | |
| 86 | pondicerianus | Grey Francolin | Least Concern | Resident | Stable |
| | (Gmelin, 1789) | | | | |
| 87 | Pavo cristatus | Indian Peafowl | Least Concern | Resident | Decreasing |
| ÷. | (Linnaeus, 1758) | | | | |
| | Phylloscopidae | | | | |
| 88 | Phylloscopus collybita | Common Chiffchaff | Least concern | Winter visitor | Decreasing |
| | (Vieillot, 1817) | | | | |
| | Picidae | TT TI 1. 1 | | | |
| 89 | Chrysocolaptes festivus | White-naped | Least Concern | Residential | Stable |
| | (Boddaert, 1783) | woodpecker | | | |
| 90 | Dinopium benghalense (Linnaeus, 1758) | Black-rumped Woodpecker | Least concern | Resident | Stable |
| | Jynx torquilla | wooupeckei | | | |
| 91 | (Linnaeus, 1758) | Eurasian wryneck | Least Concern | Winter Visitor | Decreasing |
| | Picoides nanus | Brown-capped | | | |
| 92 | (Vigors, 1832) | Pygmy Woodpecker | Least concern | Resident | Stable |
| | Ploceidae | - ,8, | | | |
| | Ploceus philippinus | | | | |
| 93 | (Linnaeus, 1766) | Baya Weaver | Least concern | Resident | Stable |
| | Psittacidae | | | | |
| | Psittacula cyanocephala | Plum-headed | _ | | |
| 94 | (Linnaeus, 1766) | Parakeet | Least concern | Winter visitor | Decreasing |
| 05 | Psittacula eupatria | Aleman Joins Developed | No thursday a d | Desident | Demociae |
| 95 | (Linnaeus, 1766) | Alexandrine Parakeet | Near threatened | Resident | Decreasing |
| 96 | Psittacula krameri | Dogo wincod Donalsoot | L sost somsom | Resident | In ana ain a |
| 90 | (Scopoli, 1769) | Rose-ringed Parakeet | Least concern | Kesidem | Increasing |
| | Pteroclididae | | | | |
| 97 | Pterocles Indicus | Painted sandgrous | Least concern | Resident | Stable |
| 71 | (Gmelin, 1789) | - | Least concern | Resident | Stable |
| 98 | Pterocles exustus | Chestnut-bellied | Least concern | Resident | Stable |
| | (Temminck, 1825) | sandgrouse | | | ~~~~~ |
| | Pycnonotidae | | | | |
| 99 | Pycnonotus cafer | Red-vented Bulbul | Least concern | Resident | Increasing |
| | (Linnaeus, 1766) | | | | _ |
| 100 | Pycnonotus leucotis | White-eared bulbul | Least concern | Resident | Decreasing |
| | (Gould, 1836) | | | | _ |
| | Strigidae Asio flammeus | | | | |
| 101 | (Pontoppidan, 1763) | Short-eared Owl | Least concern | Winter visitor | Decreasing |
| | Athene brama | | | | |
| 102 | (Temminck, 1821) | Spotted Owlet | Least concern | Resident | Stable |
| | Bubo bubo | | | | |
| 103 | (Linnaeus, 1758) | Eurasian Eagle-Owl | Least concern | Resident | Decreasing |
| | Sturnidae | | | 1 | |
| 101 | Acridotheres ginginianus | D 114 | | D | - · |
| 104 | (Latham, 1790) | Bank Myna | Least concern | Resident | Increasing |
| 105 | Acridotheres tristis | G 14 | . . | D 11. | |
| 105 | (Linnaeus, 1766) | Common Myna | Least concern | Resident | Increasing |
| 107 | Sturnus contra | Asian Dia 1.07 1 | T | D 1 - / | 0, 11 |
| 106 | (Linnaeus, 1758) | Asian Pied Starling | Least concern | Resident | Stable |
| 107 | Sturnus pagodarum | Ducharing St. 1 | Loot | Destate a | : |
| 107 | (Gmelin, 1789) | Brahminy Starling | Least concern | Resident | increasing |
| 109 | Sturnus roseus | Boon Starling | Loost concern | Winten willten | Ctobl- |
| 108 | (Linnaeus, 1758) | Rosy Starling | Least concern | Winter visitor | Stable |
| 109 | Sturnus vulgaris | Common Starling | Least concern | Winter visitor | Decreasing |



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.177 Volume 7 Issue VI, June 2019- Available at www.ijraset.com

| | (Linnaeus, 1758) | | | | |
|-----|--|------------------------------|---------------|----------------|------------|
| | Sylviidae | | | | |
| 110 | Sylvia althaea (Hume, 1878) | Hume's Whitethroat | Least Concern | Winter Visitor | Stable |
| 111 | Chrysomma sinense (Gmelin, 1789) | Yellow-eyed Babbler | Least concern | Resident | Stable |
| 112 | Sylvia curruca (Linnaeus, 1758) | Common Lesser Whitethroat | Least concern | Winter visitor | Stable |
| | Tephrodornithidae | | | | |
| 113 | Tephrodornis pondicerianus, (Gmelin, 1789) | Common Woodshrike | Least concern | Resident | Stable |
| | Tytonidae | | | | |
| 114 | Tyto alba (Scopoli, 1769) | Barn Owl | Least concern | Resident | Decreasing |
| | Upupidae | | | | |
| 115 | Upupa epops (Linnaeus, 1758) | Common Hoopoe | Least concern | Winter visitor | Stable |
| | Zosteropidae | | | | |
| 116 | Zosterops palpebrosus (Temminck, 1824) | Oriental White-eye | Least concern | Resident | Decreasing |

The category of water birds included 60 species of birds belonging to different families. Out of which 05 species Aythya nyroca, Mycteria leucocephala, Limosa limosa, Sterna aurantia and Threskiornis melanocephalus are Near Threatened, 02 species Aythya ferina and Pelecanus crispus are Vulnerable, 50 are Least Concern and 03 species "taxon has not yet been assessed for the IUCN Red List, but it is in Catalogue of Life" (Table 3).

| | | | - | |
|-------|-----------------|-------------|-------------------|-------------------|
| S.No. | IUCN Status | Water birds | Terrestrial birds | Number of species |
| 1 | Endangered | - | 02 | 02 |
| 2 | Vulnerable | 02 | - | 02 |
| 3 | Near Threatened | 05 | 01 | 06 |
| 4 | Least Concern | 50 | 108 | 158 |
| 5 | Not Evaluated | 03 | 05 | 08 |
| | Total | 60 | 116 | 176 |

According to IUCN Status, 34 species were Winter Visitor and 01 Passage Visitor and 25 were Resident (Table 4). Population trend of 29 species decreased, 20 were Stable and 07 species increased in the study area (Table 5). Population trend of 04 species were Unknown.

| S. No. | Migratory status | Water birds | Terrestrial bird | Number of Speciec |
|--------|------------------|-------------|------------------|-------------------|
| 1 | Resident | 25 | 86 | 111 |
| 2 | Winter Visitor | 34 | 25 | 059 |
| 3 | Summer Visitor | - | 04 | 04 |
| 4 | Passage Visitor | 01 | 01 | 02 |
| Total | | 60 | 116 | 176 |



International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.177

Volume 7 Issue VI, June 2019- Available at www.ijraset.com

| Table 5: Population | trend of avian | species |
|---------------------|----------------|---------|
|---------------------|----------------|---------|

| S. No. | Population trend | Water birds | Terrestrial bird | Number of Species |
|--------|------------------|-------------|------------------|-------------------|
| 1 | Increases | 07 | 12 | 019 |
| 2 | Stable | 20 | 63 | 083 |
| 3 | Decreases | 29 | 34 | 063 |
| 4 | Unknown | 04 | 07 | 011 |
| Total | | 60 | 116 | 176 |

In the category of terrestrial birds 116 species beloning to different families were observed in the study area. Out of which 02 species Aquila nipalensis and Neophron percoopterus are endangered and only one species Psittacula eupatria is near threatened. Five species were not yet been assessed for the IUCN Red List, but is in the Catalogue of Life. 108 species were least concern, 86 birds species are Resident. 25 species are Winter Visitor, 04 species Clamator jacobinus, Merops persicus, Terpsiphone paradise and Oriolus oriolus are Summer Visitor and only 01 species Coracias garrulous is Passage Visitor.

Population trend of 34 species decreases, 63 are stable and 12 species increases where as 7 species were unknown. Major threats and reason for decline in population are mining, anthropogenic activities, population growth, resource consumption, climate change and global warming, urbanisation, invasive alien species, over-exploitation of natural resources and environmental degradation.

IV. ACKNOWLEDGEMENT

We would like to express our deep gratitude and thanks to Prof. K.K. Sharma, Former Vice-Chancellor, Maharshi Dayanand Saraswati University, Ajmer (Rajasthan), for his motivational spirit. I extend my sincere thanks to Dr. Mahesh Bundele, Dean (R&D), Poornima University, Jaipur. I am highly indebted to Dr. S.P. Mehra for his timely guidance and support. I am also thankful to Department of Forest, Government of Rajasthan.

REFERENCES

- [1] Bibby, C.J., Hill, D.A. & Burgess, N.D. (2000). Bird Census Techniques. 2nd edn. Academic Press, London.
- [2] Javed, S. & Kaul, R. (2002). Field methods for bird surveys. Bombay Natural History Society; Department of Wildlife Sciences, Aligarh Muslim University, Aligarh and World Pheasant Association, South Asia Regional Office (SARO), New Delhi, India.
- [3] Ali, S. (1941). The book of Indian birds. Bombay: Bombay Natural History Society.
- [4] Ali, S. (1942). The book of Indian birds. Second edition. Bombay: Bombay Natural History Society.
- [5] Ali, S. (1945). The birds of Kutch. Bombay: Oxford University Press.
- [6] Ali, S. (1946). An ornithological pilgrimage to Lake Manasarowar and Mount Kailas. Bombay Natural History Society. 46: 284–309
- [7] Ali, S. (1977). President's letter: "mystery" birds of India: Mountain Quail. Hornbill. 3: 3-5.
- [8] Grimmett, R., Inskipp, C. & Inskipp, T. (1998). Birds of the Indian subcontinent. London: A. & C. Black/Christopher Helm.











45.98



IMPACT FACTOR: 7.129







INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 🕓 (24*7 Support on Whatsapp)