



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 7 Issue: VIII Month of publication: August 2019

DOI: <http://doi.org/10.22214/ijraset.2019.8136>

www.ijraset.com

Call: ☎ 08813907089

E-mail ID: ijraset@gmail.com

An Advance Class of Analytic Functions with Fekete-Szegő Inequality using subordination Method

S. K. Gandhi¹, Gurmeet Singh², Preeti Kumawat³, G. S. Rathore⁴, Lokendra Kumawat⁵

^{1, 3, 4, 5}Department of Mathematics and Statistics, Mohanlal Sukhadia University, Udaipur (Raj) 313001

²Department of Mathematics, GSSDGS Khalsa College, Patiala, India

Abstract: In this Paper we have introduced an advance class of analytic functions along with its subclasses by using principle of subordination and as so obtained sharp upper Bound of the function $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ belonging to these classes.

Extremal functions are also investigated.

Keywords: Bounded functions, Close to convex function, extremal function, Inverse Starlike functions, Starlike functions, Univalent functions.

I. INTRODUCTION

Let \mathcal{A} denote the class of analytic function

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad (1.1)$$

In the unit disc $= \{z: |z| < 1\}$, \mathcal{S} be the class of analytic univalent functions in \mathbb{E} .

Bieber Bach [7] proved that $|a_2| \leq 2$ for the functions $f(z) \in \mathcal{S}$. and Löwner [5] proved that $|a_3| \leq 3$ for the functions $f(z) \in \mathcal{S}$. With the above known estimates this inequality plays an important role to determining estimates of higher coefficients for some sub classes \mathcal{S} {Chhichra [11], Babalola [6]}

Using Löwner's method [5] In 1933, Fekete and szego investigated a well known relation between a_3 and a_2^2 for the class \mathcal{S}

$$|a_3 - \mu a_2^2| \leq \begin{cases} 3 - 4\mu & , \text{if } \mu \leq 0 \\ 1 + 2e^{\frac{-2\mu}{1-\mu}} & , \text{if } 0 \leq \mu \leq 1 \\ 4\mu - 3 & , \text{if } \mu \geq 1 \end{cases} \quad (1.2)$$

Let us define some subclasses of \mathcal{S}

Let \mathcal{K} denotes subclasses of \mathcal{S} of univalent convex functions $h(z) = z + \sum_{n=2}^{\infty} c_n z^n \in \mathcal{A}$ satisfying the condition

$$\operatorname{Re} \frac{(zh'(z))}{h'(z)} > 0, z \in \mathbb{E}. \quad (1.3)$$

A function $f(z) \in \mathcal{A}$ is said to be close to convex if there exist $g(z) \in \mathcal{S}^*$ such that

$$\operatorname{Re} \frac{(zf'(z))}{g(z)} > 0, z \in \mathbb{E}. \quad (1.4)$$

The class of close to convex functions introduced by Kaplan [17], and he proved that close to convex functions are univalent.

$$\mathcal{S}^*(A, B) = \{ f(z) \in \mathcal{A} ; \frac{(zf'(z))}{g(z)} < \frac{1+Az}{1+Bz}, -1 \leq B \leq A \leq 1, z \in \mathbb{E} \} \quad (1.5)$$

Where $\mathcal{S}^*(A, B)$ is a subclass of \mathcal{S}^* .

For strongly alpha quasi-convex functions Fekete-Szegő problem was studied by Abdel-Gawad [3]. The upper bound of $|a_3 - \mu a_2^2|$ for different functions in the class \mathcal{S} has been investigated by many authors including Goel and Mehrotra [13] and recently by Al-Shaqsi and Darus [4] Hayami and Owa [16], Al-Abbadi and Darus [10].

Gurmeet singh et al. [3] introduced the class of inverse Starlike functions

$g(z) = z + \sum_{n=2}^{\infty} b_n z^n \in \mathcal{A}$ satisfying the condition

$$\operatorname{Re} \left(\frac{zf(z)}{2 \int_0^z f(z) dz} \right) > 0, z \in \mathbb{E} \quad \text{i.e.} \quad \frac{zf(z)}{2 \int_0^z f(z) dz} < \frac{1+z}{1-z} \quad (1.6)$$

Gandhi et al. [14] established a new class of analytic functions with Fekete-szego inequality using subordination method.

Here introduce the class \mathcal{A} , of Univalent starlike functions $g(z) = z + \sum_{n=2}^{\infty} b_n z^n \in \mathcal{A}$ satisfying the condition

$$\left[\frac{z\{zf(z)\}'}{2f(z)} \right] < \left(\frac{1+z}{1-z} \right)^\alpha ; \alpha > 0 \quad (1.7)$$

And subclass consisting of the functions $g(z) = z + \sum_{n=2}^{\infty} b_n z^n \in \mathcal{A}$ satisfying the condition

$$\left[\frac{z\{zf(z)\}'}{2f(z)} \right] < \left(\frac{1+Az}{1+Bz} \right)^\alpha ; -1 \leq B \leq A \leq 1 ; \alpha > 0 \quad (1.8)$$

Here, Symbol $<$ stands for subordination, defined as follows:

A. Principle of Subordination

If $f(z)$ and $F(z)$ are two functions which are analytic in \mathbb{E} , then $f(z)$ is called a subordinate to $F(z)$ in \mathbb{E} , if there exists a function $w(z)$ which is analytic in \mathbb{E} satisfying the conditions

$$(i) w(0) = 0 \quad \text{and} \quad (ii) |w(z)| < 1$$

such that $f(z) = F(w(z))$, where $z \in \mathbb{E}$ and we denote it as $f(z) < F(z)$.

Let \mathcal{U} denote the class of analytic bounded functions of the form

$$w(z) = \sum_{n=1}^{\infty} d_n z^n, w(0) = 0, |w(z)| < 1 \quad (1.9)$$

With $|d_1| \leq 1, |d_2| \leq 1 - |d_1|^2$.

II. RESULTS AND DISCUSSION

1) *Theorem 1:* If $f(z) \in \mathcal{A}$, then the result

$$|a_3 - \mu a_2^2| \leq \begin{cases} 10\alpha^2 - 16\mu\alpha^2 & , \text{if } \mu \leq \frac{5\alpha-1}{8\alpha} \end{cases} \quad (1.10)$$

$$2\alpha \quad , \text{if } \frac{5\alpha-1}{8\alpha} \leq \mu \leq \frac{5\alpha+1}{8\alpha} \quad (1.11)$$

$$16\mu\alpha^2 - 10\alpha^2 \quad , \text{if } \mu \geq \frac{5\alpha+1}{8\alpha} \quad (1.12)$$

is sharp.

Proof: By using expansion method (1.7) leads to

$$1 + \frac{1}{2} a_2 z + (a_3 - \frac{1}{2} a_2^2) z^2 + \dots = 1 + 2\alpha c_1 z + 2\alpha(c_2 + \alpha c_1^2) z^2 + \dots \quad (1.13)$$

After Identifying the terms we have

$$|a_3 - \mu a_2^2| \leq |2\alpha c_2 + 10\alpha^2 c_1^2 - 16\mu\alpha^2 c_1^2|$$

This leads to

$$|a_3 - \mu a_2^2| \leq 2\alpha + [10\alpha^2 - 16\mu\alpha^2 - 2\alpha] |c_1|^2 \quad (1.14)$$

Case I: If $\mu \leq \frac{5}{8}$, then (1.14) leads to

$$|a_3 - \mu a_2^2| \leq 2\alpha + [10\alpha^2 - 2\alpha - 16\mu\alpha^2] |c_1|^2 \quad (1.15)$$

Subcase I(a): If $\mu \leq \frac{5\alpha-1}{8\alpha}$, then (1.15) leads to

$$|a_3 - \mu a_2^2| \leq 10\alpha^2 - 16\mu\alpha^2 \quad (1.16)$$

Subcase I(b): If $\mu \geq \frac{5\alpha-1}{8\alpha}$, then (1.15) leads to

$$|a_3 - \mu a_2^2| \leq 2\alpha \quad (1.17)$$

Case II: If $\mu \geq \frac{5}{8}$, then (1.14) leads to

$$|a_3 - \mu a_2^2| \leq 2\alpha + [16\mu\alpha^2 - (10\alpha^2 + 2\alpha)] |c_1|^2 \quad (1.18)$$

Subcase II(a): If $\mu \leq \frac{5\alpha+1}{8\alpha}$, then (1.18) leads to

$$|a_3 - \mu a_2^2| \leq 2\alpha \quad (1.19)$$

Subcase II(b): If $\mu \geq \frac{5\alpha+1}{8\alpha}$, then (1.18) leads to

$$|a_3 - \mu a_2^2| \leq 16\mu\alpha^2 - 10\alpha^2 \quad (1.20)$$

Combining subcase II(a) and subcase I(b), we get

$$|a_3 - \mu a_2^2| \leq 2\alpha \quad , \quad \text{if } \frac{5\alpha-1}{8\alpha} \leq \mu \leq \frac{5\alpha+1}{8\alpha} \quad (1.21)$$

This completes the theorem, therefore the result is sharp.

Extremal function for the first and third inequality is given by

$$f_1(z) = \frac{z}{(1-\alpha z)^4} \quad (1.22a)$$

And Extremal function for the second inequality is given by

$$f_2(z) = \frac{z}{(1-\alpha z^2)^2} \quad (1.22b)$$

2) *Theorem 2 :* If $f(z) \in \mathcal{A}$, then the result

$$|a_3 - \mu a_2^2| \leq$$

$$\begin{cases} (A-B)(2A-3B)\alpha^2 - 4\mu\alpha^2(A-B)^2 & , \text{if } \mu \leq \frac{(2A-3B)\alpha-1}{4(A-B)\alpha} \end{cases} \quad (1.23a)$$

$$\begin{cases} (A-B)\alpha & , \text{if } \frac{(2A-3B)\alpha-1}{4(A-B)\alpha} \leq \mu \leq \frac{(2A-3B)\alpha+1}{4(A-B)\alpha} \end{cases} \quad (1.23b)$$

$$\begin{cases} 4\mu\alpha^2(A-B)^2 - (A-B)(2A-3B)\alpha^2 & , \text{if } \mu \geq \frac{(2A-3B)\alpha+1}{4(A-B)\alpha} \end{cases} \quad (1.23c)$$

is sharp.

Proof: By using expansion method (1.8) leads to

$$1 + \frac{1}{2}a_2 z + (a_3 - \frac{1}{2}a_2^2) z^2 + \dots = 1 + (A-B)\alpha c_1 z + (A-B)\alpha(c_2 - B\alpha c_1^2) z^2 + \dots \quad (1.24)$$

After Identifying the terms in (1.24) we have

$$|a_3 - \mu a_2^2| \leq |(A-B)\alpha(c_2 - B\alpha c_1^2) + 2(A-B)^2\alpha^2 c_1^2 - 4\mu(A-B)^2\alpha^2 c_1^2|$$

This leads to

$$|a_3 - \mu a_2^2| \leq (A-B)\alpha + \{2(A-B)^2\alpha^2 - B(A-B)\alpha^2 - 4\mu(A-B)^2\alpha^2 - (A-B)\alpha\} |c_1|^2 \quad (1.25)$$

here two cases arise:

Case I: If $\mu \leq \frac{2A-3B}{4(A-B)}$, then (1.25) leads to

$$|a_3 - \mu a_2^2| \leq (A-B)\alpha + [(A-B)\{(2A-3B)\alpha-1\}\alpha - 4\mu(A-B)^2\alpha^2] |c_1|^2 \quad (1.26)$$

Under this case (1.26) two subcases arise:

Subcase I(a): If $\mu \leq \frac{(2A-3B)\alpha-1}{4(A-B)\alpha}$, then (1.26) leads to

$$|a_3 - \mu a_2^2| \leq \{(A-B)(2A-3B)\alpha - 4\mu(A-B)^2\alpha^2\} |c_1|^2 \quad (1.27)$$

Subcase I(b): If $\mu \geq \frac{(2A-3B)\alpha-1}{4(A-B)\alpha}$, then (1.26) leads to

$$|a_3 - \mu a_2^2| \leq (A-B)\alpha \quad (1.28)$$

Case II: If $\mu \geq \frac{2A-3B}{4(A-B)}$, then (1.25) leads to

$$|a_3 - \mu a_2^2| \leq (A-B) + [4\mu(A-B)^2\alpha^2 - (A-B)\{(2A-3B)\alpha+1\}\alpha] |c_1|^2 \quad (1.29)$$

Under this case again two subcases arise:

Subcase II(a): If $\mu \leq \frac{(2A-3B)\alpha+1}{4(A-B)\alpha}$, then (1.29) leads to

$$|a_3 - \mu a_2^2| \leq (A-B)\alpha \quad (1.30)$$

Subcase II(B): $\mu \geq \frac{(2A-3B)\alpha+1}{4(A-B)\alpha}$, then (1.29) leads to

$$|a_3 - \mu a_2^2| \leq \{4\mu(A-B)^2\alpha^2 - (A-B)(2A-3B)\alpha^2\} |c_1|^2 \quad (1.31)$$

Combining subcase II(a) and subcase I(b), we get

$$|a_3 - \mu a_2^2| \leq (A-B)\alpha, \text{ if } \frac{(2A-3B)\alpha-1}{4(A-B)\alpha} \leq \mu \leq \frac{(2A-3B)\alpha+1}{4(A-B)\alpha} \quad (1.32)$$

This completes the theorem therefore, the result is sharp.

Extremal function for the first and third inequality is given by

$$f_1(z) = z \left\{ 1 + \left(\frac{2A-2B}{2A-3B} \right) z\alpha \right\}^{2A-3B} \quad (1.33a)$$

And for the for second inequality is given by

$$f_2(z) = \frac{z}{(1-\alpha z^2)^{A-B}} \quad (1.33b)$$

III. CONCLUDING REMARKS AND COROLLARIES

1) *Corollary 1.1:* Taking $\alpha = 1$ in the theorem 1, we get

$$|a_3 - \mu a_2^2| \leq \begin{cases} 10 - 16\mu & , \text{if } \mu \leq \frac{1}{2} \end{cases} \quad (1.34)$$

$$|a_3 - \mu a_2^2| \leq \begin{cases} 2 & , \text{if } \frac{1}{2} \leq \mu \leq \frac{3}{4} \end{cases} \quad (1.35)$$

$$|a_3 - \mu a_2^2| \leq \begin{cases} 16\mu - 10 & , \text{if } \mu \geq \frac{3}{4} \end{cases} \quad (1.36)$$

These estimates were derived by Keogh and Merkes [1] and the results are for the class of univalent convex functions.

Further if we take $A = 1$ and $B = -1$ ($-1 \leq B \leq A \leq 1$) in the result of theorem 2, we get the result of theorem 1, therefore our result for the theorem 2 reduces to the result of the theorem 1. Hence theorem 2 is the generalization of theorem 1. And the results are sharp and also if we put $A = 1$ and $B = -1$ in extremal function of theorem 2, we get the extremal function of theorem 1. The extremal function given by [(1.22a),(1.22b)] increases as α increases and decreases as α decreases and the extremal function given by [(1.33a),(1.33b)] also increases and decreases as α increases and decreases respectively. Hence extremal function is an increasing function.

REFERENCES

- [1] F. R. Keogh, E. R. Merkes, "A coefficient inequality for certain classes of analytic functions", *Proceedings of American Mathematical Society*, 20, 8 (1989)
- [2] Gurmeet Singh, M. S. Saroa and B. S. Mehrotra, "Fekete-szegő inequality for a new class of analytic functions", *Elsevier, Proc. of International conference on Information and Mathematical Sciences*, 90 (2013)
- [3] H. R. Abdel-Gawad, D. K. Thomas, "The Fekete-Szegő problem for strongly close-to-convex functions" *Proceedings of the American Mathematical Society*, vol. 144, no.2 345(1992)
- [4] K. Al-Shaqsi, M. Darus, "On Fekete-Szegő problems for certain subclass of analytic functions", *Applied Mathematical Sciences*, vol. 2 no. 9-12, pp. 431(2008)
- [5] K. Löwner, "Über monotone Matrixfunktionen", *Math. Z* 38, 177 (1934).
- [6] Kunle Oladeji Babalola, "The fifth and Sixth coefficient of α -close-to-convex function", *Kragujevac J. Math.* 32, 5 (2009)
- [7] L. Bieberbach, "Über einige Extremal Probleme im Gebiete der Konformen Abbildung" *Math.* 77, 153 (1916)
- [8] L. Bieberbach, "Über die Koeffizienten derjenigen Potenzreihen, welche eine Schlichte Abbildung des Einheitskreises vermitteln", *Preuss. Akad. Wiss. Sitzungsber.* 940 (1916)
- [9] M. Fekete, G. Szegő, "Eine Bemerkung über ungerade schlichte Funktionen", *J. London Math. Soc.* 8, 85 (1933)
- [10] M. H. Al-Abbadi, M. Darus, "Fekete-Szegő theorem for a certain class of analytic functions", *Sanis Malaysiana*, vol. 40, no. 4, 385(2011)
- [11] P. N. Chhichra, "New Subclasses of the class of close to convex functions", *Proceedings of American Mathematical Society*, 62, 37 (1977)
- [12] R. M. Goel, B. S. Mehrotra, "A Subclass of Univalent functions", *Houston Journal of Mathematics* 8, 343 (1982)
- [13] R. M. Goel, B. S. Mehrotra, "A coefficient inequality for certain classes of analytic function", *Tamkang Journal of Mathematics* 22, 153 (1990)
- [14] S. K. Gandhi, Gurmeet Singh, Preeti Kumawat, G. S. Rathore and Lokendra Kumawat, "A new class of analytic functions with Fekete-Szegő Inequality using subordination Method", *International journal of research in advanced technology*, vol. 6, no. 9, (2018)
- [15] S. R. Keogh, E. R. Merkes, "A coefficient inequality for certain classes of analytic functions", *Proceedings of the American Mathematical Society*. Vol. 20, 8 (1969)
- [16] T. Hayami, S. Owa and H. M. Srivastava, "Coefficient inequalities for certain classes of analytic and univalent functions", *J. Ineq. Pure and Appl. Math.* 8(4), 1 (2007)
- [17] W. Kaplan, "Close-to-convex schlicht functions", *Michigan Mathematical Journal* 1, 169 (1952)



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)