



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 7 Issue: XI Month of publication: November 2019

DOI: <http://doi.org/10.22214/ijraset.2019.11105>

www.ijraset.com

Call: ☎ 08813907089

E-mail ID: ijraset@gmail.com

Analysis of Different Method of Solving Quadratic Equations

Vishal V. Mehtre¹, Ajit Kumar Mishra²

¹Assistant Professor, ²Student, Bharati Vidyapeeth Deemed (to be) University, College of Engineering, Pune, Maharashtra, India.

Abstract: In this research paper we are going to detail about how to calculate approximate value of roots of the polynomial equation by different ways. We are going to study three methods i.e, equation by Bisection, Regular-Falsi and Newton-Raphson methods and how these methods differ from each other. There are some different method to solve the same problem but the question arises that which method is better, more time efficient and more accurate. These methods are evolving and improved on regular basis. Less time consumption and easiness of solving equation can be brought by changing variable and by some small change in calculation. At the conclusion we will elaborate on the differences in solving a particular polynomial equation by using variety of method, what are the differences among these methods and difficulty level of them. We will get to know how these methods differ from each other

I. INTRODUCTION

The fact that the roots of the polynomial can be obtained immediately using computer program as MATLAB, does not diminish the importance of gaining the new ways for solving the polynomial equation, simpler than that of current ones. In this paper we are going to get detail information about the different methods such as Bisection, Secant and Newton-Raphson methods and the difference between the ways of solving and the difference between each of them. The root finding problem is one of the most relevant computational problem. It arises in a wide variety of practical application in Physics, Chemistry, Biosciences, etc. Different methods converge the roots at different rates.[2] That is Some methods are faster in converging to the roots than others. The rate of convergence could be linear, quadratic or otherwise. The higher the order, the faster the method converges. The whole study is comparing the rate of performance of Bisection, and Newton-Raphson methods of finding roots. It can be seen that Newton-Raphson may converge faster than any other method but when we compare the performance, it is needful to consider both cost and speed of convergence. An algorithm that converges quickly but takes a few second per iteration may take more time overall than an algorithm that converges more slowly, but takes a few milliseconds per iteration. As Secant method and Newton-Raphson method are almost same, from geometric perspective. The difference is that Newton-Raphson method uses a line that is tangent to one point, while the Scent method uses a line that is secant to two points. In Newton-Raphson, the derivation of a function at a point is used to create the tangent line, whereas in Secant method, a numerical approximation of the derivative based on two points is used to create the secant line. In comparing the rate of convergence if Bisection, Secant and Newton-Raphson methods used C++ program language to calculate cube root of number from 1 to 25, using the three methods. They observed that rate of convergence is in the following order: Bisection Methods < Newton-Raphson <

Secant Method. They conclude that Newton-Raphson method is 7.6786 times better than the bisection method while Secant method is 1.3894 times better than Newton method.[5]

II. METHODS TO OBTAIN ROOTS OF EQUATION

Given a function $f(x)=0$, continuous on a closed interval $[a_1, b_1]$, such that $f(a_1).f(b_1)<0$

A. Bisection Method




Fig 2.1: Graphical representation of Bisection method

The essential condition for bisection method is $f(a).f(b) < 0$. we find two values in the given interval such that the multiplication of them should be less than zero. After when we have to find the values of “a” and “b” we take the out the value of $c = (a+b)/2$ and we put this value of c in the original equation.[1] Let us take an example to understand the concept-

$F(x) = x^3 - 2x - 5$ (This is the given polynomial equation and we have to find the roots of the equation)

$F(1) = 1-2-5=-6$, $f(2) = 8-4-5=-1$, $f(3)=27-6-5=16$. Hence we can see that the $f(1). f(3)<0$ and the condition get satisfied. So the roots will lie between [2,3]. Therefore 1st iteration $a+b$ 2+3

$$c = \frac{2}{2} + \frac{3}{2} = 2.5 \text{ then we have to find the}$$

value of $f(2.5)=5.625 > 0$

$F(2). f(2.5) < 0$ this condition again get full filled again we have to find the value of

2nd iteration

$c = \frac{2+2.5}{2} = 2.25$ and again we find the value of $f(2.25)=1.890 > 0$ this process continues this until we find the value of c which is same after 3

place of decimal

Table 2.1: Result of Bisection Method

Number of iteration	Value of a	Value of b	$C = \frac{a + b}{2}$	$F(c) =$
1st	2	3	2.5	5.625
2nd	2.5	2	2.25	1.890
3rd	2.25	2	2.125	0.34
4th	2.125	2	2.0625	-0.3513
5th	2.0625	2.125	2.09375	-0.0089
6th	2.09375	2.125	2.109375	0.166835
7th	2.09375	2.109	2.101562	0.078562
8th	2.10156	2.093	2.09765	0.03464
9th	2.09765	2.093	2.0957	0.012827
10th	2.0957	2.093	2.0957	0.001936

Hence, we find the value till we don't get the value of c till 3 places of decimal and hence we get it and the 10th iteration.

B. Regula-falsi Method




Fig 2.2: Graphical Representation of Regula-Falsi Method

Regula-Falsi method is very similar to secant method. In secant method the approximate function $f(x)$ by a straight line. The point at which the line crosses x (ax1) is called approximate value of that function.[3] If $x(0)$ & $x(1)$ are the initial approximation then approximate line to the function $f(x)$ passes through point $[X0, f(X0)]$ & $[X1, f(X1)]$ If approximation are select in such a way that if $f(x) < 0$ then secant method is called as RegulaFalsi method. $f(X) \cdot f(X_{n+1}) < 0$ crosses zero at certain point hence method converges fast.

$$X_{n+1} = X_n - \frac{X_n - X_{n-1}}{f(X_n) - f(X_{n-1})} * f(X_n)$$

let us understand this method by an example- $F(x) = x^3 - 2x - 5$ (This is the given polynomial equation and we have to find the roots of the equation)

$$F(0) = -5, f(1) = -6, f(2) = -1, f(3) = 16$$

Hence $f(2) \cdot f(3) < 0$, $X_0 = 2$ & $X_1 = 3$ and $f(X_0) = -1$ & $f(X_1) = 16$ 1st iteration

For $n=1$

$$X_2 = X_0 - \frac{X_1 - X_0}{f(X_1) - f(X_0)} * f(X_0)$$

$$= 2.058823$$

$$F(X_2) = -0.390805$$

2nd iteration For $n=2$

$$X_3 = X_2 - \frac{X_2 - X_1}{f(X_2) - f(X_1)} * f(X_2)$$

$$= 2.081263423$$

$$F(X_3) = -0.147244$$

Table 2.2: Result of Regula-Falsi Method

Num ber of itera tion	Value of X_0	Value of X_1	$X_{n+1} = X_n - \frac{X_n - X_{n-1}}{f(X_n) - f(X_{n-1})} * f(X_n)$	$F(X_{n+1})$
1st	2	3	2.058823	-0.390 805
2nd	2.058823	3	2.081263423	-0.205 5
3rd	2.081263423	2.058823	2.09483784	0.00277
4th	2.09483784	2.081263423	2.094579431	2.06*10-4

As we can see that value of 4th iteration and the value of 5th iteration are matching after 4 place of decimal point.

C. Newton-Raphson Method

Let x_0 be the initial root of the equation. X

$$X_1 = X_0 + h \text{-----(1)}$$

Using Taylor series for equation $F(x_1) = f(x_0) + h \cdot f'(x_0) + \frac{h^2}{2} \cdot f''(x_0) + \dots$. Therefore, h is small so neglect higher order terms.

$$F(X_1) = f(X_0) + h \cdot f'(X_0) = 0 \Rightarrow h = -f(X_0)/f'(X_0)$$

substitute value of h in equation (1)

$$X_1 = X_0 - \frac{f(X_0)}{f'(X_0)}$$

In general, the formula is:

$$X_{m+1} = X_m - \frac{f(X_m)}{f'(X_m)}$$

(*) To check approximate is correct or not—

(1) If $f'(X_0) = 0$ change initial value

(2) Use the condition $f(X) \cdot f'(X) > 0$

Let us understand this with an example...

$$F(x) = x^3 - 2x - 5$$

$$F'(x) = 3x^2 - 2$$

$$F(1) = -6, F'(1) = 1$$

$$F(2) = -1, F'(2) = 10$$

$$F(3) = 16, F'(3) = 25$$

$$H = -16/25 = 0.64$$

1st iteration

$$X_0 = 3$$

$$X_1 = 3 + 0.64 = 2.36$$

$$X_2 = 2.36 - \frac{3.4242}{14.7088} = 2.127200$$

Table 2.3: Result of Newton-Raphson Method

Number of iteration	Values of X	Value came out from formula
1st	X_1	2.36
2nd	X_2	2.127200
3rd	X_3	2.095136
4th	X_4	2.09455167
5th	X_5	2.094551482

As we can see that value of 4th iteration and the value of 5th iteration are matching after 4 place of decimal point.

III. CONCLUSION

We can conclude on the basis of detailed analysis carried out through this paper is that Newton Raphson method and Regula-Falsi method are easy to use and implement for finding roots of any equation as compared to Bisection method. In Bisection method we need to carry the calculations up to 10th iteration which is troublesome while contrastingly, In Newton Raphson method and RegulaFalsi method we will get the desired result in the 5th iteration. Hence, we can conclude that Bisection method is more prone and fallible to error since the convergence is slow and number of iterations is more. In Newton – Raphson, as we calculate the derivative of the given function, the calculation is converged at faster rate with fewer number of iterations. With help of Regula–Falsi method the result obtained are same like Newton – Raphson method, which may change with the function used. Depending on the function used the number of iterations and convergence rate changes. We have also concluded that due to fast convergence rate to find the roots of any equation, Newton – Raphson method is most widely used.



REFERENCES

- [1] P.Deuffhard, "Newton Methods for Nonlinear Problem and Adaptive Algorithms", Springer Series in Computational Mathematics, Vol. 35. Springer, Berlin, 2004
- [2] A family of regula falsi root-finding methods by Sérgio Galdino, Polytechnic School of Pernambuco University ,
- [3] "Comparative study of Bisection method, Newton-Raphson and Regula-Falsi method" "Ehiwario, J.C., Aghamie, S.O, ISOR Journal of Engineering, ISSN (e): 2250-3021, Vol. 04, Issue 04, April 2014.
- [4] Numerical Solution of Root-Finding Problems by Professor Biswa Nath Datta, Department of Mathematical Sciences Northern Illinois University



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)