



IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 8 Issue: IV Month of publication: April 2020

DOI: http://doi.org/10.22214/ijraset.2020.4079

www.ijraset.com

Call: 🕥 08813907089 🔰 E-mail ID: ijraset@gmail.com



Opportunistic Invasive Fungal Diseases: Types, Symptoms, Treatment, and Laboratory Diagnosis

Saqib Bashir¹, Aasia Sofi², Dr. Hanan³, Dr. Jaya Bharti⁴

^{1, 2} MSc. Student, Department of Medical Laboratory Technology, Mewar University, Chittorgarh, Rajasthan, India ^{3, 4} Assistant Professor, Department of Paramedical Sciences, Mewar University, Chittorgarh, Rajasthan, India

Abstract: Invasive fungal infections (opportunistic) are significant wellbeing complications in immuno compromised patients and in terminal stages of chronic diseases. The signs and manifestations differ and can range from colonization in hypersensitive bronchopulmonary sickness to dynamic infection by local infective agents. Many factors impact the seriousness and infection causing capacity of the fungus, for example, chemicals, dimorphic development in some Candida species, melanin creation and poison creation. Infection is affirmed when histopathology study is finished with the assistance of special stains shows fungal tissue involvement or when the infective agent is gotten from clinical examples by culture. Both obtained and intrinsic immunodeficiency may be related with increased weakness to systemic infections. Fungal infection can't be treated because antifungal therapy for Candidiasisis still controversial and for molds, the clinician must have to affirm that the species acquired from the culture medium is the pathogen. Convenient inception of antifungal treatment is a basic segment influencing the results. The general spreading fungal infection requires the utilization of foundational operators with or without surgical activities, and in some cases immunotherapy is additionally used. Preclinical and clinical examinations show a decent connection between medicate portion and treatment result. Medication portion movement is essential to ensure that therapeutic levels are accomplished for ideal clinical results. The goal of this audit is to examine opportunistic fungal infections, diagnostic methods and the administration of these diseases.

Keywords: Fungal disease, Medication, Laboratory methods, Opportunistic fungal diseases.

I. INTRODUCTION

Some saprophytic fungus do not cause malady yet they can cause infection under some significant conditions for example, immunocompromised persons and in terminal stages of chronic disease. These are called opportunistic fungi and the infection caused by them is called invasive fungal infections (IFI).

Invasive fungal infections (IFI) (opportunistic) are spreading rapidly because due to advances in clinical consideration, widespread utilization of anti-microbials, corticosteroids and due to immunosuppressive medications. These types of contagious species are present in soil, plant debris and other natural substances, and make up roughly 7 per cent (611,000 species) of every single eukaryotic species on earth, out of which only about 600 species are human pathogen. [1]

The variables responsible for invasive fungal infections (IFI) include patients with neutropenia, hematological malignancies, bone marrow transplantation, patients taking wide –range anti-toxins, and delayed utilization of corticosteroids for treatment and chemotherapy, patients with HIV disease, invasive clinical procedures, and because of resistant immune suppressive agents. Other risk factors are diabetes mellitus, lack of healthy nutrition, solid organ transplantation and significant medical procedure.[2-3]There are also reports of the presence of infection in Immunocompetent patients[4]without signs or side effects of conditions related with immunocompromised status.

Disease can be transmitted by the inward breath of spores, percutaneous immunization in cutaneous and subcutaneous diseases, entrance into the mucosa by commensal living beings for example, *Candida albicans*, and the ingestion of a poison in contaminated nourishment or drink (gastrointestinal disease).[5]

II. CANDIDIASIS

The causative agent of this fungal infection is candida albicans (80-90% of cases). It is yeast which live in small amounts in our mouth and belly or on skin. Candidiasis is a disease of skin, mucosa and also affects internal organs. [6]

A. Thrush (Oropharyngal Candidiasis)

Type of candidiasis when the yeast spreads to mouth and throat and can cause an infection known as thrush. This infection can affect mostly newborns, old age groups and persons with weak immune system. And also can infect adults who are being treated with cancer, taking medications like corticosteroids and broad- spectrum anti-toxins and also infect diabetic patients. [7]



International Journal for Research in Applied Science & Engineering Technology (IJRASET)

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429 Volume 8 Issue IV Apr 2020- Available at www.ijraset.com

- 1) Symptoms:
- a) White or yellow patches on tongue, lips, gums and in inner cheeks (Fig: 1.1)
- b) Rush in mouth and in throat (Fig: 1.2)
- c) Mouth cracking on corners (Fig: 1.3)
- d) Pain during swallowing (if spreads to throat)
- 2) *Treatment:* Thrush can be treated with antifungal medicines like clotrimazol, fluconazole, itraconazole and amphoteric B and washing of mouth with chlrohexidine (CXN) can prevent the infection in persons with weak immune system. [8]





Fig: 1.2



Fig: 1.3

B. Genital Yeast Infection (Genital Candidiasis)

These are the candida species that can cause infection in the genital tract of women the vagina and in men the penis will be affected with this infection .This happens when too much yeast grows in vagina and is more common in women's as compared to men. [9] This occurs typically when the vaginal balance changes, which is caused by pregnancy, diabetes and due to use of spermicides.

- 1) Symptoms:
- a) Vaginal itching
- b) Pain and burning during urination
- c) Pain during sexual intercourse
- d) Swelling of vagina and vulva [10]
- A man with this infection has a itching on his penis.
- 2) Treatment: The antifungal tablet or cream will cure the infection. Fluconazole an antifungal medicine can be prescribed.

C. Invasive Candidiasis

This type of infection is caused when the yeast spreads to bloodstream by medical equipment's or devices. This infection affects heart, brain, blood, eyes and also bones. The risk factors are weak immune system, kidney failure and use of antibiotics. [11]



International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429

Volume 8 Issue IV Apr 2020- Available at www.ijraset.com

- 1) Symptoms: The symptoms of this infection include chills and fever. The infection can be treated with the use of antifungal medicines.
- D. Laboratory Diagnosis
- 1) Specimen:
- *a)* Urine in case of urinary tract infection(UTI)
- b) Vaginal discharge in case of genital infection
- c) Cerebrospinal fluid in case of central nervous system (CNS)infections
- 2) *Direct Microscopy:* Gram staining and potassium hydroxide (KOH) mounts shows gram positive yeast cell under microscope (Fig: 1.4).Colonization is detected by demonstration of pseudohyphae (Fig: 1.5) [12]
- Culture: Candida species can grow on sabouard dextrose agar (SDA) media very well and also in ordinary bacteriological media like blood agar. The optimum temperature is 25-37^oC.Creamy and pasty colonies will appear after 24 hours of incubation. [12]



III. ASPERGILLOSIS

The Aspergillosis is caused by a mold (fungus) which is present everywhere. The aspergillus fumigatus is the main causative pathogen. The fungus mainly affects respiratory tract.

The Aspergillosis is caused by the inhalation of Aspergillus conida or mycelial fragments present in soil, in decaying matter. This fungus can cause three types of diseases:

A. Allergic Reaction

Some persons with asthma can develop such type of reactions on exposure to aspergillus mold. These individuals develop a disease known as allergic bronchopulmonary Aspergillosis. [13]

- 1) Treatment: These allergic reactions can be treated by oral corticosteroids to prevent the existing asthma or cystic fibrosis.
- 2) Symptoms
- a) Fever
- b) Cough with blood

B. Aspergilloma

This is a chronic lung disease in which fungus (aspergillus) colonizes in pulmonary cavities and grows into a tangled mass known as aspergillomas. Also known as fungus ball. (Fig: 1.6).

1) Treatment: The aspergillomas cannot be treated with antifungal medicines, so by surgery the fungal mass is removed.



Fig: 1.6



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429 Volume 8 Issue IV Apr 2020- Available at www.ijraset.com

- 2) Symptoms:
- *a)* Cough with blood(hemoptysis)
- b) Wheezing
- c) Difficulty in breathing
- d) Weight loss
- e) Fatigue

C. Invasive Aspergillosis

This is the most severe Aspergillosis and can spread to brain, heart, and kidneys and also infects skin. Invasive Aspergillosis occurs in persons with weak immune system due to cancer chemotherapy, bone marrow transplatation. The untreated Aspergillosis can cause death. [14]

- 1) Treatment: The invasive Aspergillosis can be treated with antifungal medicines like vericonazole, amphoteric B. [15]
- 2) Symptoms:
- *a*) Fever and chills
- b) Coughing with blood
- c) Chest pain
- d) Headache
- e) Skin lesions
- D. Laboratory Diagnosis
- 1) Specimen:
- *a)* Sputum(in case of respiratory infection)
- b) Biopsy
- c) Bronchoalveolar washingS
- 2) *Imaging Test:* A chest X-ray is done which produces the detailed images of the fungal mass in case of aspergillomas infection. (Fig: 1.7). [16]
- 3) Respiratory Secretion (Sputum) Test: The sputum sample is stained and is checked under microscope for the presence of Aspergillus filaments (Fig: 1.8). If the fungus is seen than the sample is cultured to confirm the diagnosis.
- 4) *Biopsy:* In case of invasive Aspergillosis the tissue from the infected lungs is examined under microscope to confirm the diagnosis. [17]



Fig 1.7

Fig: 1.8



International Journal for Research in Applied Science & Engineering Technology (IJRASET)

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429 Volume 8 Issue IV Apr 2020- Available at www.ijraset.com

IV. MUCORMYCOSIS

Mucormycosis (previously called zygomycosis) is caused by Mucor, Rhizopus and Licheimia. These fungi are saprophytes of soil and decaying vegetables. The Mucormycosis are life threatening infections and the risk factors include diabetes, neutropenia and ketoacidosis. These fungus mainly affects the upper respiratory system, digestive tract, brain and also keratin tissue. [18]

A. Symptoms

The symptoms depend on the type of Mucormycosis disease.

- 1) Rhino cerebral Mucormycosis: The symptoms include fever, headache, reddish skin over nose, swelling of eyes (Fig: 1.9), sinuses.
- 2) Pulmonary Mucormycosis: Symptoms include coughing with blood, fever, difficulty during breathing.
- 3) GI Mucormycosis: Symptoms include abdominal pain, vomiting with blood.
- 4) *Cutaneous Mucormycosis:* Symptoms include reddish and swollen skin initially and later it becomes an ulcer with dark Centre and sharp edges (Fig: 1.10). [19]





Fig: 1.10

- B. Laboratory Diagnosis
- 1) Specimen:
- a) Scraping from the site of infection
- b) Nasal exudates
- c) Pus and sputum
- 2) *Direct Microscopy:* KOH (potassium hydroxide) mounts of samples shows non-septate hyphae. Tissues are stained with H and E stain and then seen under microscope for the demonstration of hyphae.
- *Culture:* The samples can be cultured on sabouard dextrose agar (SDA) media after that the colonies are identified.Lactophenol cotton blue (LCB) preparations show branched sporangiophores along aerial mycelium. [20]

C. Treatment

The Mucormycosis can treat by antifungal medicines such as amphoteric B, posaconazole and isavuconazole are given through vein to the patient with Mucormycosis. These medicines are effective against this fungal disease. [21]

V. PENECILLOSIS

This fungal infection is caused by dimorphic fungi known as penicillium rnarneffei.It is more common is persons which have the history of HIV (Human-immunodeficiency-virus) infection. This fungal infection is endemic in Southeast Asia and also in southern parts of China. The penicillium marneffei is a saprophytic fungus and can cause Penicillosis, keratitis, otomycosis and deep infections

Previously this infection was named as Talaromycosis and the causative fungus was called Talaromyces marneffei. But know a day it is called Penicillosis and the fungus is known as Penicillium marneffei. [22]

- A. Symptoms
- 1) Bumps on skin, face and also in neck(Fig: 1.11)
- 2) Fever
- 3) Cough
- 4) Enlargement of liver and spleen
- 5) Abdominal pain and diarrhea
- 6) Shortness of breathing[23]



International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429 Volume 8 Issue IV Apr 2020- Available at www.ijraset.com



Fig: 1.11

- B. Laboratory Diagnosis
- 1) Direct Microscopy: The samples from different infected body parts are collected such as skin, bone marrow, lung or lymph node are cultured in laboratory in suitable media and the growth of septate hyphae of fungus is seen under microscope by making the smears.
- 2) *Culture:* The fungus can grow on sabouard dextrose media (SDA) media and give the white colour colonies and later the colonies become powdery and blue in colour. [24]

C. Treatment

The Penicillosis can be treated by antifungal medicines such as amphoteric B, itraconazole, vericonazole and terbinafine. These medicines are seen effective against this infection. [25]

VI. CONCLUSION

The fungal infections are life threatening complications to every individual who is being infected. So the best approach to avoid these life threatening infections is the early detection of the causative agents, after the signs and symptoms are being identified. So that the appropriate treatment can be given as soon as possible. Clinicians should be familiar about all the diagnostic methods and they have to use the suitable antifungal agents, because they can impact the largest outcome of patients. After the localized fungal infection the appropriate antifungal medicines should be advised by every physician.

REFERENCES

- [1] Mora C, Tittensor DP, Adel S, Simpson AG, Worm B. How many species are there on Earth and in the ocean? PLoS Biol. 2011; 9:e1001127. [PMC free article] [Pub Med] [Google Scholar]
- [2] Brown GD, Denning DW, Levitz SM. Tackling human fungal infections. Science. 2012; 336:647. [Pub Med] [Google Scholar]
- [3] Ribes JA, Vanover-Sams CL, Baker DJ. Zygomycetes in human disease. Clin Microbial Rev. 2000; 13:236–301. [PMC free article] [Pub Med] [Google Scholar]
- [4] Baddley JW. Clinical risk factors for invasive aspergillosis. Med Mycol. 2011; 49(Suppl 1):S7–S12. [PubMed] [Google Scholar]
- [5] Badiee P, Alborzi A, Farhoudi F. A case of Candida mediastinitis after dental extraction. J Infect Dev Ctries. 2011; 5:75-8. [Pub Med] [Google Scholar]
- [6] Traci C, Johnson MD 2018, Candidiasis .Fungal Diseases. United States: Centers for Disease Control and Prevention 2019
- [7] Buchacz K,Lau B,Jing Bosh Abraham AG,Gil MJ.Incidence of AIDS- defining opportunistic infections in a multicolor analysis of HIV-infected persons in United States and Canada 2000-2010.
- [8] Pappas PG, Kauffman CA, Andes DR, Clark CJ, Marr KA, and Ostrosky-Zeichner Clinical practice guideline for the management of candidiasis: 2016 update by the Infectious Diseases society of America.
- [9] Conclaves Ferreira Calves CT, Henrique's Azeredo Silva S.Vulvovaginal candidiasis: epidemiology, microbiology and risk factors 2016.
- [10] Sobel JD 2007.
- [11] Kullberg BJ, ArendrupMC.CDC Invasive candidiasis 2015.
- [12] Segal and Elad 2005, Deorukhkar and Saini 2015, Odds and Bernaerts 1994, Horvath 2003.
- [13] Denning,David W,Pashley,Catherine,Hartl,Domink,Wardlaw,Andrew,Godet,Cendrine,Del Giacco,Stefano,Dalhaes,Laurence,Sergejeva,Svetlana 2015,Fungal allergy in asthma- state of the art and research in needs.
- [14] Goel.Ayush.Pulmonary aspergillosis 2015.



International Journal for Research in Applied Science & Engineering Technology (IJRASET)

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429

Volume 8 Issue IV Apr 2020- Available at www.ijraset.com

- [15] Kontoyiannis DP,Lionakis MS, Lewis RE, C hamilos Healy Perego Safdar A, Kantarjian H,Champlin R,Walsh TJ 2005.Zygomycosis in a tertiary-care cancer in the era of Aspergillus-active antifungal therapy.
- [16] Curtis A, Smith G, Ravin CI 1979. Air crescent sign of invasive aspergillosis.
- [17] Kradin RL, Mark EJ, 2008. The pathology of pulmonary disorders due to Aspergillus Sapp.
- [18] Lee F.Y, Mossad S.B.Adal K.A 1999.Rinaldi M.G 1989.Zygomucosis.
- [19] Nancy F Crum-Cain lone MD MPH 2008, Auluck A 2007, Spellberg B, Edwards J, Ibrahim A 2005, Mucormycosis.
- [20] Michaela Lackner, Rita Caramalho, CornelianLass-Florl, diagnosis of Mucormycosis 2014.
- [21] Lyndsay Mayer 2017, Rebecca J.Frey 2008, Dhurat, Rachita, Kothavade, Rajendra J, Kumar, Anand 2018, Food and Drug Administration, First-line antiretroviral therapy resistant HIV patient with rhinoentomophthoromycosis.
- [22] Yilmaz N, Visagie CM, Houbraken J, Frisvad JC, Samson RA 2014, Vanittanakom N, Cooper CR, Jr Fisher MC, Sirisanthana Penicillium marneffei infection and recent advances in the epidemiology and molecular biology 2006.
- [23] Supparatpinyo K, Khamwan C, Baosoung V, Nelson KE, Wong SY, Wong KF, Penicillium maeneffei infection.
- [24] Sirisanthana Penicillium infection in AIDS patients 2001.
- [25] Le T, Kinh NV, Cuc NTK, A trail of Itraconazole and Amphoteric B for HIV Associated Talaromycosis 2017.

FIGURE SOURCES

- 1) Figure 1.1: Thrush on tongue in child who had taken antibiotics by James Heilman, MD 2020 Wikipedia.
- 2) Figure 1.2: Scientific Figure on ResearchGate.https://www.researchgate.net 2018.Possible risk factors for Candida Esophagitis in Immunocompetent.
- 3) Figure 1.3: Parkland Natural Health and Wellness-Studio.co.in
- 4) Figure 1.4: Article by Acharya Tankeshwar on Candida albicans 2016.https://microbeonline.com
- 5) Figure 1.5: Article by Acharya Tankeshwar on Candida albicans 2016.https://microbeonline.com
- 6) Figure 1.6: Multiple aspergillomas within large cavity lesions of tuberculosis origin. Wikipedia, Medical Dictionary
- 7) Figure 1.7: Review by Sanjay G Revankar, MD, Wayne State University School of Medicine updated on 2019.
- 8) Figure 1.8: Invasive pulmonary aspergillosis .Wikipedia.com
- 9) Figure 1.9: Orbital Mucormycosis in an Immunocompetent Individual by Parisa Badiee in 2012
- 10) Figure 1.10: Wikimedia.org/Wikipedia.com, Mucormycosis
- 11) Figure1.11: Penicillium marneffei Infection AIDS. https://www.researchgate.net/figure/mollusum-contagiosum











45.98



IMPACT FACTOR: 7.129







INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 🕓 (24*7 Support on Whatsapp)