



IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 5 Issue: VII Month of publication: July 2017 DOI:

www.ijraset.com

Call: 🛇 08813907089 🕴 E-mail ID: ijraset@gmail.com



Metavolcanics and Its Litho Structural Relationship with Assosiated Rock Types around Badi Lake, Udaipur, Rajasthan

Diksha Pandey¹, Narendra Kumar Chauhan², Abhay Dashora³ ^{1,2,3} Department of Geology, MLS University, Udaipur-313 001

Abstract : In Indian shield, Aravalli mountain belt is in the northwestern part and uphold a well developed Proterozoic sequence which can be unearthed from approximately 2500 my to 500my. In the present study area, where different types of metasediments are present, exists basic volcanics representing the different episodes of Proterozoic volcanism. Some work has been done on various aspects of geology but detailed structural and petrographical studies on these volcanic have not been carried out in this area.

The study of these volcanics is an excellent opening to understand and unfold the evolutionary history of Aravalli belt. In the present study area, the basic volcanic rocks are intercalated with cleaned washed quartzites and other shallow water sediments. Basically, this sequence of volcano- sedimentary unit overlies the basement of Archean age.

The present work shows that the rock sequence of the Badi lake area can be divided into broad lithological groups representing the different environmental conditions: (l) a composite suite of basic volcanics and tuffs (metamorphosed to green schists), orthoquartzites, conglomerates, brecciated ferrugenious quartzite and dolomites, (2) phyllites with minor intercalatory quartzites, (3) phyllites and schists with abundant intercalatory quartzites and minor impure dolomites.

I. INTRODUCTION

The present study investigation area is falling under Girwa tehsil in NW border of the city of Udaipur district, Rajasthan, India. It belongs to Udaipur division and located about 15 km towards west from district headquarters Udaipur on Udaipur – Badi Lake road. The rocks of the Badi Lake Group occur in between Kharpena to Madar in its south to north extension respectively, covering a distance of 30 km with the width of 3 to 5 km.

The Badi Lake Group represents a volcano-sedimentary assemblage comprising coarse to fine grained clastic sediments, synsedimentational basic volcanics, pyroclastic and volcaniclastic rocks overlying the Udaipur Group. The sequence of meta sedimentary-meta volcanic in the Aravalli Supergroup lies unconformably on the Banded Gneissic Complex (BGC), Heron (1953). The meta- sediments present in the BGC basement include quartzite, marbles, calc-silicate rocks, ironstone formations and mica schists. Out of all these metasediments, only quartzite and marble occur as mappable units on regional scale. Other metasediments are very thin and are highly weathered.

The stratigraphic succession of the rocks occurring in Udaipur district is given in Table1. Aravalli Supergroup has been divided into two groups- Lower Aravalli Group and Upper Aravalli Group (Roy, 1988), separated by an unconformity marked by conglomerate and red beds. Extensive residual weathering leading to the development of ferricrete, silcrete, laterite, ochreous shales and cave breccias in the topmost beds of the carbonate sequence (of the lower Aravalli Group) indicate the span of hiatus between the two groups. Gupta et al (1980, 1992, 1997) figured out the development of the volcanic belt near Badi Lake and hence named it as Badi Lake Group (Table 1).

Table 1. Stratigraphic succession showing the study area in Aravalli Supergroup by Heron (1953), Gupta et al (1997) and Roy

and Jakhar (2002)



International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor:6.887

Volume 5 Issue VII, July 2017- Available at www.ijraset.com

Heron (1953)	Gupta et al (1997)			Roy and Jakhar (2002)		
Delhi System (2000-700m.y)	Delhi Supergroup					Delhi Supergroup
"Raialo series" Aravalli System (Archean)	ARAVALLI SUPERGROUP	Jharol Group Barilake Group Udaipur Group Debari Group (Babarmal Formatio	Samlaji Formation Goram Formation Khamnor Formation Varla Formation Sajjangarh Formation Udaipur Sector Banswara Formation Nimach Mata Formation Balicha Formation Eklinggarh Formation Sabina Formation	ARAVALLI SUPERGROUP	Upper Middle Lower	Lakhawali Formation Jharol Formation Kabita Dolomite Debari Formation Tidi Formation Bowa Formation = Machla Magra Formation Mochia Formation = Zawar Formation Udaipur Formation Jhamarkotra Formation Delwara Formation
Banded Gniess Complex (> 2500 m.y)	Bhilwara Supergroup			' Mewar Gneiss		

A. Meta Volcanics and its Association around Badi Lake

In the north of Udaipur, Heron (1953) reported acquisition of volcanic rocks at the base of the Aravalli System, the ideal exposures being found on the east of Badi Lake (Badi Formation is considered to be equivalent to the Delwara Formation). All these volcanic rocks are supposedly metamorphosed and they show metamorphism of green schist facies.

Here, the Banded Gneissic Complex appears at the base of the ridge quartzite. These ridge quartzites are succeeded by the meta – volcanics upward. Thin bands are also present in it of impure limestone and discontinuous quartzites with sheared granite. The broad belt of amygdular meta – volcanics nearly 3 km wide, passes upward into dull fine-grained light green chlorite-schist which Heron (1953) believed to be the product of volcanic tuffs topping the lava forming most of the remainder of the sequence of meta-volcanics. Some pelitic bands have also been recognised within these meta-volcanics (Fig 1 & 2).

Quartz vien with the invariable association of calcite and epidote are common throughout the exposures and in the amygdules. Heron considered the whole association of meta-volcanics of biotite – chlorite schist, amphibolites and hornblende schists, etc. as the original volcanic effusive but it appears certain that in this belt of rocks mapped as volcanic by Heron (1953), identified as quartzites, pelites, limestones and marly sediments which are metamorphosed. Heron was impressed by the amygdaloidal nature of



International Journal for Research in Applied Science & Engineering Technology (IJRASET)

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor:6.887 Volume 5 Issue VII, July 2017- Available at www.ijraset.com

these rocks at places and classified them as volcanic rocks. Elongation of amygdules and quartz pebbles of quartzite are in the same direction exhibits the same episode of deformation.

Two set of cleavage are usually present in the meta-volcanics (Plate 1D). Intersectional cleavages are always clear and form one of the most common lineations in these rocks. Somewhere in fresh surfaces mineral streaks of biotite and chlorite are present.

On the basis of different litho variation in associated rocks, metavolcanics in the Badi Lake area is divided into three different associations.

B. Metavolcanics with Amygdules and Tuffs

- Massive Amygdular Metavolcanic Association: This association is found in the eastern side of the Badi Lake overlying the basement and extending upto Varla village, depicted by its amygdular character revealing its volcanic nature. The rock is hard and massive and of dark green colour and shows a feebly developed cleavage. At some places brownish green colour is due to the severe effect of weathering which makes it soft and friable in nature. The amygdules seen vary from 2 20 mm long, are filled with quartz, calcite minerals and rarely epidote deformed and elongated showing the preferred orientation in the direction of dip. They are in general fine grained and greenish black to greyish black in colour. (Plate 1A & 1B)
- 2) Conglomerate- Greenschist Association: This association is a characteristic feature of volcanic rocks which are represented by chlorite schist with amygdular and vesicular character in some greenschists revealing its volcanic percentage (lava flows and ash fall tuffs, cf. Heron, 1953; Roy and Paliwal, 1981). The belt of greenschist occupy a large area in the western side of Badi Lake along with diamictic conglomerate, occurs in isolated outcrops occupying the cores of anticlinal folds (antiformal anticline, cf. Ramsay, 1967).

Pebbles and boulders, mostly of quartzite and vien quartz, sometimes its longer axis (X) is as large as 50 cm (Plate 1C). Pebbles of comparatively smaller size (between 0.5-2cm X axis are seen in the outcrops of greenschist near Badi Lake. The schistosity in greenschist is wrapping the flattened pebbles and large clasts.

The contact between the two units in the east is sharp possibly fault controlled. A narrow band of greenschist, at places, alternates with the conglomerate – quartzite near the base of the basal formation.

C. Greenschist Associated with Quartzite and Granite

The contact between greenschist and quartzite is clearly visible in (Plate 2A & 2B) western part of the Badi Lake overlying the amygdular greenschist. An interesting feature associated with this, is the occurrence of pyrophyllite schist at the contact of granite with quartzites (Plate 1E). It precisely follows the contact between the granite and quartzites. The pyrophyllite schists have preserved all the minute details of polyphase deformation. Since pyrophyllite is restricted to a specific time surface, the granite – quartzite contact, therefore it marks the invaluable marker horizon.

 Quartz – Pyrophyllite – Chlorite – Sericite Association: The proportions of mineral present in this association are not constant and vary from place to place. A fine grained rock is slightly foliated and all the constituents minerals are mixed together as they are fine grained. Chlorite is light green to dark green or light yellowish green to green in colour. There are some pockets of pyrophyllite along the contact of granites with the basal conglomerate – quartzite. It is a light green schistose rock with greasy touch, appearing crenulated and puckered in nature (Plate 1D & 1E). It contains quartz, chlorite, pyrophyllite and sericite. Pyrophyllite is white with occasional greenish shade

D. Metavolcanics in Association With Dolomites

These rocks are light green to greenish black in colour and show very strong cleavage. The cleavages are closely spaced and lineations are at steep angle. Somewhere on the cleavage surfaces, glistening flakes of black mica are seen. Somewhere, the rock is like phyllites of the Aravalli System and is indistinguishable from it.

Dolomites appear as grey or rusty brown coloured, massive bodies crisscrossed by viens of quartz. These are poorly bedded rocks comprising of 90% dolomite with minor calcite along with quartz, chlorite, talc, plagioclase and biotite. Fine thin laminated bands are frequently seen in the outcrops (Plate 1F). Sporadic occurrences of silcrete and ferricrete accompanied by laterites, in the outcrops are indicators of paleo-weathering activity (Roy and Paliwal, 1981; Roy and Jakhar, 2002). The carbonates show uniform dolomitic composition suggesting an early dolomitization.

1) Greenschist – Carbonate – Quartzite Association: Quartzites and greenschists are two prominent rock types belonging to study area in this association. The sequence in all starts with the quartzite, which rarely grades into conglomerates. Greenschist bands



International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor:6.887

Volume 5 Issue VII, July 2017- Available at www.ijraset.com

are either interstratified with or overlie conformably the quartzites (Plate 2B). The greenschist is a field term applied to the chlorite schists. Presence of deformed amygdules at some places of these rocks confirms their volcanic percentage.

2) Carbonate association: The basal sequence of conglomerate – quartzite and greenschist passes upwards into the rocks of a calcareous facies, grouped here under this association. The contact between the two is gradational in nature. The dominant rock unit of this association is dolomite, the ferrugineous dolomite, chert, argillaceous limestone and phyllites constitutes a very small portion of this association. Several pockets of talc have also been developed near the intrusive contacts, as a result of hydrothermal alteration of dolomites. Somewhere because of the extremely rapid facies variation from dolomitic limestone to orthoquartzite it is not mappable at this scale the two units separately.

II. PETROGRAPHY

The basal Aravalli volcanic are greenish to dark green coloured, fine to medium grained rocks possessing schistose structure. The green schists are amygdular in places (deep green in colour) and are highly schistose. The schistosity is well marked by preferred dimensional orientation of chlorite, actinolite and or hornblende and subordinate amounts of albite, quartz, epidote and calcite. They are composed of typical greenschist facies mineral assemblage and exhibit a low grade regional metamorphism. These volcanics show some relict minerals, vescicles or amygdules which are preserved despite of pervasive recrystallization, and show porphyritic characters.

A representative specimen from the eastern banks of Badi lake is an actinolite-albite quartz schist containing amygdules filled with brown biotite or pale grey quartz. The vesicles are elongated in the direction of foliation and are 2 - 10mm long.

These metavolcanics in thin sections comprises chlorite, actinolite, hornblende, and albite, orthoclase with grains of quartz, epidote and sphene. The orthoquartzites and conglomerates interstratified with the green schists occur as discontinuous outcrops forming narrow ridges.

Mineralogically, the greenschist comprise chlorite, quartz and feldspar with or without actinolite/hornblende as the chief constituents, whereas epidote, biotite, sericite and sphene are less commonly occurring minerals. Major opaque minerals include magnetite, hematite, pyrite and ilemenite.

Chlorite is the most dominant mineral in greenschists generally occur in the form of irregular flakes and prisms of varying dimensions, slightly elongated showing preferred orientation parallel to S1 plane. The grains are strongly pleochroic in different shades of green.

Amphiboles (actinolite/tremolite or hornblende) occurs as elongated needles or tabular laths. They are present in the form of xenoblast and porphyroblast at some places. Presence of ilemenite, rutile with hornblende, epidote and clino-zoisite suggest the retrogression of pyroxene to amphibole by uralitization. Hornblende is euhedral to subhedral, green in colour, pleochroic in nature (X- pale yellow to brown, Y- light green, Z- dark green).

Pyroxene present is commonly clino-pyroxene (aegerine-augite) with the perfect two set of cleavages. Brown augite is showing one set of cleavage, phenocryst being euhedral to subhedral in shape.

Relict plagioclases occur rarely as medium sized or coarse grains, highly weathered and altered to sericitized aggregates euhedral to subhedral in shape.

Quartz usually occurs as granular aggregates of equidimensional grains. It also occurs as small irregular inequant grains. The grains are flattened and are aligned with actinolite and chlorite. It defines schistosity in meta volcanics.

Epidote occurs more or less rounded grains, as well as elongated laths. Generally as already mentioned, it is a secondary mineral in the vesicles or in the groundmass together with chlorite.

Calcite in some rocks occurs in granular aggregates within the groundmass. Besides, there are some euhedral crystals of calcite and also some calcite viens in the greenschist following either schistosity or fracture planes.

A. Metavolcaincs with Amygdules and Tuffs

- Under microscope, greenschist show a schistose texture (Plate 2E, 3B & E). In rare instances, volcanic flow structure has been
 preserved. Some greenschist show vesicles and amygdules of varying shape and size. These vesicles are invariably flattened and
 stretched (Plate 2C & D). The minerals filling the amygdules are calcite, coarsely crystalline epidote, quartz, feldspar and iron
 oxide.
- 2) At some places, the vesicles do not show any deformation and are spherical in shape except for that the groundmass of the rock shows a sort of pressure shadow effect. There are thin flakes of biotite present which by their parallel orientation define the



International Journal for Research in Applied Science & Engineering Technology (IJRASET)

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor:6.887 Volume 5 Issue VII, July 2017- Available at www.ijraset.com

cleavage in the rock. Quartz and plagioclase are also present. Presence of quartz particularly in the groundmass around the amygdale region, suggests its replacement origin. Arrangement of chlorite flakes are showing folding of second generation which is identified by folding of cleavage plane (Plate 2E).

B. Greenschist Associated with Quartzite and Granite

The rock is dark green coloured and shows schistose texture. The rock shows fairly well developed two set of cleavages (Plate 3B & E). Vesicles and amygdules are elongated in the direction of dip. Some igneous characteristics are shown by flow structure and amygdule structure. While clastic characters and metamorphic characters are shown by parallel arrangements of chlorite and quartz mineral (Plate 2F) in quartzites associated with it. There is an interesting feature present in this association is the occurrence of pyrophyllite schist at the contact of granite with quartzites.

- 1) The rock show perfect cleavage developed by the parallel orientation of biotite flakes and dimensionally elongated quartz, calcite and very subordinate plagioclase feldspar. Chlorite appears distinctly pleochroic from light green to green. Both pyrophyllite and chlorite contains some tiny yellow rutile grains and shows moderate pleochroism. It occurs as small flakes and show grey to greyish brown polarization colours. Sericite also occurs as tiny colourless flakes.
- 2) In a moderately foliated rock, quartz form lenticles parallel to schistosity and sericite changes to muscovite (Plate 3A). In some rocks, calcite is present which is recrystallised into patches and lenticles within chlorite mass
- *3)* In highly foliated, lenticles of quartz are common embedded in chlorite and muscovite demarcating the prominent schistosity of the rock (Plate 3B & E).

C. Metavolcanics in Association with Dolomites

- 1) In the least metamorphosed dolomite, a cluster of very fine grained aggregates embedding small dolomitic spars represents micritic texture. The impact of low grade green schist metamorphism is evident from the occurrence of chlorite fibres, fine needles of penninite-chlorite and biotite. Very few grains of amphibole such as hornblende are also present. Another evidence of upper green schist facies metamorphism is the occurrence of colourless to faint green pleochroic prismatic fibrous laths of tremolite-actinolite which show incipient impact of high grade metamorphism.
- 2) It is a fine grained rock with a very well developed cleavage which is unaffected by any later movements. Sometimes, there are deciphered bands in which the grain size and mineralogical composition is variable. The minerals present are quartz, sericite, chlorite, calcite, magnetite and plagioclase feldspar.
- *3)* The carbonate material is essentially in the form of vein or patches (Plate 3D). Some chlorite has recrystallised at places and the recrystallised chlorite appears to be later to the development of cleavage in the rock. Impure varieties of dolomite contain, besides dolomites and quartz, a few grains of chlorite, phlogopite, rarely tremolite, feldspar and tourmaline.
- 4) Feldspar grains are fresh and have irregular grain boundaries. A few feldspar grains show twinning on albite laws and range in composition between albite and oligoclase (Plate 3C). Feldspar grains show effects of deformation. Unstrained quartz grains occur in clusters, varying in size. Larger grains of quartz show undulose extinction (Plate 3C & D). In some rocks tremolite are seen embedded in the dolomitic matrix and shows sieve structure with pores filled in with dolomites.

III. CONCLUSION

In this area, the basic volcanic rocks occur as lava flows, intercalated with clean washed quartzite and other shallow water sediments. These volcano-sedimentary sequences overlie the basement complex of Archean age popularly known as Banded Gneissic Complex (BGC) and are overlain by a volcanoclastic/conglomerate one. This belt of metavolcanics coincides roughly with the boundary between the shelf and the deep-sea sedimentary facies associations of the Aravalli Supergroup.

On the eastern side of these meta-volcanics, present the high ridge quartzites and on its western side, there are thin bands of impure limestone within these meta volcanic with some quartzites (which are not mappable) which passes upwards to diamictic comglomerate (quartz pebble conglomerate) and some sheared granite.

Petrographically, the rocks display almost uniform mineral assemblage and textural relations. These are generally composed of typical greenschist facies mineral assemblages predominantly comprise serpentinites, which show extensive alteration to talccarbonate, talc and chlorite schist rocks. These metabasic volcanic rocks show iron enrichment.



International Journal for Research in Applied Science & Engineering Technology (IJRASET)

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor:6.887 Volume 5 Issue VII, July 2017- Available at www.ijraset.com

In general the growth of metamorphic minerals has obliterated the primary minerals and textures. However, few relict minerals such as pyroxene and plagioclase and the magmatic features including vesicles and amygdules and ophitic textures are also preserved favours that these are metamorphosed, volcanogenic sediments mixed with clastic seiments.





PLATE 2





F. Sheared quartzite with fine to coarse grained quartz with the development of micaceous mineral along cleavage.





International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor:6.887

Volume 5 Issue VII, July 2017- Available at www.ijraset.com

E. Chlorite schist with quartz, muscovite, biotite and chlorite which together form a fine grain schistose fabric. It exhibits S1 and S2.

REFERENCES

- [1] Ahmad, T., Rajamani, V., 1991. Geochemistry and petrogenesis of the Basal Aravalli volcanic near Nathdwara, Rajasthan, India. Precambrian Research 49, 185-204
- [2] Bhu, H., Sarkar, A., Purohit, R., Banerjee, A., 2006. Characterization of fluid involved in ultramatic rocks along the Rakhabdev Lineament from southern Rajasthan, northwest India. current science, vol. 91, no. 9.
- [3] Chauhan, D.S., 1970. Pyrophyllite deposits of Barors village, Udaipur district, Rajasthan. Bulletin Indian Geological Association 3, 9-12.
- Chauhan, N.K., Sharma, A., Sharma, V., 2004. Geology and Structures of rocks exposed north-east of Gogunda, District Udaipur, Rajasthan. Rajasthan Mineral Bulletin DMG 25, 10-15.
- [5] Chauhan, N.K., Sharma, B.L., Mohemmad Sabah A., 1996. Structural Geometry and strain history of the Early Proterozoic Aravalli rocks of Gorimari Udaipur distric, Rajasthan, Journal Geological society India. 47, 59-74.
- [6] Goel, O.P. 1988. Petrogenesis of Basal Aravalli metabasalts, east of Udaipur, Rajasthan. In: Roy, A.B. (Ed.) Precambrian of Aravalli Mountain Rajasthan, Geol. Soc. India. Mem 7: 317-326.
- [7] Heron, A.M., Geology of Central Rajputana. Memoir Geological Survey India 79, 339.
- [8] Heron, A.M., 1953. Mem. Geol. Surv. India, 79, 1–389.
- [9] Ramsay JG 1967. Folding and fracturing of rocks. Mc Graw Hills Book Co NY, and London, 568.
- [10] Roy, A.B., 1988. Mem. Geol. Soc. India, 7, 3-31.
- [11] Roy, A.B., Jhakar, S.R., 2002. Geology of Rajasthan (Northwest India) Precambrian to Recent. Scientific Publishers (India), Jodhpur, 421 pp.
- [12] Gupta, S.N., Arora, Y.K., Mathur, R.K., Iqbaluddin, Prasad, B., Sahai, T.N. and Sharma, S.B., 1980. Lithostatigraphic map of Aravalli region. Geological Survey of India, Hyderabad.
- [13] Gupta SN, Mathur RK, and Arora YK 1992. Lithostra-tigraphy of Proterozoic rocks of Rajasthan and Gujarat A review. Rec Geol Surv Ind, 115; 7-8; 63-85,
- [14] Gupta, S.N., Arora, Y.K., Mathur, R.K., Iqbaluddin, Prasad, B., Sahai, T.N. and Sharma, S.B. 1997. Mem. Geol. Surv. India, 123, 1–262.
- [15] Naha, K. and Halyburton, R.V. 1974. Early Precambrian stratigraphy of Central and Southern Rajasthan, India. Precamb. Res. 1: 55-73.
- [16] Pandit, M.K., de Wall, H., Chauhan, N.K., 2008. Paleosol at the Archean- Proterozoic contact in NW India revisited evidence for oxidising conditions during paleosol weathering? Journal of Earth System Science117, 201- 210.
- [17] Purohit, R., Sanyal, P., Roy, A.B., Bhattacharya, S.K., 2010. 13C enrichment in the Paleoproterozoic carbonate rocks of the Aravalli Supergroup, northwest India: influence of depositional environment. Gondwana Research 18, 538-546.
- [18] Ram
- [19] say, J.G. (1967): Folding and fracturing of rocks. Mc Graw Hills Book Co NY, and London, 568.Roy, A.B. and Paliwal, B.S., Shekhawat, S.S., Nagori, D.K., Golani, P.R. and Bejarnia, B.R. 1988. Stratigraphy of Aravalli Supergroup in the type area. In: Roy, A.B. (ed.) Precambrian of the Aravalli Mountain, Rajasthan Geol. Soc. India Mem. 7: 121-138,
- [20] Sinha Roy, 1988. Proterozoic Wilson Cycle in Rajasthan. In: Roy, A.B. (ed.). Precambrian of the Aravalli Mountain, Rajasthan. Geol. Soc. India Mem. 7: 95-107
- [21] Sinha-Roy, S., Malhotra, G., Mohanty, M.K., 1998. Geology of Rajasthan. Geological Society India, 278.

FIG: 1 Simplified geological map of Aravalli Supergroup in the type area (After Roy et al. 1988) showing distribution of various lithological units. Scale 5 Km Index **Deep water sediments:** Mica schist, phyllites and thin Bands of quartz 1 **Debari Formation:** Conglomerate-arkose-quartz 8888 Udaipur Formation: Greywackes **Delwara Formation:** Basic volcanic and quartzites JHAMARKOTRA Banded Gneissic Complex: (Pre-Aravalli basement rock) 111















45.98



IMPACT FACTOR: 7.129







INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 🕓 (24*7 Support on Whatsapp)