



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 11 Issue: III Month of publication: March 2023

DOI: <https://doi.org/10.22214/ijraset.2023.50009>

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

Minerals Rich Organic Manure (MROM)

Dr. Shri Pal¹, Ravinder Kumar², Lalit Sharma³

¹Managing Partner, Bunkerman, Plot No 20, Bhatoli Kalan, Baddi (HP)-173205, INDIA

²Partner, Bhandora Organics, Behsuma, Dist Meerut (UP), INDIA

³Chemist (R&D), Bunkerman, Behsuma, Dist Meerut (UP), INDIA

Abstract: *Indiscriminate use of synthetic chemical fertilizers over past few decades in India and other countries of the world, has become a cause of great concern to maintain long term soil fertility, the soil environment, and its components. The soil microbial community plays a major and important role in fostering soil health and plant growth. While the use of synthetic chemical fertilizers has a profound impact on plant growth, it significantly alters the structure of the microbial community towards a detrimental degradation.*

Sustainable farming practices help in reducing the depletion of natural resources and maintain both productivity and soil fertility. The use of natural minerals that contain fertilizer nutrients in their native state is a very promising approach to reducing emissions associated with the manufacturing industries. Organic material from natural sources (biodegradable agricultural waste, food waste, Cow Dung, Gomutra (Cow Urine), etc.) and the waste obtained from saturated filters of Bunkerman CO₂/TVOC Removal Systems, acts as a source of microbial culture and encourages the release of nutrients into the soil during mineral weathering. The combination of nutrient based minerals and their biological weathering agents together with organic matter, has the potential to remediate, restore, and sustain depleted agricultural soils.

In the present invention, four types of "Minerals Rich Organic Manures (MROM-1 to MROM-4)" have been manufactured by making use of the waste obtained from saturated filters of Bunkerman CO₂/TVOC Removal Systems, the biodegradable agricultural waste, Cow Dung, Gomutra (Cow Urine) and other important minerals like Rock Phosphate, Zeolite, Bentonite, Gypsum, Kaolin, Carbon, Silica etc, available in nature.

The emphasis has been on the significance of sustaining agricultural productivity and microbial diversity in the rhizosphere, the region of soil in the vicinity of plant roots in which the chemistry and microbiology is influenced by their growth, respiration, and nutrient exchange.

Keywords: *mineral fertiliser, organic manure; sustainable farming; pollution, microbes, phosphate*

I. INTRODUCTION

It is a well-known fact that normal atmospheric air generally contains 79.03% Nitrogen, 20.94% Oxygen and 0.03% Carbon dioxide by volume. Nitrogen is not absorbed by lungs. Exhaled air contains an average of 4.38% Carbon dioxide [1,2,3]. It is a well known fact that the carbondioxide (CO₂) level in the atmospheric air was naturally maintained around 280 ppm over the past many centuries till about 1776 or a little later until beginning of the Industrial Era. However, due to industrial revolution, the CO₂ level has started increasing exponentially over the past few decades and it has already reached a world average of about 420 ppm. If it is not controlled by innovative solutions, it is likely to cross a figure of about 520 ppm by the year 2050.

To overcome the above problem of CO₂ increase, Bunkerman in India, has recently invented an indigenous technology for which the patent has been filed vide Docket No 128656 dated 17 Nov 2022 with Controller General of Patents, Designs & Trade Marks, India [1,2]. It is observed that when the CO₂, TVOC and other pollutants are absorbed/adsorbed in the filter material of Bunkerman a reasonably high value minerals and compounds are generated inside the saturated filters[1,2]. On saturation the filters may be replaced by the new filters and the saturated material of old filters can be utilised as a raw material to obtain "Minerals Rich Organic Manure (MROM)" by mixing and reacting it with other organic materials and compounds such as biodegradable agricultural waste, Cow Dung, Gomutra (Cow Urine), Rock Phosphate, Zeolite, Bentonite, Gypsum, Kaolin, Carbon, Silica etc.

Minerals are also an important component of the soil; they are the skeleton of the soil and the source of mineral elements[4]. Minerals play an important role in the improvement of soil's physical and chemical properties and the growth and metabolism of microorganisms [5 to 8]. However, the beneficial effects of the use of appropriate minerals in the soil have long been neglected. In addition, our research has found that forms of mineral weathering, such as silicate weathering, are often accompanied by the formation of secondary carbonate minerals in the process of biological weathering, which undoubtedly increases the potential of cultivated soil carbon sinks.

Further studies have shown that secondary minerals formed with mineral weathering have a good remediation effect on heavy metal pollution[9,10]. In addition to the formation of secondary minerals, the cations released by mineral weathering can also combine with the soil's organic complexes through co-precipitation, which in turn mediates the formation of soil aggregates, preserving soil organic carbon, thereby reducing the potential of soil carbon depletion. Therefore, "Minerals Rich Organic Manure (MROM)" has a positive impact on agriculture, soil health and the ecological environment. It provides a new vision and a new dimension for the development of sustainable agriculture in line with reducing the environmental pollution and help in arresting the climate change and global warming.

II. EMBODIMENT

A. Embodiment 1

MROM-1. The material obtained from the saturated BUNKERMAN filters has been found to contain a high value of minerals like N, P, K, Ca, Mg, S, C etc. The constituent and composition depends on the design of the filter adopted and the area in which it has been used. The constituents of the filters used in residential area may well differ from those used in industries and high toxic areas. However, in general, the material contains a good amount of minerals like N, P, K, Ca, Mg, S and C which are useful for the growth of the plants and to maintain biodiversity and clean environment. Therefore, the simplest method of preparing MROM from this waste material is just to crush it into a powder form in dry condition and test its constituents. Additional quantity of minerals or even some additional minerals like Fe, Cu etc can also be added at this stage, if required. The final product is tested in the laboratory to check whether it meets the requirements of the MROM and packed as per the laid down norms for packaging. The Process Flow Diagram is given at Figure 1 below.




Figure 1 : Simplified Process of Manufacturing MROM-1 From Waste of Saturated Filters of BUNKERMAN's CO₂/TVOC and Pollution Removal System

B. Embodiment 2

MROM-2. The preparation of MROM-2 involves simple biological and physical methods as indicated in Figure 2. In this method, the biodegradable agricultural waste, Cow Dung, Gomutra (Cow Urine), the waste obtained from saturated filters of Bunkerman CO₂/TVOC Removal Systems and other important minerals like Rock Phosphate, Zeolite, Bentonite, Gypsum, Kaolin, Carbon, Silica etc, depending on the local availability, are used to manufacture MROM.




Figure 2 : Simplified Process of Manufacturing MROM-2 From Waste of Saturated Filters and Biodegradable Agricultural Wastes

The detailed stepwise procedure is listed below:-

- 1) Cut open the Filter cover and collect the waste (absorbents/adsorbents/molecular sieves and other ingredients) from the filter into an open tray.
- 2) Grind the waste in a grinder to form a fine powder.
- 3) Carry out laboratory tests to determine its constituents.
- 4) Collect the biodegradable agricultural wastes available (such as tree leaves, sugarcane leaves, rice husk, rice plant (parali) , sugarcane bagas, grass, wheat husk, animal feed waste, food waste, fruits & vegetable waste etc). Spread this waste in a composting pit or bed in 3 to 5 layers, one by one and spread a mixture of cow dung, cow urine and water on it so that complete waste material becomes property wet by this fluid mixture.
- 5) Now spread a fine layer of waste material obtained from filters as explained in steps 2 and 3 above.
- 6) Spread a thin layer of any additional minerals such as Rock Phosphate, if required.
- 7) Repeat the steps 4 to 6 above in three to five layers depending on the amount of material available and quantity of MROM required to be manufactured.
- 8) Cover the entire bed or pit with about 5 to 10 cm thick soil layer all around and leave it for natural fermentation/composting for 30 to 45 days.
- 9) After 30 to 45 days, remove the soil from a portion of the bed/pit and check whether the agricultural waste has been properly composted. If not, cover it with soil again and wait for few more days. If yes, then remove the top soil layer from the bed/pit and cut and mix the composted material with a spade or any other tool to obtain a nearly uniform material.
- 10) Add any additional minerals or nutrients, if required and carry out the final quality check in the laboratory for the prepared MROM.
- 11) Your MROM-2 is now ready for packaging, sale and use in organic farming.

C. Embodiment 3

MROM-3. In this method, first the vermi compost is prepared from the agricultural waste by using cow dung and the earth worms. Then the waste obtained from BUNKERMAN filters is processed and mixed with this Vermi compost in appropriate proportion in a mixing bed. Additional minerals/nutrients are then added to it, if required. The prepared MROM is then tested in the laboratory for quality check and then sent for packaging and further use in organic farms. The manufacturing process is illustrated in Figure 3 below.




Figure 3 : Manufacturing Process of MROM-3 From Waste of Saturated Filters and Vermi Compost

D. Embodiment 4

Experiments conducted on Earthworms have shown that the earthworms grow better and faster on mineral rich diet. Earthworms need a continuous supply of calcium. South Australian research found that earthworm numbers doubled when pH(CaCl₂) in their feed/soil rose from 4.1 to 6.7. Experiments were conducted by adding the waste of BUNKERMAN's saturated filters to the feeding material supplied to the earth worms and it was found that the size and number of earthworms was considerably increased in this process. The resulting vermi compost was also found to be rich in mineral contents. Therefore, the fourth type of MROM i.e. MROM-4 was prepared as per the process illustrated in Figure 4 below.




Figure 4 : Manufacturing Process of MROM-4 From Waste of Saturated Filters and Agricultural Wastes by Using Earth Worms




Photo 1 : Photo of Vermicomposting Bed Showing How Worms Convert the Waste Material into MROM-4



Photo 2 : Composting Pits for MROM-2 During Experiments



Photo 3 : Vermi Composting Beds for Manufacturing MROM-4



Photo 4 : Final packaging of MROM-3

III. RESULTS

A. Test Results

Table 1 : Soil Health Card & Soil Test Results : Sample 1 (Soil Only)

किसान का विवरण			
नाम	भरपुरा अर्जुनचंद		
पता	भरपुरा		
ग्राम	भरपुरा (सूर्यपुरा)		
खेती	मसुरा		
जिला	मेरठ		
पिनकोड			
अन्वय संस्था			
मोबाईल संख्या			

मृदा नमूना विवरण	
मृदा नमूना संख्या	480 Farm 161
मृदा एकत्र करने की तिथि	17/3/2023
सर्व संस्था	294-32000-110-4
संस्था नम्बर	
सैल का क्षेत्रफल	
ग्रु-सिद्धि (GPS)	अक्षांश देशान्तर
सिंचित / अतिरिक्त जूनि	सिंचित

संदर्भ उपज के लिए उर्वरक सिफारिशें (जैविक खाद के साथ)			
क्रमिक	काल व दिग्ग	उर्वरक	एन.पी.के. लिए उर्वरक संयोजन-1
1.	गई	सुरिया	एन.पी.के. लिए उर्वरक संयोजन-2
		पोटास	सुरिया 272
		सीपी	एन.पी.के. 140
2.	फसल	सुरिया	पोटास 30
		पोटास	सुरिया 400
		सीपी	एन.पी.के. 150
		सीपी	पोटास 25

प्रयोगशाला का नाम :- कृषि विज्ञान केन्द्र हस्तिनापुर				
सॉयल परीक्षण परिणाम				
क्रमांक	पैरामीटर	परिणाम	इकाई	आकलन
1	पी.एच. (pH)	7.55	जलयीन < 6.5 सामान्य 6.5-7.5 क्षारीय > 7.5	
2	ई.सी. (EC)	0.27	1.0 से कम	
3	जैविक कार्बन (OC)	35	कम 0-0.20 उत्तम 0.21-0.40 मध्यम 0.41-0.60 अच्छा 0.61-0.80	L
4	उपलब्ध नाइट्रोजन (N)	267.6	उत्तम > 250 मध्यम 250-500 अच्छा > 500	
5	उपलब्ध फास्फोरस (P)	24.9	कम 0-10 उत्तम 11-30 मध्यम 31-50 अच्छा 51-100	M
6	उपलब्ध पोटेशियम (K)	120	कम 0-50 उत्तम 51-100 मध्यम 101-150 अच्छा 151-250	M
7	उपलब्ध सल्फर (S)	4.9	कम 10 से कम उत्तम 11 से 15 मध्यम 15 से अधिक अच्छा 15 से अधिक	
8	उपलब्ध जिंक (Zn)	2.8	कम 0.6 से कम उत्तम 0.61-12 मध्यम 12 से अधिक अच्छा 12 से अधिक	
9	उपलब्ध बोरॉन (B)	1.48	कम 0.25 से कम उत्तम 0.26-0.5 मध्यम 0.5 से अधिक अच्छा 0.5 से अधिक	
10	उपलब्ध ऑयन (Fe)	1.9	कम 4 से कम उत्तम 4.1 से 8.0 मध्यम 8 से अधिक अच्छा 8 से अधिक	
11	उपलब्ध मैंगनीज (Mn)	57	कम 0.21 से कम उत्तम 0.21 से 0.4 मध्यम 0.4 से अधिक अच्छा 0.4 से अधिक	
12	उपलब्ध कॉपर (Cu)	52	कम 0.2 से कम उत्तम 0.21 से 0.4 मध्यम 0.4 से अधिक अच्छा 0.4 से अधिक	

Table 2 : Soil Health Card & Soil Test Results : Sample 2 (Soil + 10% Vermi Compost)

किसान का विवरण			
नाम	भरपुरा अर्जुनचंद		
पता	भरपुरा		
ग्राम	भरपुरा (सूर्यपुरा)		
खेती	मसुरा		
जिला	मेरठ		
पिनकोड			
अन्वय संस्था			
मोबाईल संख्या			

मृदा नमूना विवरण	
मृदा नमूना संख्या	442 Farm 161
मृदा एकत्र करने की तिथि	17/3/2023
सर्व संस्था	
संस्था नम्बर	
सैल का क्षेत्रफल	
ग्रु-सिद्धि (GPS)	अक्षांश देशान्तर
सिंचित / अतिरिक्त जूनि	सिंचित

संदर्भ उपज के लिए उर्वरक सिफारिशें (जैविक खाद के साथ)			
क्रमिक	काल व दिग्ग	उर्वरक	एन.पी.के. लिए उर्वरक संयोजन-1
1.	गई	सुरिया	एन.पी.के. लिए उर्वरक संयोजन-2
		पोटास	सुरिया 161
		सीपी	एन.पी.के. 140
2.	फसल	सुरिया	पोटास 30
		पोटास	सुरिया 310
		सीपी	एन.पी.के. 150
		सीपी	पोटास 25

प्रयोगशाला का नाम :- कृषि विज्ञान केन्द्र हस्तिनापुर				
सॉयल परीक्षण परिणाम				
क्रमांक	पैरामीटर	परिणाम	इकाई	आकलन
1	पी.एच. (pH)	7.65	जलयीन < 6.5 सामान्य 6.5-7.5 क्षारीय > 7.5	
2	ई.सी. (EC)	0.35	1.0 से कम	
3	जैविक कार्बन (OC)	59	कम 0-0.20 उत्तम 0.21-0.40 मध्यम 0.41-0.60 अच्छा 0.61-0.80	M
4	उपलब्ध नाइट्रोजन (N)	272.24	उत्तम > 250 मध्यम 250-500 अच्छा > 500	
5	उपलब्ध फास्फोरस (P)	22.4	कम 0-10 उत्तम 11-30 मध्यम 31-50 अच्छा 51-100	M
6	उपलब्ध पोटेशियम (K)	175	कम 0-50 उत्तम 51-100 मध्यम 101-150 अच्छा 151-250	M
7	उपलब्ध सल्फर (S)	15.9	कम 10 से कम उत्तम 11 से 15 मध्यम 15 से अधिक अच्छा 15 से अधिक	
8	उपलब्ध जिंक (Zn)	10.9	कम 0.6 से कम उत्तम 0.61-12 मध्यम 12 से अधिक अच्छा 12 से अधिक	
9	उपलब्ध बोरॉन (B)	58	कम 0.25 से कम उत्तम 0.26-0.5 मध्यम 0.5 से अधिक अच्छा 0.5 से अधिक	
10	उपलब्ध ऑयन (Fe)	2.4	कम 4 से कम उत्तम 4.1 से 8.0 मध्यम 8 से अधिक अच्छा 8 से अधिक	
11	उपलब्ध मैंगनीज (Mn)	62	कम 0.21 से कम उत्तम 0.21 से 0.4 मध्यम 0.4 से अधिक अच्छा 0.4 से अधिक	
12	उपलब्ध कॉपर (Cu)	59	कम 0.2 से कम उत्तम 0.21 से 0.4 मध्यम 0.4 से अधिक अच्छा 0.4 से अधिक	

Table 3 : Soil Health Card & Soil Test Results : Sample 3 (Soil+10% MROM-1)

कृषि स्वस्थता कार्ड		प्रयोगशाला का नाम :- कृषि विज्ञान केन्द्र हस्तिनापुर			
किसान का विवरण		सॉयल परीक्षण परिणाम			
नाम	अशोक कुमार	क्रमांक	पैरामीटर	परिणाम	इकाई
पता	अशोक	1	पी.एच. (pH)	7.60	अपनी < 6.5 सामान्य 6.5-7.5 अधिक > 7.5
ग्राम	अशोक (कल्याण)	2	ई.सी. (EC)	1.0	1.0 से कम
तहसील	नूतान	3	जैविक कार्बन (OC)	4.0	कुल कम < 0.30 उच्च 0.31-0.40 सामान्य 0.41-0.80 अधिक 0.81-0.80
जिला	मेरठ	4	उपलब्ध नाइट्रोजन (N)	220.12	कम < 250 सामान्य 250-500 अधिक > 500
पिनकोड		5	उपलब्ध फॉस्फोरस (P)	18.8	कुल कम 0-10 उच्च 11-20 सामान्य 21-30 अधिक 31-40
अवधार संख्या		6	उपलब्ध पोटैशियम (K)	160	कुल कम 0-50 उच्च 51-100 सामान्य 101-200 अधिक 201-250
मोबाईल संख्या		7	उपलब्ध सल्फर (S)	11.7	कम 0 से कम उच्च 11 से कम सामान्य 12 से अधिक अधिक 13 से अधिक
मृदा नमूना विवरण		8	उपलब्ध जिंक (Zn)	6.8	कम 0.0 से कम उच्च 0.01-12 सामान्य 13 से अधिक अधिक 14 से अधिक
मृदा नमूना संख्या	49/ Fashw'A'	9	उपलब्ध बोरॉन (B)	3.7	कम 0.15 से कम उच्च 0.16-0.5 सामान्य 0.5 से अधिक अधिक 0.6 से अधिक
नमूना एकत्र करने की तिथि	17/3/2023	10	उपलब्ध आयरन (Fe)	3.2	कम 0 से कम उच्च 0.1 से 0.4 सामान्य 0.4 से अधिक अधिक 0.5 से अधिक
सर्वे संख्या		11	उपलब्ध मैंगनीज (Mn)	61	कम 0.21 से कम उच्च 0.21 से 0.4 सामान्य 0.4 से अधिक अधिक 0.4 से अधिक
खेत का नाम		12	उपलब्ध कॉपर (Cu)	5.7	कम 0.2 से कम उच्च 0.21 से 0.4 सामान्य 0.4 से अधिक अधिक 0.4 से अधिक
खेत का क्षेत्रफल					
ग्र-स्थिति (GPS)	अक्षांश देशान्तर				
निष्कर्ष / अतिरिक्त टिप्पणी	सिंचित				
सर्वम उपज के लिए उर्वरक सिफारिशें (जैविक खाद के साथ)					
क्रमांक	कमत व किसिम	उर्वरक	एन.पी.के. लिए उर्वरक संयोजन-1	एन.पी.के. लिए उर्वरक संयोजन-2 (3:2:1)	
1.	गहूँ	सुरिया	210	सुरिया 213	
		पेटासा	68	एन.पी.के. 192	
		दोसी	132	पेटासा 15	
2.	चना	सुरिया	382	सुरिया 387	
		पेटासा	68	एन.पी.के. 203	
		दोसी	143	पेटासा 13	

Table 4 : Test Results of Vermi Compost : Sample 1



NEON		NEON INDUSTRIAL TESTING & RESEARCH LABORATORY		
An ISO 9001 : 2015, ISO 14001 : 2015, ISO 45001 : 2018 Certified		Laboratory : 205 A-8 , Rajpura, Mizwana Road Near Subhash Inter College Meerut		
Mobile : 8218221131, 8445233689, Website-www.neonitri.com, E-mail : neonitri@gmail.com				
TEST REPORT				
VERMICOMPOST ANALYSIS REPORT				
REPORT NO.	VC/NITRI/50720230321-01	PAGE	1 of 1	
SAMPLE CODE:	VC/NITRI/507	SERVICE REQUEST DATE:	21.03.2023	
REPORT ISSUE DATE:	23.03.2023	SAMPLE RECD. ON:	21.03.2023	
NAME & ADDRESS OF CUSTOMER		REF. NO. - NITRI/MSP-21/FR-W/FMT-7.8 SAMPLE DETAILS		
M/S - Bankerman Plot No. 20 Himadra, Batolikalan, Baddi Industrial Area, Solan, Himachal Pradesh, 173205		Description : One Vermicompost Sample received by Neom Representative 21.03.2023 Sample Marking : Vermicompost Sample Analysis Done on : 21.03.2023 to 23.03.2023		
TEST RESULT				
S.No.	Parameter	Result	Unit	Test Method
1.	pH 10 % Sol.	12.98	---	NEON/VC/SOP/005
2.	Moisture	50.36	%	NEON/VC/SOP/008
3.	Organic Matter & Organic Carbon	1.9	%	NEON/VC/SOP/010
4.	TKN	0.083	%	NEON/VC/SOP/012
5.	Sulphur	328	Mg/kg	NEON/VC/SOP/013
6.	Calcium	0.55	%	NEON/VC/SOP/014
7.	Phosphorous	0.041	%	NEON/VC/SOP/016
8.	Potassium	0.2	%	NEON/VC/SOP/018
9.	Magnesium	0.026	%	NEON/VC/SOP/015
10.	Iron	94.2	Mg/kg	NEON/VC/SOP/026
11.	Zinc	6.40	Mg/kg	NEON/VC/SOP/027
12.	Copper	3.22	Mg/kg	NEON/VC/SOP/032
"End of Report"				
				
<p>Notes:-</p> <ol style="list-style-type: none"> 1. Sample will be retained for two weeks from the date of issue of test report, unless specified by the customer. 2. The results given above are related to the tested sample and mentioned parameters. Endorsement of Product is neither inferred nor implied. 3. Total Liability of our works is limited to invoiced amount. 4. This report can not used as evidence in a court of law without the written approval of the lab. 5. Certificate shall not be reproduced except in full, without the written approval of the laboratory. 6. Any sort of play by the customer with the data of this certificate shall be illegal. 				

Table 5 : Test Results of Vermi Compost Plus Waste of BUNKERMAN Filter Material (MROM-1) : Sample 2



NEON INDUSTRIAL TESTING & RESEARCH LABORATORY
 An ISO 9001 : 2015, ISO 14001 : 2015, ISO 45001 : 2018 Certified
 Laboratory : 205 A-B, Rajpura, Mawana Road Near Subhash Inter College Meerut
 Mobile : 8218221131, 8445233689, Website-www.neonitrfl.com, E-mail : neonitrfl@gmail.com



TEST REPORT
VERMICOMPOST ANALYSIS REPORT

REPORT NO.	VC/NITRL/50720230321-02	PAGE	1 of 1
SAMPLE CODE:	VC/NITRL/507	SERVICE REQUEST DATE:	21.03.2023
REPORT ISSUE DATE:	23.03.2023	SAMPLE RECD. ON	21.03.2023
M/S - Bunkerman Plot No. 20 Himnada, Batolikaln, Baddi Industrial Area, Solan, Himachal Pradesh,173205		REF. NO. NITRL/MSP. 21/TR.W/FMT-78	SAMPLE DETAILS
		Description : One Vermicompost Sample received by Neon Representative 21.03.2023 Sample Marking : Vermicompost Sample Analysis Done on : 21.03.2023 to 23.03.2023	

TEST RESULT

S.No.	Parameter	Result	Unit	Test Method
1.	pH 10 % Sol.	9.32	----	NEON/VC/SOP/006
2.	Moisture	56.22	%	NEON/VC/SOP/008
3.	Organic Matter & Organic Carbon	1.76	%	NEON/VC/SOP/010
4.	TKN	0.095	%	NEON/VC/SOP/012
5.	Sulphur	480	Mg/kg	NEON/VC/SOP/013
6.	Calcium	0.51	%	NEON/VC/SOP/014
7.	Phosphorous	0.046	%	NEON/VC/SOP/016
8.	Potassium	0.16	%	NEON/VC/SOP/018
9.	Magnesium	0.023	%	NEON/VC/SOP/015
10.	Iron	66.4	Mg/kg	NEON/VC/SOP/026
11.	Zinc	480	Mg/kg	NEON/VC/SOP/027
12.	Copper	248	Mg/kg	NEON/VC/SOP/032

"End of Report"

Note:-

1. Sample will be retained for two weeks from the date of issue of test report, unless specified by the customer.
2. The results given above are related to the tested sample and mentioned parameters. Endorsement of Product is neither inferred nor implied.
3. Total liability of our works is limited to involved Amount.
4. This report can not used as evidence in a court of law without the written approval of the lab.
5. Certificate shall not be reproduced except in full, without the written approval of the laboratory.
6. Any sort of play by the customer with the data of this certificate shall be illegal.

IV. DISCUSSION

The test results conducted on soils (Table 1 to 3) indicate that the mineral contents of the soil get improved by addition of Vermi Compost and MROM-1. The test results shown in Tables 4 and 5 indicate that the mineral contents of Vermi Compost get significantly enhanced by addition of MROM-1 to the Vermi Compost. This mixture of Vermi Compost and MROM-1, in fact, forms MROM-3 discussed above. The tests results of MROM-2 and MROM-4 also showed a significant amount of mineral content in the manure which are useful for the plant growth.

V. CONCLUSIONS

The "Minerals Rich Organic Manure (MROM)" is manufactured from the biodegradable agricultural waste, Cow Dung, Gomutra (Cow Urine), the waste obtained from saturated filters of Bunkerman CO₂/TVOC Removal Systems and other important minerals like Rock Phosphate, Zeolite, Bentonite, Gypsum, Kaolin, Carbon, Silica etc, available in nature. The emphasis has been on the significance of sustaining agricultural productivity and microbial diversity in the rhizosphere. In this manner, the nitrogen, phosphorus, potassium and total nutrient of the soil, improve greatly with use of MROM in farming. Four types of "Minerals Rich Organic Manures (MROM-1 to MROM-4)" have been manufactured by making use of the waste obtained from saturated filters of Bunkerman CO₂/TVOC Removal Systems, the biodegradable agricultural waste, Cow Dung, Gomutra (Cow Urine) and other important minerals like Rock Phosphate, Zeolite, Bentonite, Gypsum, Kaolin, Carbon, Silica etc, available in nature.

REFERENCES

- [1] Indoor Air Quality Management System Under Closed Conditions, An Invention Under Atm Nirbhar Bharat, Technical Report, Prepared by Bunkerman, Plot No 20, Bhatoli Kalan, Baddi (HP)=173205, INDIA.
- [2] Indoor Air Quality Management System Under Button Up Conditions International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538 Volume 10 Issue XII Dec 2022- Available at www.ijraset.com
- [3] TM-5-858-7 Designing Facilities to Resist Nuclear Weapon Effects, Facilities Support Systems, US Army Corps of Engineers, Headquarters, Dept of the Army, Washington DC, October 1983.
- [4] Bio-Organic Mineral Fertilizer for Sustainable Agriculture: Current Trends and Future Perspectives, Shameer Syed et al, MDPI <https://doi.org/10.3390/min11121336>.
- [5] Ditta, A.; Muhammad, J.; Imtiaz, M.; Mehmood, S.; Qian, Z.; Tu, S. Application of rock phosphate enriched composts increases nodulation, growth and yield of chickpea. *Int. J. Recycl. Org. Waste Agric.* 2018, 7, 33–40.
- [6] Niamat, B.; Naveed, M.; Ahmad, Z.; Yaseen, M.; Ditta, A.; Mustafa, A.; Xu, M. Calcium-enriched animal manure alleviates the adverse effects of salt stress on growth, physiology and nutrients homeostasis of *Zea mays* L. *Plants* 2019, 8, 480.
- [7] Masruroh, A.; Minardi, S. Rock phosphate, zeolite and quail manure to enhance potassium uptake and yield of soybean on alfisols. *Asian J. Soil Sci. Plant Nutr.* 2019, 5, 1–9.
- [8] Li, Y.; Liu, X.M.; Zhang, L.; Xie, Y.H.; Cai, X.L.; Wang, S.J.; Lian, B. Effects of short-term application of chemical and organic fertilizers on bacterial diversity of cornfield soil in a karst area. *J. Soil Sci. Plant Nutr.* 2020, 20, 2048–2058.
- [9] Wang, Q.; Wang, R.; He, L.; Sheng, X. Location-related differences in weathering behaviors and populations of culturable rock-weathering bacteria along a hillside of a rock mountain. *Microb. Ecol.* 2017, 73, 838–849.
- [10] Yang, X.; Lian, B.; Zhu, X.L.; An, Y.L.; Chen, J.; Zhu, L.J. Effects of adding potassium-bearing mineral powder on nitrogen, potassium and potassium contents of chicken manure compost. *Earth Environ.* 2012, 40, 286–292.
- [11] potassium and potassium contents of chicken manure compost. *Earth Environ.* 2012, 40, 286–292.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)