



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 5 Issue: XI Month of publication: November 2017

DOI:

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

Spatial Variations in the Benthic Diatom Flora and Its Diversity in the Highland-Plateau Rivers (Central India)

Sarika Grover¹, Jyoti Verma², Prateek Srivastava¹, Ambrina Sardar Khan¹

¹Amity Institute of Environmental Sciences, Amity University Uttar Pradesh Sec-125 Noida- 201303

²Department of Zoology, University of Allahabad, Allahabad- 211002

Abstract: This study examines spatial distribution of benthic diatom communities from the rivers Chambal, Ken and Tons in Central India. Their floras consist of total 386 taxa, 110, 206 and 212 taxa in Chambal, Ken and Tons, respectively and are quite similar. Flora is highly similar within each river but varies among their corresponding sections (upper to mouth). Richness decreases gradually from source to mouth because substrate heterogeneity declines. Similarity in richness also declines in each river, but increases mildly amongst their corresponding sections, because substrate differs spatially. Alpha diversity is high and similar for the rivers. Species-rich genera are same. Elements of pristine conditions prevail in the flora. In light of these observations the benthic diatom communities are examined for the spatial distribution of flora and species diversity in the highland - plateau rivers Ken and Tons and Chambal. Studies will be useful to understand the regional patterns of diatom diversity and formulate reference conditions in these and other rivers of this region. The study will serve to assess the impacts of the proposed Ken-Betwa Link and climate change.

Keywords: Indian Peninsula, ecoregion, diatom, plateau rivers, Principal Component Analysis

I. INTRODUCTION

The freshwater biodiversity is distributed in a fundamentally different pattern from that in marine or terrestrial systems. Organisms on land or in the sea live in media that are more or less continuous over extensive regions and species adjust their ranges to some degree as climate or ecological conditions change. But freshwater habitats are relatively discontinuous, and many freshwater species do not disperse easily across the land barriers that separate river drainages into discrete units. The loss of biodiversity in an ecosystem has important implications, including diminished resistance and resilience to disturbance, system simplification, and loss of ecological integrity (WRI et al., 1992). In such ecosystems, monitoring and assessment using the resident biota provides both an integrative view of the effects of human influences and a rich variety of signals that can be used to diagnose the causes of degradation (Karr 2006). Biological assessment has thus been recommended for ecological assessment of such ecosystems as it is more reliable, relatively inexpensive and provides a synergistic and holistic approach (Chutter 1998). Various groups of organisms such as phytoplankton, zooplankton, macro invertebrates, periphyton, macrophytes and fish have been recognized as efficient biomonitors (Hering et al. 2006, Resh 2008, Hughes et al. 2012, Wu et al. 2014, Na et al. 2014). However, macro invertebrates and periphyton are the most recommended groups that can integrate the effects of multiple environmental stressors over time (Stevenson and Pan 1999, Kar and Chu 2000), thereby reflecting the ecological status of aquatic ecosystems. Amongst the periphytic communities, diatoms have been established as robust biomonitors and for long have been used for assessment of environmental conditions in streams and rivers (Stevenson et al. 2010). Diatoms constitute over 80% of the primary producers in the in the highland stream and river ecosystems (Nautiyal et al., 1997; 2004 a & b).

Little progress has been made to document the diatom flora, measures of species diversity and assemblages in the highland streams/rivers of India (Rout & Gaur 1994; Nautiyal 2001 & 2005; Nautiyal & Nautiyal 1999; Nautiyal et al., 2000; 2004 a & b; Jüttner & Cox 2001; Nautiyal & Verma 2009; Verma & Nautiyal 2009). The interest is more recent because the uplands, like Himalaya have witnessed a lesser degree of human interference compared to Alps and other mountain chains and can serve as reference sites (Ormerod et al. 1994; Cantonati et al., 2001 & Nautiyal et al., 2004 b) and because increased developmental activities, consequent urbanization is affecting these rather pristine waters (Nautiyal et al., 1996 & 2000). The proposed Ken-Betwa Link (Bundelkhand region, Uttar Pradesh and Madhya Pradesh, NWDA 2006; Krueger et al., 2007) is one such contentious issue in the water starved Bundelkhand region (Central India). The Chambal River is also an intense issue because diatoms from this river have never been explored, though other rivers of the central highland region have received attention (Nautiyal and Nautiyal, 2014; Verma

and Nautiyal, 2016; Nautiyal et al., 2017; Verma et al., 2016). Both are important rivers in the lower section of the Yamuna river system in Central India.

Studies aiming to generate information on reference conditions are needed in the Peninsula and north-eastern parts of the country. In light of these observations the benthic diatom communities are examined for the spatial distribution of flora and species diversity in the highland - Plateau Rivers namely Chambal, Ken and Tons. Studies will be useful to understand the regional patterns of diatom diversity and formulate reference conditions in these and other rivers of this region.

II. METHODOLOGY

A. Study area

The Chambal River originates from the summit of Janapav hill of the Vindhyan range at an altitude of 854 m above the msl at 22°27' N and 75°37' E in Mhow, located in Madhya Pradesh of Central India. It flows through three large states of India namely Madhya Pradesh (M.P.), Rajasthan and Uttar Pradesh (U.P.) whereas Ken and Tons flow between 23°30' to 26°N latitude and 78°30' to 82°30'E, longitude. Both the Rivers are located in a north-eastern part of the Bundelkh and plateau and Vindhya mountain ranges in the Central Highlands ecoregion located to north of the Tropic of Cancer between Narmada in the south and Gangetic Plains in the north. They are separated by smaller rivers (<100 km long) originating in the hills close to the Gangetic Plain. Both of them are ca. 300~350 km long. They find their source in the Kaimur series of the Upper Vindhyan range north of the river Narmada. The Ken flows northwards and joins the right bank of the Yamuna ca. 100 km before its confluence with the Ganga at Allahabad. The Tons lies to east of the Ken and flows in north-easterly direction to join the Ganga, nearly 50 km downstream of Allahabad. Both rivers flow from low to higher latitudes and cut across the plateau as they move towards the Plains. Upper sections of the Ken and Tons are about 20 km apart and the distance gradually increases (~150 km) towards their mouth sections. The Chambal River has three major tributaries namely the Parbati, Kali Sindh, and Banas rivers. Between 1960 and 1972 four multipurpose dams namely Gandhi Sagar, Jawahar Sagar, Ranapratap Sagar and Kota Barrage have been constructed on the upper reaches of Chambal River which have affected its flow considerably (Hussain and Choudhury 1992).

In order to decipher the spatial features of benthic diatom community one sampling station each was selected in the upper, middle, lower and mouth sections of the Ken and Tons (Figure 1) and four sites of National Chambal Sanctuary region of Chambal River (Figure 1). In each section the sampling station was selected on the ease of access by road. The geographical positions and physico-chemical characteristics of the rivers at these stations are presented in Table 1. Agriculture land use prevails on the river banks at all sampling locations, except patch of dry deciduous forest near Panna National Park in the Ken River. The area is largely rural. Urban areas include Banda, and Satna, located 2 to 10 km away respectively from the Ken and Tons in their middle section.

III. MATERIAL AND METHODS

Diatoms were collected from all the sampling sites of three rivers. Benthic diatom flora and communities were obtained by scraping 3 x 3 cm area of 4-5 palm sized stones (cobble) with a toothbrush following standard procedures (Kelly et al. 1998). All substrata were gently shaken and the resulting suspensions were pooled to form a single sample, which was then put in a labeled plastic bottle. All diatom samples were homogenized and fixed with 4% formaldehyde. In the laboratory, diatoms samples were first cleaned with hot HCl & KMnO₄ to remove organic coatings. The treated samples were repeatedly washed to remove all traces of acid and then cleaned with hydrogen peroxide. This method is based on Hasle (1978) and adapted by Round et al. (1990). It has been found suitable for cleaning diatom samples collected in India (Karthick et al. 2010). Diluted samples were mounted in Naphrax for examining the flora using standard literature (Hustedt 1985; Krammer & Lange-Bertalot 1986-1991; Gandhi 1998; Lange-Bertalot 2001; Krammer 2002; 2003; Werum & Lange Bertalot 2004; Metzeltin et al., 2005). The identification and counting of taxa were carried out under a Light Microscope (Leica DM750) at a 100x magnification using immersion oil in accordance with CEN standards (2001). The samples and mounts are stored at the Amity Institute of Environmental Sciences, Amity University, Noida. In each sample 500-600 valves were counted for computing relative abundance, measures of diversity for the community within and among rivers using PAST v.2.03 and CAPS v. 4.0 (Hammer et al., 2001; Seaby & Henderson 2007).

IV. RESULTS & DISCUSSION

A. Diatom communities: Spatial trends in richness and species diversity

The benthic diatom communities of the Chambal, Ken and Tons consists 110, 206 and 212 taxa (species, varieties, forms) respectively in their flora. The total flora of both rivers amounts to 276 taxa from 49 genera (Table 3). Though the flora is significantly similar in these rivers, but not highly similar as expected despite close proximity to each other, same terrain and

ecoregion. Stancheva *et al.*, (2007) also attributed high floristic similarity (0.737) among the Bulgarian rivers, it and Osum (length, 314 km, 189 km respectively; 1800 to 2030 m a.s.l., Balkan Range) to natural factors, such as similar geological characteristics and relief of the catchments of both rivers. Differences, in the flora of the Alaknanda (1 station 3 sites at 500 m a.s.l., 193 taxa, 23 genera, Nautiyal & Nautiyal, 1999 b) and Nagni Gad (4 stations from source to mouth 2000 to 400 m.a.s.l., 136 taxa, 25 genera, Verma & Nautiyal, 2009) that flow through lesser Himalaya, India shows that despite one geological entity they vary in total number of taxa because of their glacierfed and spring fed character and stream order which influence their flows.

Richness of total as well as individual flora from the Chambal, Ken and Tons sampled at 12 locations is considerably high compared with the total flora (114 taxa, 36 genera) from Bulgarian rivers Vit (98 taxa) and Osum (94 taxa) from 28 sites. The richness of the Ken and Tons is also higher than Nagni gad and a Plateau River Damodar (120 taxa, 22 genera, Nautiyal & Nautiyal, 1999 a), but is at par with Alaknanda-Ganga. Except Vit and Osum that are temperate rivers, rest are tropical to sub tropical, so richness cannot be merely attributed to location of the river in tropics and temperate, a geographical factor but also to primary and secondary factors (Begonet *et al.*, 1990). This receives support from the observations on the diversity also. The diversity is higher in all region (H 5.561) and individual rivers the Ken and Tons (Table 2) compared with the Himalayan and Plateau rivers (Nautiyalet *et al.*, 1996, 2004; Jüttneret *et al.*, 1996; Rothfritzet *et al.*, 1997; Nautiyal & Nautiyal, 1999 a & b). The Shannon diversity (H) of the Chambal River was found to be maximum at 2.316 (C2). The Shannon diversity > 4.0 have been reported from the River Indus, Pakistan (2.82-7.95 Ali *et al.*, 2003); Shillong stream (1.6-4.1, Rout & Gaur, 1994); Himalaya and Alps (2.6 to 4.1, Cantonatiet *et al.*, 2001). Each station contains 73% to 88% of river's flora as the richness varies from 90 to 110 taxa in the Chambal, 151 to 182 taxa in the Ken and 156 to 186 taxa in the Tons. The richness is notably higher in the upper and middle stretch and declines in the lower and mouth stretch. A general longitudinal decline in the species richness and diversity is hence evident in these rivers (Table 2) attributed to reduction in substrate size and its heterogeneity from source to mouth. Besides reduction in particle size the alluvium deposited by the Ganga and Yamuna in mouth sections of the Ken and Tons also contributes to decline in substrate heterogeneity. The decrease in diversity also corresponds with declining the altitude of rivers. In the Himalaya Rivers' the Shannon diversity increased longitudinally with decreasing altitude (Nautiyalet *et al.*, 1997; Nautiyalet *et al.*, 2004 a).

The longitudinal variation is minimal because 58 to 70% of the flora is common from source to mouth and very few taxa are added along the course (compared with K1 16 new taxa occur at K2 and 4 taxa at K3). Similarly, some taxa are present at one station only causing decline downstream of its source. Hence, of 182 taxa at K1, 169, 155 and 136 taxa occur at K2, K3, and K4, respectively in the Ken and of 177 at taxa T1, 169, 136 and 128 taxa occur at T2, T3 and T4 in the Tons. The similarity decreases mildly from source to mouth resulting in low intra-river variation (Table 4).

B. Spatial distribution of floral elements

Detrended correspondence analysis (DCA) was performed on the diatom community data to determine the length of the gradient. The gradient length was less than 3 standard deviation units, which suggested the use of linear ordination techniques. Hence, PCA was used to determine the relationship between diatom assemblage composition and sampling sites using the software CANOCO version 4.5 (terBraak 1986).

A PCA was performed on the whole data set with respect to diatom assemblage composition and results obtained (Figure 2). The first four axes accounted for 76.44% of the total variability. The first and second principal axes together explained 54.00% of the total variation, which had an eigen value of 0.372 and 0.167 respectively. Species such as GPAR, ADCK, NCTE and NANT were positively associated with Tones sites which showed high loading values on axis 1 and negatively associated with Ken sites. Kens sites were dominated by the species such as NLIN, NROS, SULN, AMSC and NVIR, which showed high loading values on axis 2 whereas BVIT, NACI and SRUM were dominated by the Chambal River sites.

Brachysiravitrea was dominated by the Chambal River, followed by Nitzschia acicularis, Syne drarumpens and Navicula cryptonella. The flora of the Ken (K) and Tons (T) were dominated by species and varieties of Navicula Bory (K 27, T 29 taxa), Nitzschia Hassal (K 21, T 25 taxa) and Cymbella Agardh (K 21, T 20 taxa), followed by Synedra Ehrenb., Gomphonema Ehrenb., Cymboplectra Krammer, Achnanthisium Kütz. and Amphora Ehrenberg having 8 to 15 taxa (Table 4). This is different from the Himalayan highlands where Navicula Bory (50 taxa), Achnanthisium Kütz. (49 taxa), Cymbella Agardh (39 taxa, includes Cymboplectra), Nitzschia Hassal (28 taxa) are species rich genera (Nautiyal *et al.*, 2004) and Bulgarian ranges where Nitzschia Hassal (27 taxa), Navicula Bory and Gomphonema Ehrenb. (13 taxa each), are species rich (Stancheva *et al.*, 2007). In the Damodar R. the species-rich genera are Synedra Ehrenb., Gomphonema Ehrenb., Cymbella Agardh, Achnanthis Bory, Navicula Bory, Nitzschia Hassal (Nautiyal & Nautiyal, 1999 a). Among some genera that occur either in the Ken or Tons, the genera Frustulia Rabenhorst, Gomphocymbelopsis and Mastogloia Thwaites are exclusive in the Ken while Anomoeoneis Pfitzer and Bacillaria Gmelin in the Tons.

The species of these genera are not common in India compared with species of *Aulacoseira* Thwaites, *Achnanthes* Bory, *Stauroneis* Ehrenberg, *Epithemia* Kützing, *Rhopalodia* O. Muller and *Hantzschia* Grunow (Table 3) that are well known from other parts of the country. Hence, former are truly exclusive compared to the latter that are relatively common.

In the Ken, the species-rich genera *Nitzschia* Hassal, *Synedra* Ehrenb., *Achnantheidium* Kütz. and *Gomphonema* Ehrenb. show decline in the number of taxa from the upper to mouth sections, *Navicula* Bory remain constant and *Cymbella* Agardh shows alternate decrease and increase. In the Tons *Navicula* Bory, *Nitzschia* Hassal, *Synedra* Ehrenb. and *Gomphonema* Ehrenb. show decrease while *Achnantheidium* Kütz. and *Cymbella* Agardh remain constant. Most of the taxa generally decline downstream of the upper section while few remain constant.

V. CONCLUSIONS

The study examines spatial distribution of benthic diatom flora, species richness and diversity among the Chambal, Ken and Tons rivers located in Central India. The region supports higher richness (276 taxa, 46 genera) compared to 110, 206 and 212 taxa in the Chambal, Ken and Tons, respectively, which are considerably similar, as 31 to 33% taxa are specific to respective rivers. Richness decreases gradually from source to mouth because substrate heterogeneity declines. Flora is highly similar within each river but varies among their corresponding upper, middle, lower and mouth sections, despite proximity of rivers. Similarity in richness declines downstream of source in each river, but increases mildly among them, though distance between these rivers increase progressively, attributed respectively to continuum and different substrates in corresponding sections, water quality being similar. *Navicula* Bory, *Nitzschia* Hassal and *Cymbella* Agardh are species rich genera in both rivers. Species richness of most genera decline downstream of the source. Genera representing pristine, oligotrophic water conditions prevail over high conductivity and brackish and pollution tolerant flora.

REFERENCES

- [1] Begon M, Harper JL and Townsend CR, *Ecology: Individual, Populations and Communities*. Cambridge, Massachusetts, USA, (1990).
- [2] Cantonati M, Corradini G, Jüttner I and Cox EJ, Diatom assemblages in high mountain streams of the Alps and the Himalaya. *Beiheft zur Nova Hedwigia*, 123 (2001): 37-61.
- [3] Chutter FM, *Research on the Rapid Biological Assessment of Water Quality Impacts in Streams and Rivers*. Water Research Commission. Pretoria WRC Report No 422/1/98 (1998).
- [4] Gandhi HP, *Fresh water Diatoms of Central Gujarat (With a review and some others)*. Bishen Pal Singh, Mahendra Pal Singh, Dehradun, India, 324 (1998). Hammer O, Harper DAT and Ryan PD, PAST: Paleontological Statistics Software Package for Education and Data Analysis. *Palaeontologia Electronica*, 4(1) (2001): 9.
- [5] Hasle GR, *Some specific preparations: Diatoms In Sournia A (Ed.), Phytoplankton Manual*, UNESCO, Paris, 1-314 (1978).
- [6] Hering D, Johnson RK, Kramm S, Schmutz S, Szoszkiewicz K, Verdonschot PFM, Assessment of European streams with diatoms, macrophytes, macroinvertebrates and fish: a comparative metric-based analysis of organism response to stress. *Freshwater Biology* 51 (2006): 1757-1785.
- [7] Hughes RM, Herlihy AT, Gerth WJ, Pan Y, Estimating vertebrate, benthic macroinvertebrate, and diatom taxa richness in raftable Pacific Northwest rivers for bioassessment purposes. *Environment Monitoring and Assessment* 184 (2012): 3185-3198.
- [8] Hussain SA and Choudhury BC, *The Gangetic dolphin and the status of its habitat in National Chambal Sanctuary*. Proceedings of the "Seminar on conservation of river dolphins of the Indian subcontinent". New Delhi August 18-19. (1992).
- [9] Hustedt F. *The Pennate Diatoms*. Koeltz Scientific Books, Koenigstein, 918 (1985).
- [10] Jüttner I. and Cox EJ, Diatom communities in streams from the Kumaon Himalaya, north-west India. In: A. Economou - Amilli (ed.) *Proceeding of 16th International Diatom Symposium*. Athens and Aegean Islands. Amvrosiou Press, University of Athens, Greece, 237-248 (2001).
- [11] Jüttner I, Rothfritz H and Ormerod SJ. Diatoms as indicators of river quality in the Nepalese Middle Hills with consideration of the effects of habitat-specific sampling. *Freshwater Biology*, 36 (1996): 475-486.
- [12] Karr JR, *Seven foundations of Biological Monitoring and Assessment*. *Biologia Ambientale* 20(2) (2006): 7-18. Karr JR and Chu EW, Sustaining living rivers. *Hydrobiologia* 422/423 (2000): 1-14.
- [13] Karthick B, Taylor JC, Mahesh MK, Ramachandra TV, *Protocols for Collection, Preservation and Enumeration of Diatoms from Aquatic Habitats for Water Quality Monitoring in India*. *The IUP Journal of Soil and Water Sciences* 3(1) (2010): 1-36.
- [14] Kelly MG, Use of the trophic diatom index to monitor eutrophication in rivers. *Water Research* 32 (1998): 236-242.
- [15] Krammer K, *Achnantheaceae, Kritische Ergänzungen zu Navicula (Lineolatae) und Gomphonema*. Vol. 5 English and French translations of the keys and supplements. Stuttgart and Heidelberg, 2(4) (1991): 1-437.
- [16] Krammer K. *Diatoms of Europe - Vol 3, The genus Cymbella*, A. R. G. Gantner Verlag K. G., 2001, 584 (2002).
- [17] Krammer K, *Diatoms of Europe - Vol 4, The genus Cymboplectra, Delicata, Navicymbula, Gomphocymbellopsis, Afrocybella*, A.R.G. Gantner Verlag K.G., 2003, 164 plates, 520 (2003).
- [18] Krammer K and Lange-Bertalot H, *Bacillariophyceae. Die Süßwasserflora von Mitteleuropa*. Vol. 2/1 Naviculaceae, p.1-876 mit 206 pl. Vol. 2/2 Bacillariaceae, Epithemiaceae, Surirellaceae, p. 1-596 (1988). Vol 2/3 Centrales, Fragilariaceae, Eunotiaceae, 2 (1986-1991): 1-576.
- [19] Krueger K, Segovia F and Toubia M, *Assessment of the India River linking plan: a closer look at the ken-betwa pilot link*. By A practicum submitted In partial fulfilment of the requirements For the degree of Master of Science Natural Resources and Environment At the University of Michigan, (2007).

- [20] Lange-Bertalot H. Diatoms of Europe - Vol 2, The genus *Naviculasensu stricto* 10 Genera separated from *Naviculasensu lato* Frustulia, A. R. G. Gantner Verlag, (2001).
- [21] Metzeltin D, Lange-Bertalot H. and Garcia-Rodriguez F, Diatoms of Uruguay. Taxonomy-Biogeography-Diversity. *Iconographia Diatomologica* 15, A.R.G. Gantner Verlag Komanditgesellschaft, FL 9491 Ruggell, 15 (2005): 726.
- [22] Na Y, Erchao L, Dexiang F, Baicai X, Chaoqun W, Meiling Z, Liqiao C, Correlations between zooplankton assemblages and environmental factors in the downtown rivers of Shanghai, China. *Chinese Journal of Oceanology and Limnology* 32(6) (2014): 1352-1363.
- [23] National Water Development Agency, Terms of Reference for Preparation of the Detailed Project Report: Interlinking of Rivers. [Online]. Available: <http://nwda.gov.in/writereaddata/linkimages/9.pdf>. Komanditgesellschaft, FL 9491 Ruggell, 518(2006).
- [24] Nautiyal P, Ecology of mountain streams and rivers in Garhwal Himalaya, In Kumar A ed. Ecology and Conservation of Lakes, Reservoirs and Rivers. ABD Publishers, Jaipur, (2001): 393-417.
- [25] Nautiyal P, Taxonomic Richness in the fish fauna of the Himalaya, Central Highlands and the Western Ghats. *International Journal of Ecology & Environmental Science*, 31(2)(2005): 73 -92.
- [26] Nautiyal P and Singh HR, Fish diversity in some highlands of the Indian subcontinent (Oriental region). In: H R Singh & Nautiyal P [eds] Biodiversity & Ecology of Aquatic Environments, Narendra Publishing House, Delhi, pp 107-156 (2009).
- [27] Nautiyal P and Verma J, Longitudinal variation in the composition of diatom flora in a Central Highland River of Vindhya region, the Ken. *Bull National Institute of Ecology*, 19 (2009): 1-4.
- [28] Nautiyal P and Nautiyal R, Altitudinal variations in the relative abundance of epilithic diatoms in some glacier and springfed Himalayan tributaries of the Ganga River System in the Garhwal region, In: John J [ed.] Proc. 15 Internat. Diatom Symp. Perth, 143-151, Perth, ARG Gantner Verlag K.G., pp 143-151 (2002).
- [29] Nautiyal P, Bhatt JP, Kishor B, Rawat VS, Nautiyal R, Badoni K and Singh HR, Altitudinal variations in phytobenthos density and its components in the cold water mountain river Alaknanda-Ganga. *Phykos*, 36 (1&2) (1997): 81-88.
- [30] Nautiyal P, Kala K and Nautiyal R, A preliminary study of the diversity of diatoms in streams of the Mandakini basin Garhwal Himalaya. In: Proceedings of 17th International Diatom Symposium (Ed. M. Poulin.), Ottawa, Canada, 2002 Biopress, Bristol. pp. 235-269 (2004b).
- [31] Nautiyal P, Nautiyal R, Kala K, and Verma J, Taxonomic richness in the diatom flora of Himalayan streams (Garhwal, India). *Diatom*, 20(2004a): 123-132.
- [32] Nautiyal R and Nautiyal P, Altitudinal variations in the pennate diatom flora of the Alaknanda-Ganga river system in the Himalayan stretch of Garhwal region. In: S. Mayama, M. Idei and I. Koizumi (eds.) Proceedings of Fourteenth International Diatom Symposium Koeltz Scientific Books, Koenigstein, pp. 85-100 (1999a).
- [33] Nautiyal R and Nautiyal P. Spatial distribution of diatom flora in Damodar river system of Chhota Nagpur. In: The Fourth Indian Fisheries Forum (Ed. M. Joseph), Kochi. pp. 17-22 (1999b).
- [34] Nautiyal R., Nautiyal P, and Singh HR, Impact of sewage on the diatom communities of river Alaknanda (Srinagar, Garhwal). *International Journal of Ecology & Environmental Science*, 22(1999b): 289-296.
- [35] Nautiyal R, Nautiyal P and Singh HR, Species diversity of epilithic diatom community. *Tropical Ecology*, 41(2) (2000): 255-258.
- [36] Ormerod SJ, Rundle SD, Wilkinson SM, Daly GP, Dale KM and Juttner I, Altitudinal trends in the diatoms, bryophytes, macroinvertebrates and fish of a Nepalese river system. *Freshwater Biology*, 32 (2)(1994): 309-322.
- [37] Resh VH, Which group is best? Attributes of different biological assemblages used in freshwater biomonitoring programs. *Environment Monitoring and Assessment* 138 (2008): 131-138.
- [38] Rothfritz H, Juttner I, Suren AM and Ormerod SJ, Epiphytic and epilithic diatom communities along environmental gradients in the Nepalese Himalaya implications for the assessment of biodiversity water quality. *Arch. Hydrobiologia*, 138 (1997): 465-482.
- [39] Round FE, Crawford RM, Mann DG, The Diatoms: Biology and Morphology of the Genera. Cambridge University Press, Cambridge 747 (1990).
- [40] Rout J and Gaur JP, Composition of dynamics of epilithic algae in a forest stream at Shillong (India). *Hydrobiologia*, 291 (1994): 61-74.
- [41] Seaby R, and Henderson P, Community analysis (Package 4.0) Searching for structure in community data: PISCES Conservation Ltd., Lymington, England, (2007).
- [42] Stevenson RJ, and Pan Y, Assessing environmental conditions in rivers and streams with diatoms. In 'The Diatoms. Applications for the Environmental and Earth Sciences'. (Eds E. Stoermer and J. P. Smol.) (Cambridge University Press: Cambridge.) 11-40 (1999).
- [43] Stevenson RJ, Pan Y, and van Dam H, Assessing environmental conditions in rivers and streams with diatoms. Smol JP, Stoermer EF (eds) The Diatoms: Applications for the Environmental and Earth Sciences. 2nd edition. Cambridge University Press, Cambridge. 57-85 (2010).
- [44] ter Braak CJF, Canonical correspondence analysis: a new eigenvector technique for multivariate direct gradient analysis. *Ecology* 67 (1986): 1167-1178.
- [45] Verma J and Nautiyal P, Longitudinal patterns of distribution of epilithic diatoms in a lesser Himalayan stream. *Journal of Hill Research*, 2(2)(2009): 105-109.
- [46] Werum Mand Lange-Bertalot, Diatoms in springs from Central Europe and else where under the influence of hydrogeology and anthropogenic impacts. *Ecology Hydrogeology-Taxonomy, Iconographia Diatomologica*, A.R.G. Gantner Verlag., Komanditgesellschaft, FL 9491 Ruggell, 13 (2004): 1-480.
- [47] World Resources Institute (WRI), World Conservation Union (IUCN), and United Nations Environment Programme (UNEP). Global Biodiversity Strategy: Guidelines for Action to Save, Study, and Use Earth's Biotic Wealth Sustainably and Equitably. Washington, D.C.: World Resources Institute, (1992).
- [48] Wu N, Schmalz B and Fohrer N, Study Progress in Riverine Phytoplankton and its Use as Bio-Indicator – a Review, *Austin Journal of Hydrology* 1(1) 2014: 1-9.

Table 1 Geographical co-ordinates of the sampling stations and Physical and chemical characteristics (minimum and maximum) at different stations of the Central Highlands region. (AT, Air temperature; WT Water temperature; pH; C1, Pinahat; C2, ; C3, Etawah; C4, ;K1, Shahnnagar; K2, Panna; K3, Banda; K4, Chilla; T1, Amdara; T2, Maihar; T3, Satna; T4, Chakghat).

River	Ken River				Tons River					Chambal River			
Stations	K1	K2	K3	K4	T1	T2	T3	T4	C1	C2	C3	C4	
Latitude(N)	23°59'28.92"	24°44'17.38"	25°28'38.25"	25°46'15.49"	24°6'30.83"	24°16'14.13"	24°33'42.88"	25°2'1.06"	26°8'6.38"	24°8'6.55"	26°7'0.19"	26°6'9.91"	
Longitude(E)	80°18'17.7"	80°0'41.16"	80°18'51.62"	80°31'36.99"	80°36'1.34"	80°48'18.11"	80°54'26.34"	81°43'51.75"	78°33'8.6"	78°36'77"	78°93'51"	78°93'66"	
Altitude(m asl)	180	135	131	80	360	326	290	94	135	138	197	201	
AT(°C)	23-32	11.5-25	20-24.5	13.4-21.5	28-33	17-30	17.5-28	17-27	14-30	17-28	10-26	12-25	
WT(°C)	20.5-27	15-21.5	17-22	15-24.5	28-26.5	17-30	17.5-28	17-27	19-22	18-21	17.2-24	16.1-20	
pH	7.2-7.5	7.0-7.2	7.2-7.5	7.2-7.5	7.2-7.6	7.0-7.6	7.4-7.6	7.4-7.6	7.4-8.9	7.8-8.3	7.5-8.2	7.4-8.2	

Table 2 Measures of species diversity in the Chambal, Ken and Tons rivers.

	Shannon H	Simpson 1-D
C1	2.229	0.894
C2	2.316	0.899
C3	2.255	0.895
C4	2.094	0.853
K1	5.048	0.992
K2	4.982	0.992
K3	4.853	0.99
K4	4.704	0.989
T1	5.01	0.992
T2	5.031	0.992
T3	4.799	0.99
T4	4.715	0.989

Table 3 Spatial variations in the number of species occurring in various genera recorded from the Plateau Rivers Ken and Tons.

	GENERA	Chambal	Ken	Tons
Thalassiosiraceae	Aulacoseira	1	1	
	Cyclotella	2	1	2
Fragilariaceae	Diatoma	1	2	4
	Fragilaria	1	1	2
	Staurosira	1	1	1
	Synedra	2	13	15
	Tabellaria	1	1	1

Eunotiaceae	Eunotia	3	3	4
Achnanthaceae	Achnanthes	1	1	
	Achnanthidium	8	8	11
	Planothidium	2	4	4
	Cocconeis	3	3	3
Naviculaceae	Amphora	5	10	8
	Anomoeoneis	1		1
	Brachysira	1	1	2
	Caloneis	3	5	5
	Cymbella	8	21	20
	Cymbopleura	8	12	10
	Diploneis	1	5	4
	Encyonema	2	4	4
	Frustulia		1	
	Gomphocymbelopsis	1	1	
	Gomphonema	6	2	3
	Gyrosigma	1	11	12
	Mastogloia	1	1	
	Navicula	12	27	29
	Naviculas. l.		3	3
	Adlafia		1	2
	Craticula	2	4	3
	Diadesmis	1	1	1
	Fallacia		1	2
	Geissleria	1	1	1
Hippodonta		1	2	
Luticola	1	6	7	
Placoneis	2	2	2	
Sellaphora	3	5	5	
Neidium	1	2	2	
Pinnularia	2	3	3	
Stauroneis	1	2		
	Scoliopleura		1	
Epithemiaceae	Epithemia	2		1
	Rhopalodia	1		2
Bacillariaceae	Bacillaria	1		1
	Denticula	1	1	1
	Hantzschia	2	1	
	Nitzschia	12	21	25
Surirellaceae	Surirella	3	8	5
Total Genera		41	42	38
Total species		110	206	211




Figure 2: Principal Component Analysis (PCA) biplot showing the relationship between diatom composition assemblages and selected sampling sites recorded from the Chambal, Ken and Tons River. Acronyms are presented in Appendix I.

Appendix I

Summary of the results derived from Principal Component Analysis (PCA) applied on diatom assemblages and selected sampling sites.

	Axis 1	Axis 2	Axis 3	Axis 4
Eigen value	0.3725	0.1675	0.1285	0.0959
Explained variation (cumulative)	37.25	54.00	66.85	76.44



Appendix Ii

CODE	SPECIES NAME
ADMI	Achnanthydiumminutissimum
AMSC	A. m. v. scotica
ADCK	Achnanthydiumchitrakootense
ALIN	Achnanthydiumlinearis
NCTE	Naviculacryptotenella
SULN	Synedra ulna
CDIV	Cymbelladiversa
AWTE	Amphora twentiana
NCPR	Naviculacapitatoradiata
CTUM	Cymbellatumida
CATL	Cymbellaaustralica
NANT	Naviculaantonii
CEXC	Cymbella excise
NROS	Navicularostellata
NLIN	Nitzschialinearis
GPAR	GomphonemaParvulum
MGRA	Melosira granulate
NVIR	Naviculavidula
BVIT	Brachysiravitrea
SRUM	Synedrarumpens
NACI	Nitzschiaacicularis



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)