



# **iJRASET**

International Journal For Research in  
Applied Science and Engineering Technology



---

# **INTERNATIONAL JOURNAL FOR RESEARCH**

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

---

**Volume: 6      Issue: 1      Month of publication: January 2018**

**DOI: <http://doi.org/10.22214/ijraset.2018.1093>**

**[www.ijraset.com](http://www.ijraset.com)**

**Call:  08813907089**

**E-mail ID: [ijraset@gmail.com](mailto:ijraset@gmail.com)**

# Distribution of ostracode Fauna and Environment of Deposition of Early Miocene (Aquitanian-Burdigalian) Sediments in Kachchh, Gujarat

Nidhi bhanat<sup>1</sup>, M. L. Nagori<sup>2</sup> Maya Chaudhary<sup>3</sup>

<sup>1, 2, 3</sup> Department Of Geology, Mohanlal Sukhadia University 51, Saraswati Marg, Udaipur- 313 002, India

**Abstract:** Kachchh is a pericratonic basin in which Early Miocene sedimentation took place in a shallow marine environment. During the course of study of ostracods from these beds a detailed sampling carried out from a hill section (N 23°24', E 68°49') exposed about eight kilometres from Rampar village, towards Goyela-Matanomadh road. The sediment logical data and ostracod assemblage together reveals the fluctuation in relative sea-level. The change in ostracod assemblage in middle part indicating deepening i.e. middle shelf environment. This is followed by progressive shallowing of the basin marked by deposition of marl. The top part composed of limestone with full of mega fauna indicates again deeper water condition prevail in the area. The paleoecology of the genera viz. *Actinocythereis*, *Neomonoceratina*, *Alocopocythere*, *Cytherelloidea*, *Pontocypris*, *Propontocypris*, *Pseudocythere*, *Semicytherura*, *Xestoleberis*, *Cytheropteron*, *Miocyprideis*, *Pokornyella* and *Occultocythereis* is discussed in the paper. The overall evidence suggests that majority of the genera thrive well in near-shore to shallow marine environment and hence it is inferred that the beds of Early Miocene might have been deposited in near-shore to shallow marine conditions.

**Keywords:** Kachchh, Early Miocene, Khari nadi Formation and Chhasra Formation, Paleoecology and Ostracods.

## I. INTRODUCTION

In Kachchhbasin the Tertiary rocks occur as narrow strip fringing the Mesozoic outcrops of the highlands. The rocks of Tertiary period are exposed in the western part of the basin. The Early Miocene rocks composed of mixed carbonate-siliciclastic sediments. The sedimentation in this basin took place largely in shallow marine environment. Lithostratigraphically it is referred as Khari Nadi Formation at the base and Chhasra Formation at the top. The Khari Nadi Formation uncomfortably overlies the Maniyara Fort Formation of Oligocene age, while the upper contact of the Chhasra Formation is unconformable with Sandhan Formation of Pliocene. On the basis of larger foraminifera, the Khari Nadi Formation is assigned Aquitanian age, while Chhasra Formation is of a Burdigalian age (Raju, 1974, 1991). During the course of study of ostracods from these beds a detailed sampling carried out from a hill section, which is about 55 feet in height (N 23°24', E 68°49') exposed about eight kilometres from Rampar village, towards Goyela-Matanomadhroad (Fig. 1). The sedimentation at this hill section ((N 23°24', E 68°49') commenced, with deposition of clay at the base, followed by siltstone, marl, siltstone, marl and limestone at the top.




Fig. 1. Location and Geological map of the study area, Kachchh (after Biswas, 1992)

## II. PALEOECOLOGY

The sediments at this hill section commenced with deposition of clay (ADK1/1 to ADK1/4) at the base upto six feet in height. In the lower part it is devoid of any fauna, while in upper part rare occurrence of genera like *Alocopocythere*, *Actinocythereis*, *Miocyprideis*, *Paijenborchellina*, *Stigmatocythere* etc. appeared first time.

The ecological data pertaining to certain Recent species of *Alocopocythere* from the Persian Gulf (Bate, 1971; Paik, 1977) and the Mandvi Beach (Jain, 1978) shows that the genus occurs commonly in lagoonal and shallow marine environments, while the studies of living species of the genus *Actinocythere* is from the Persian Gulf (Paik, 1977) and the Mandvi Beach (Jain, 1978) suggest that the genus thrive well in near-shore to shallow marine environments. The living species of the genus also occurs in the off central Bay of Bengal from inner shelf and outer shelf ranging at the depth 47 meters to 105 meters from sandy clay substrate (Nishath et al., 2017).

The genus *Miocyprideis*, according to Ducasse and Moyes (1971), is characteristic of littoral environment. A species of the genus has been observed to inhabit the coastal waters of Gulf of Kachchh, near Okha Port, Saurashtra (Khosla, 1988). Studies of certain living species of the genus *Paijenborchellina* from the Abu Dhabi Lagoon, Persian Gulf (Bate, 1971) shows that it occurs from near-shore shelf to lagoonal environments. The living species of the genus also occurs in the off central Bay of Bengal from Lower-bathyal, ranging at the depth 1487 meters from silty clay substrate (Nishath et al., 2017). Of the various ostracods which occur in this section, little is known about the ecology of the genus *Stigmatocythere* as this does not occur in the present day. The lithological and above ostracod appearance indicates that first marine incursion took place in this region.

This is overlain by 2-3 feet thick siltstone bed (ADK1/5 to ADK1/6) with moderate occurrence of above genera, with appearance of the genus *Cytheropteron*. The genus inhabits in marine waters of all the depths in the present day seas. Deep water species are generally thin-shelled and smooth (Morkhoven, 1963). The species of the genus occurs in the Sendai Bay region on the inner shelf at depths less than 60 meters and middle shelf at depth near the 70 meters from the Pacific coast of north-eastern Japan, (Ikeya and Itoh 1991). Ikeya and Suzuki, 1992, recorded *Cytheropteron* species from outer-shelf (80-150m deep) from off Shimane Peninsula, south western Japan sea. The genus *Cytheropteron* is a eurybathic, reported from the Gaidorf formation of Mühlbach, Molasse Basin, Lower Austria from deeper environments by Zorn (2003). The living species of the genus also occurs in the off central Bay of Bengal from Inner-shelf to upper-abyssal ranging at the depth 47 meters to 2386 meters from clayey silt substrate (Nishath et al., 2017). This is indicative of slight deepening of the basin i.e. inner-shelf to outer-shelf. The siltstone bed is overlain by approximately 7 feet thick marly (ADK1/7 to ADK1/10) bed, with moderate occurrence of all the above genera, with first appearance of genus *Cytherelloidea*. The genus *Cytherelloidea* is essentially a marine form but can also adapt to polyhaline conditions (Morkhoven, 1963; Keen, 1977; Omatsola, 1972; Sohn, 1962, 1964). The genus occurs in various temperatures and depth ranges in marine environments and is a good paleo temperature indicator and in the present day seas it does not survive in temperature less than 10°C. Nishath et al., 2017, recorded the living species of the genus from the off central Bay of Bengal from outer-shelf and middle-bathyal ranging at the depth 749 meters from silty clay substrate.

In the overlying marl (sample ADK 1/9), *Pokornyella* appeared first time and the genus is suggestive of epi-neritic environment (Morkhoven, 1963). The ecological data pertaining to certain living species of the genus from the Uranouchi Bay, Japan (Ishizaki, 1968) shows that it occurs in depths from 2-25 metres, being commonly found in depths from 14 to 32 metres. In upper part firm marine conditions established and abundant occurrence of all the above genera.

In the upper marly bed (ADK 1/12) rare to moderate occurrence of the genera listed earlier, indicates possibly marine regression took place. In addition, genus *Occultocythereis* appeared first time, which shows that the genus is marine form and mostly inhabits in epi-neritic water (Morkhoven, 1963). Above to this 20 feet thick sequence of alternating bands of siltstone and silty shale (ADK 1/13 to ADK 1/24) with abundant earlier ostracods with emergence of new genera like *Xestoleberis*, *Semicytherura*, *Pseudocythere*, *Propontocypris* and *Pontocypris*. The genus *Xestoleberis* though occurs commonly in littoral to epi-neritic environments, can also survive even in brackish water conditions (Morkhoven, 1963; Bold, 1971). Swain (1955), while describing the food of ostracods near La Jolla and San Deigo, California, observed that the genus is a plant feeder and favours a turbulent environment. Benson (1964), while studying the Recent Cytheracean ostracods found the presence of most of the *Xestoleberis* species upto depths of 75 metres. The same genus also reported by Elewa (2004) from Eocene of Cairo, Egypt, indicates a typical steep continental margin of the shallow marine environment. *Xestoleberis* has also been reported by Pezelj et al., (2007) from Late Badenian of Croatia which indicates Middle shelf / Outer shelf environment.

The genus *Semicytherura* is characteristics of shallow marine i.e. littoral environment (Morkhoven, 1963). The same genus also reported by Elewa (2004) from Eocene of Cairo, Egypt, indicates a typical steep continental margin of shallow marine environment.




The representatives of the genus *Pseudocythere* there in the present day sea inhabit at all depths (Morkhoven, 1963), Maddocks (1969), recorded certain living species of the genus *Propontocypris* and *Pontocypris* from Indian ocean which are characteristics of shallow-water near shore environments, especially those offering abundant phytal and coralline micro habitats and are never found at bathyal or abyssal depths and only rarely in offshore shelf sediments. The living species of the genus *Propontocypris* reported from the off central Bay of Bengal from the inner shelf to middle bathyal ranging at the depth 29 meters to 763 meters from sandy clay substrate by Nishath et al., (2017). From above it is concluded that again marine regression took place in the area.

The upper part of siltstone (ADK 1/24 to ADK 1/27) the frequency of ostracods reduces, mainly represented by genera *Actinocythereis*, *Alocopocythere*, *Cytherelloidea*, *Pontocypris*, *Propontocypris*, *Loxoconcha*, *Paijenborchellina* and *Stigmatocythere*, indicating a regressive phase of deposition of sediments. The overlying marl (ADK1/28 to ADK 1/32) having rare occurrence of ostracods like *Actinocythereis*, *Alocopocythere*, *Miocyprideis*, *Pokornyella*, *Xestoleberis*, *Cytherelloidea* etc., indicating that the above beds might have been deposited in a regressive phase, with a very shallow depth. This is overlain by about 15 feet thick, hard limestone bands, which is rich in megafauna indicates again slight deeper water conditions in the area.

From the above distribution it is evident that genera like *Actinocythereis*, *Alocopocythere* and *Neomonoceratina* occurs almost throughout in the Khari Nadi formation and Chhasra formation, while genera like *Pontocypris*, *Propontocypris*, *Pseudocythere*, *Semicytherura*, *Xestoleberis*, *Cytheropteron* appears in the middle part. Ostracods in the top marly part represented by rare occurrence of genera like *Actinocythereis*, *Neomonoceratina*, *Alocopocythere*, *Cytherella*, *Cytherelloidea* etc. (Plate 1).

On the basis of ostracods and lithological association it is concluded that the sediments of the Early Miocene, ages in Kachchh region (as is true of other marine coastal deposits in India) were formed as a result of marine transgression. The deposits thus formed are generally characterized by (1) shallow water depositional environment (2) coarse nature of sediments (3) moderate thickness of beds and (4) their occurrence as thin strips near the existing coast line. The marine transgressions commenced with a gradual advance of shallow epeiric seas over slowly submerging land. Such transgressions bring with them the littoral environment, which is subsequently replaced by comparatively deeper infra-littoral or shallow circa-littoral environments.

**Plate- I**



Explanation of Plate I

1. Cytherelloidea, male carapace, length 0.56mm, left valve view, X64; 2. Miocyprideis, male carapace, length 0.55mm, right valve view, X73; 3. Archicythere is, male carapace, length 0.47mm, left valve view, X87; 4. Occultocythereis, male carapace, length 0.33mm, right valve view, X115; 5. Actinocythereis, male carapace, length 0.91mm, left valve view, X66; 6. Chrysocythere, male carapace, length 0.47mm, right valve view, X102; 7. Stigmatocythere, male carapace, length 0.65mm, right valve view, X72; 8. Alocopocythere, male carapace, length 0.90mm, right valve view, X71; 9. Ruggieria, male carapace, length 0.64mm, left valve view, X81; 10. Aurila, male carapace, length 0.44mm, left valve view, X107; 11. Pokornyella, male carapace, length 0.51mm, right valve view, X82; 12. Loxoconcha, male carapace, length 0.42mm, left valve view, X107; 13. Paijenborchellina, male carapace, length 0.66mm, right valve view, X74; 14. Semicytherura, male carapace, length 0.37mm, right valve view, X119; 15. Xestoleberis, female carapace, length 0.36mm, right valve view, X114; 16. Pseudocythere, male carapace, length 0.29mm, right valve view, X145; 17. Pontocypris, male carapace, length 0.57mm, left valve view, X84; 18. Propontocypris, male carapace, length 0.58mm, right valve view, X74; 19. Phlyctenophora, male carapace, length 0.94mm, right valve view, X89; 20. Cytheropteron, female carapace, length 0.28mm, right valve view, X136.

### III. ACKNOWLEDGMENT

Authors are grateful to Head and Coordinator, SAP, Department of Geology, Mohanlal Sukhadia University, Udaipur for the financial support to carryout field work and for extending SEM, laboratory and library facilities.

### REFERENCES

- [1] Bate, R.H. (1971). The distribution of Recent Ostracoda in the Abu Dhabi Lagoon, Persian Gulf. *Paléocologie des Ostracodes, Colloque Pau (1970), Centre Rech. Pau-SNPA, Bull., vol. 5 suppl., 239-256.*
- [2] Benson, R.H. (1964). Recent marine podocopid and platycopidostracodes of the Pacific. *Publ. Staz. Zool., Napoli, vol. 33, suppl., pp. 387-420.*
- [3] Bold, W.A. Van Den. (1971). Ostracode associations, Salinity and Depth of deposition in the Neogene of the Caribbean region. *Paléocologie des Ostracodes, Colloque Pau (1970), Centre Rech. Pau-SNPA, Bull., vol. 5, suppl., pp. 449-460.*
- [4] Ducasse, O. and Moyes, J. (1971). Intérêt des ostracodes dans une séquence paléogéographique du Tertiaire Nord-Aquitain. *Paléocologie des ostracodes, Colloque Pau (1970), Centre Rech. Pau-SNPA, Bull., vol. 5, suppl. pp. 489-514.*
- [5] Elewa, A.M.T. (2004). Quantitative analysis and palaeoecology of Eocene Ostracoda and benthonic foraminifera from Gebel Mokattam, Cairo, Egypt. – *Palaeogeogr., Palaeoclimatol., Palaeoecol., vol. 211, pp. 309-323.*
- [6] Ikeya, N. and Itoh, H. (1991). Recent Ostracoda from the Sendai Bay region, Pacific coast of northeastern Japan. *Reports of the faculty of Science, Schizuoka University, vol. 25, pp. 93-145.*
- [7] Ikeya, N. and Suzuki, C. (1992). Distributional patterns of modern ostracodes off Shimane Peninsula, southwestern Japan Sea. *Reports of the faculty of Science, Schizuoka University, vol. 26, pp. 91-137.*
- [8] Ishizaki, K. (1968). Ostracodes from Uranouchi Bay, Kochi Prefecture, Japan. *Sci. report, Tohoku Univ., 2<sup>nd</sup> ser. (Geol.), vol. 40, pp. 1-45*
- [9] Jain, S.P. (1978). Recent Ostracoda from Mandvi Beach, west coast of India. *Indian Geol. Assoc., Bull., vol. 11, no. 2, pp. 89-139.*
- [10] Keen, M.C. (1977). Ostracod assemblages and the depositional environments of the Headon, Osborne, and Bembridge Beds (Upper Eocene) of the Hampshire Basin. *Palaeontology, vol. 20, pt. 2, pp. 405-445, pls. 46-49.*
- [11] Khosla, S.C. (1988). Tertiary and Recent species of Miocyprideis from India. In: Hanai, T., Ikeya, N. and Ishizaki, K. (eds.): *Evolutionary biology of Ostracoda, its fundamentals and applications, Development in Paleontology and Stratigraphy, Elsevier, Kondansha, Tokyo, pp. 93-103.*
- [12] Maddocks, R.F. (1969). Recent ostracodes of the family Pontocypridae chiefly from the Indian Ocean. *Smithsonian Contr. Zool., vol. 7, pp. 1-56.*
- [13] Morkhoven, F.P.C.M. Van. (1963). Post-Paleozoic Ostracoda. *Amsterdam Elsevier Publ. Co., vol. 2, pp. 1-478.*
- [14] Nishath, N. M, Hussain, S.M., Neelavnnan, K., Thejasino, S., Saalim, S. and Rajkumar A. (2017). Ostracod biodiversity from shelf to slope oceanic conditions, off central Bay of Bangal, India. *Paleogeography, Paleoclimatology, Palaeoecology, vol. 483, pp. 70-82.*
- [15] Omatsola, M.E. (1972). Recent and Subrecent Trachyleberididae and Hemicytheridae (Ostr. Crust.) from the western Niger delta, Nigeria. *Geological Instn. University Uppsala, Bulletin, N. ser., vol. 3, pp. 37-120.*
- [16] Paik, K. Ho. (1977). Regionale Untersuchungen zur Verteilung der Ostracoden im Persischen Golf und im Golf, von Oman. "Meteor" Forsch. Ergebnisse, Reihe C, no. 28, pp. 37-7
- [17] Pezelj, Đ., Sremac, J. and Sokač, A. (2007). Paleocology of the Late Badenian foraminifera and ostracoda from the SW Central Paratethys (Medvednica Mt., Croatia). *Geologia Croatia, vol. 60/2, pp. 139-150*
- [18] Raju, D.S. N. (1974). Study of Indian Miogypsinoidae. In: Drooger, C.W. (Ed.), *Utrecht Micropaleontology, Bull., vol. 9, pp. 13-53*
- [19] Raju, D.S. N. (1991). Miogypsina scale and Indian chronostratigraphy. *Geoscience Journal, vol. 12, pp. 53-65.*
- [20] Sohn, I.G. (1962). The Ostracoda genus *Cytherelloidea*, a possible indicator of paleotemperature. *U.S. Geological Survey, Article No. 162, pp. 144-147.*
- [21] Sohn, I.G. (1964). The Ostracoda genus *Cytherelloidea*, a possible indicator of paleotemperature. *Publ. Staz. Zool. Napoli, vol. 33, pp. 529-534*
- [22] Swain, F.M. (1955). Ostracoda of San Antonio Bay, Texas. *Journal of Paleontology, vol. 29, pp. 561-646.*
- [23] Zorn, I. (2003). Ostracoda from the Gaiendorf Formation (Middle Miocene, Lower Badenian) of Mühlbach (Molasse Basin, Lower Austria). *Annalen des Naturhistorischen Museums in Wien, vol. 104A, pp. 77-84.*





10.22214/IJRASET



45.98



IMPACT FACTOR:  
7.129



IMPACT FACTOR:  
7.429



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24\*7 Support on Whatsapp)