



# IJRASET

International Journal For Research in  
Applied Science and Engineering Technology



---

# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

---

**Volume: 6**

**Issue: II**

**Month of publication: February 2018**

**DOI:**

[www.ijraset.com](http://www.ijraset.com)

Call:  08813907089

E-mail ID: [ijraset@gmail.com](mailto:ijraset@gmail.com)

# Effect of Salinity on Growth, Photosynthetic Pigments and Antioxidant Activity in Watermelon (*Citrullus lanatus* (L.))

Venkatachalam Vasudevan<sup>1</sup>, Markandan Manickavasagam<sup>2</sup>

<sup>1,2</sup>Department of Biotechnology, Bharathidasan University, Tiruchirappalli, India

**Abstract:** The aim of this study was to examine the effect of different concentrations (0, 50, 100, 150, 200, 250, 300mM) of sodium chloride (NaCl) on growth response, photosynthetic pigments and major antioxidant enzyme activity in watermelon cv. Arkamanik. Salt stressed watermelon plants were analysed after 15 days of treatment, for growth, photosynthetic pigments and antioxidant enzymes activity. The results indicated that growth traits such as plant height, leaf number, total fresh weight, total dry weight and tap root length were decreased, the photosynthetic pigment contents (chlorophyll a, chlorophyll b and carotenoid) also decreased and all the antioxidant enzymes such as superoxide dismutase (SOD), catalase (CAT), peroxidase (POD), guaiacol peroxidase (GPX), ascorbate peroxidase (APX) and glutathione reductase (GR) in leaves were enhanced with increasing salt concentrations. These data showed that antioxidant enzymes appear to protect seedlings against stress-related damage, and play an important role in salinity tolerance of watermelon.

**Keywords:** Salinity; Growth; Photosynthetic pigments; Antioxidants activity; Watermelon

## I. INTRODUCTION

Crop plants are exposed to environmental stresses in field conditions, and about 50% of yield loss caused by biotic and abiotic stresses [1]. Plant development and productivity are adversely affected due to increase in salinization of lands. Overall 6 % of land throughout the world is affected by salinity, it is about 22 % of the total agricultural land [2]. Plants exposed to drought or salinity environment results in considerable changes in physiological, biological and gene expression levels. Plants adopt numerous molecular mechanisms to prevent themselves from the negative impact of salt stress.

Development of salinity tolerant cultivars via genetic engineering is an efficient approach to solve this environmental problem. However these salt-tolerant transgenic plants are failed to survive under field condition with high salinity. To overcome this problem, it is necessary to understand the stress influencing factors for specific genotype during the development under field saline condition. With the development of new molecular biology tools, it is feasible to understand the molecular mechanism behind plant response to salt stress. Hence, the knowledge of candidate salt tolerance genes is a prerequisite for utilization of modern techniques for development of salinity tolerant plants.

The prolonged exposure of plants to soil salinity leads to lethal effect due to osmotic stress and ionic toxicity ([3],[4],[5]). High salinity normally stimulates the excess production of reactive oxygen species (ROS) ([6], [7]). These ROS are systematically regulated by both enzymatic and non-enzymatic antioxidants ([8], [9]). High salinity stress can weaken these regulatory system, results in the lethal effect of plants [10]. Salinity stress also diminishes the activity of photosynthetic system which leads to reduction of chlorophyll contents, stomata number and stomatal conductance [11].

Watermelon (*Citrullus lanatus* L.), is an economically important fruit crop of high nutritious value worldwide and this plant is moderately sensitive to salt stress. The fruit contains 70% of water, due to large leaf area the plant consumes more amount of water naturally ([12], [13],[14]). The cultivation of watermelon is heavily dependent on irrigation, especially during the fruit development stage [15]. China is the largest producer of watermelon with 1,839,750 hectares in 2013 [16].

The plants grown in high salinity leads to limited growth and yield, which also affects the seed viability and germination, but it activates the antioxidant enzyme system. Till now, the physiological and biochemical aspects of salt tolerance in economically important crop plants are relatively unclear. There is not much information about the effects of salinity on watermelon species in the current literature, with the exception of a few studies. Hence, the present study was aimed to examine the growth response, photosynthetic pigment content and major antioxidant enzymes activity in watermelon Arkamanik in response to different salt treatments.

## II. MATERIALS AND METHODS

### A. Plant material

The experiments were carried out in green house, Department of Biotechnology, Bharathidasan University, Tiruchirappalli. The seeds of watermelon cv. Arkamanik were obtained from IIHR, Bangalore, India.

### B. Salinity treatments

Healthy seeds, were selected then washed under tap water with soap solution (Tween 20). Twenty seeds were grown in each plastic pot, containing equal quantities of vermiculite and sand. The seeds were allowed to grow under greenhouse condition, 16/8 hrs (day/night), the temperature was 28–35°C in the daytime and 24–30°C at night and 60-70% relative humidity. Germinated seeds with same growth were transferred to plastic pots with one seed per pot with 1:1 (v/v) autoclaved mixture of sand and vermiculite. The seeds were grown for 4 weeks, the age of plant for treatment was followed by [17],[18].

Different concentrations of sodium chloride (0, 50, 100, 150, 200, 250 and 300 mM) were used to create the salinity conditions. Pots were irrigated with nutrient solution (1x solution) in three days of interval for 4 weeks as described in the earlier work [19]. The seedlings were separated into seven groups, each group was treated with one concentration of NaCl. The nutrient solution alone supplied for control plants. The treatment of concentrations were increased gradually from lower to higher up to the determined concentration of NaCl in nutrient solution. About 200 ml of NaCl solution was given with three days of interval. Samples were collected on the 15th day from the start of treatment, the leaves from three seedlings in each group were analyzed.

### C. Growth measurements

Growth measurements, for the plants exposed to saline treatments, were taken after 15 days of treatment. For each treatment three replicates were used to calculate the mean values. The following measurements were taken after treatment: Plant height (Shoot length), number of leaves in plant, fresh weight and dry weight (plants were dried at 70°C in hot air oven) and the tap root length.

### D. Photosynthetic Pigments

The leaves were extracted with acetone for quantification of photosynthetic pigments [Chlorophyll a (Chl a), chlorophyll (Chl b), and total carotenoids (Caro)] [20] with minor modifications as detailed by the earlier work [21]. The pigment concentrations were expressed as  $\mu\text{g}$  per g fresh weight.

### E. Antioxidant Activity

The harvested leaves were frozen in liquid nitrogen, and used for crude protein extraction as described in the earlier literature [22]. The specific activity of superoxide dismutase (SOD), catalase (CAT), ascorbate peroxidase (APX), peroxidase (POD), glutathione peroxidase (GPX), and glutathione reductase (GR), were quantified in the protein sample. SOD activity was determined by the inhibition of photoreduction of nitrobluetetrazolium (NBT), quantified by spectrophotometer at 560 nm with the influence of riboflavin as superoxide radicals source [23].

The inhibition of 50% of NBT photoreduction was considered as one unit of SOD. CAT activity was measured by consumption of  $\text{H}_2\text{O}_2$  in the protein extract, which showed decrease in absorbance at 240 nm as described earlier [24]. The amount of enzyme required to decompose  $\text{H}_2\text{O}_2$  (1  $\mu\text{mol}$ ) per minute at optimum temperature of 25°C was calculated as one unit of CAT. The oxidation of ascorbate in the reaction was measured by decrease of absorbance (290 nm) under spectrophotometer; from these values the activity of APX was calculated [25].

The amount of enzyme required for consumption of ascorbate (1  $\mu\text{mol}$ ) per minute at optimum temperature of 25°C was calculated as one unit of APX. The activity of GR was measured by the reduction in absorbance peak at 340 nm, due to the oxidation of NADPH [26]. The one unit of GR was considered as the quantity of enzyme required to oxidize 1  $\mu\text{mol}$  of NADPH per minute at optimum temperature of 25°C. Specific POD activity was calculated by the oxidation of benzidine by spectrometric absorbance at 530 nm according to the previous report [27]. Glutathione peroxidase activity was assayed by the protocol followed by [28]. The activities are expressed as  $\mu\text{g}$  GSH consumed /minute/mg protein.

### F. Statistical Analysis

All data obtained were analyzed by one-way analyses of variance (ANOVA) using SPSS 20.0 software. Significant difference from control among treatments were calculated by DMRT (Duncan's multiple range tests) ( $P=0.05$ ).

### III. RESULTS AND DISCUSSION

#### A. Effect of NaCl Stress on Growth of Watermelon

The effect of salinity (NaCl) on growth parameters was analyzed by measuring the plant height, number of leaves, fresh weight and dry weight, and the tap root length of watermelon. The results showed that, the plant height, number of leaves, fresh and dry weight of plants were significantly reduced and the tap root length moderately while increasing the NaCl concentration from 50 to 300 mM (Figure.1). It was noted that the treatment of high salinity (300 mM NaCl) reduced plant growth parameters to more than 55% in comparison with control (Fig. 1). If the concentrations increased above 300 mM of NaCl the plant growth stunted and died after treatment of 15 days (data not shown). Maximum growth inhibition was observed at 300 mM NaCl after 15 days of treatment with 68% of reduction in plant height with respect to control. There was a gradual decrease in plant height with growth inhibition percentage of 17%, 32%, 41%, 52%, 66% and 68% at 50, 100, 150, 200, 250, 300 mM NaCl treatments respectively.

The plant growth was drastically affected by abiotic stress (salinity), it decreasing the water potential and cause nutrient imbalances which leads to plant death ([29],[30],[31]). Therefore, investigating the molecular mechanism behind salt tolerance is prerequisite for plant breeding and genetic manipulation of plants to develop salinity tolerant plants [30]. The number of leaves reduced drastically with reduction percentage of 11%, 21%, 32%, 42%, 52% and 63% after exposed to 50, 100, 150, 200, 250, 300 mM NaCl treatments respectively. The maximum number of leaves was reduced at 300 mM (63%, 7 leaves), when compared to control (19 leaves). The shoot fresh weight was decreased significantly by 20%, 34%, 46%, 53%, 73% and 80% at 50, 100, 150, 200, 250, 300 mM NaCl respectively, with respect to control. The maximum fresh weight reduction occurred at 300 mM (80%, 3g FW) in comparison with control (15 g FW).

The dry weight of shoot decreased gradually by 9%, 20%, 32%, 52%, 62% and 77% at 50, 100, 150, 200, 250, 300 mM NaCl treatments respectively, comparing to control. The maximum dry weight reduction observed at 300 mM (77%, 0.8 g DW), compared to control (3.5 g DW). The tap root length was reduced dramatically by 11%, 22%, 38%, 50%, 62% and 67% at 50, 100, 150, 200, 250, 300 mM NaCl treatments respectively, compared to control. The treatment with 300 mM NaCl showed maximum percentage of root inhibition (67%, 6 cm) with respect to control (18 cm). In the present study, the result conclude that increasing the concentrations of NaCl remarkably decreased the plant height, subsequently fresh and dry weight of the plants. Similar to our results, other studies done in moth bean *Vigna aconitifolia* L. [32], in radish plant, *Raphanus sativus* L. [33], and in black gram *Vigna mungo* L. [34], concluded that increasing the concentrations of NaCl developed a decline in the lengths of the plants. In contrary to our results in maize *Zea mays* L. [35], in rice seedlings *Oryza sativa* L. [36], in cowpea, *Vigna unguiculata* L. [37] and in *Brassica campestris* L. [38] the researchers reported that the length of plants increased in lower concentration of NaCl treatment, at higher concentration the growth of plants drastically reduced.

#### B. Photosynthetic Pigments

The effect of different concentrations of salt stress, on the photosynthetic pigments (chlorophyll 'a', 'b' and carotenoids) of the watermelon plants was investigated. The results showed that an inverse correlation between NaCl concentration and chl. 'a', and 'b' content (Figure. 2). Whenever the concentration of sodium chloride was increased, the chl. 'a', and 'b' content decreased and reached its lowest content, at 300 mM, compared to control plant fresh weights, respectively.

The results showed that when compared to control chl. a content was significantly decreased by 6%, 10%, 13%, 17%, 20% and 25% at 50, 100, 150, 200, 250, 300 mM NaCl treatments respectively. The maximum reduction of chl. a content was observed at 300 mM (25%, 96  $\mu\text{g/g}$  FW) that the control (128  $\mu\text{g/g}$  FW). Similarly, the chl. b content was gradually decreased by 6%, 15%, 24%, 31%, 35% and 41% at 50, 100, 150, 200, 250, 300 mM NaCl treatments respectively. The maximum percentage of chl. b reduction was observed at 300 mM (41%, 32  $\mu\text{g/g}$  FW), compared to control (54  $\mu\text{g/g}$  FW). Our results showed that a decrease in chlorophyll 'a', 'b', and carotenoids, agree with the earlier report [39], barley seedlings exposed to 0, 120 and 240 mM of NaCl decreased the photosynthetic pigments. In addition to this, the studies in *Paspalum vaginatum* (L.) [40] and in *Centaurea erythraea* (L.) [41] explained that chlorophyll 'a', 'b' and total chlorophyll decreased with the increase of salt concentrations. A decrease in chlorophyll content under salt stress, is due to inhibition of Rubisco and PEP carboxylase enzymes, involved in biosynthesis of chlorophyll [42], and the chlorophyll degradation also occurs by the activation of chlorophyllase [43]. In contrary to our results, reported that salt stressed rice seedlings showed increase in the chlorophyll content of 15 days old seedlings [36]. Further it was confirmed in *Beta vulgaris* L. in which the treatment of increased concentration of NaCl, simultaneously increased the leaf total chlorophyll content [33]. The effect of salinity stress on carotenoid content in leaves at different concentrations are given in Figure. 2. The carotenoid content was decreased gradually, with the increasing concentrations of NaCl. There was an inverse relationship between salt concentration and carotenoid content have been observed from the present experiments (Fig 2). The carotenoid content was

gradually decreased by 8%, 16%, 24%, 28%, 40% and 44% at 50, 100, 150, 200, 250, 300 mM NaCl treatments respectively. The reduction percentage of carotenoids was maximum at 300 mM (44%, 28  $\mu\text{g/g}$  FW), with respect to control (50  $\mu\text{g/g}$  FW). The differences of carotene content in the statistical analysis were significant at all the used concentrations of sodium chloride. The reduction of carotenoid content in our study agrees with results in barley seedlings showed drastic reduction of carotenoids [39]. On the other hand, contrary to our results, there was an increase in carotene content in rice seedling exposed to salt stress [36].

### C. Antioxidant enzymatic assay

We observed these significant changes in the antioxidant activity including SOD, CAT, POD, APX, GR and GPX under salinity stress. In the present study, there were significant differences in leaf SOD, POD, GPX, GR, APX and CAT activity under different NaCl concentrations. In addition, all the antioxidant enzyme activity of the leaf was increased gradually with respect to salt concentration, those values were higher in all salt treated plants than that in the control.

The activity of CAT and SOD under salinity stress of different concentrations was presented in Figure 3. There was a significant difference in superoxide dismutase and catalase activity at 50 to 300 mM of salt stressed and untreated seedlings. Compared with control (without salt treatment), CAT activity gradually increased by 4%, 10%, 19%, 24%, 31% and 42% after treated with 50, 100, 150, 200, 250, 300 mM NaCl respectively. The activity of CAT was higher at 300 mM NaCl treatment with increased percentage of 42% ( $145 \text{ U g}^{-1} \text{ min}^{-1} \text{ FW}$ ), than that of control plant ( $102 \text{ U g}^{-1} \text{ min}^{-1} \text{ FW}$ ). CAT is an efficient antioxidant produced during stress conditions, it prevents the plants from oxidative damage by degrading the hydrogen peroxide ([44], [45]). During unfavourable stress conditions the catalase activity was improved to act as a protective agent against oxidative stress caused by toxic effect of  $\text{H}_2\text{O}_2$ , produced throughout the metabolic processes of plant cells ([45], [46], [47]). Similar to our results, in *Ipomoea pescaprae* the catalase activity increased with increasing the concentrations of NaCl [48]. Increasing in catalase activity under different salt stress was reported in *Cassia angustifolia*, maize, wheat and *Sesamum indicum* ([49], [50], [51], [52]). SOD is an important antioxidant, with the capability to repair oxidative damage caused by ROS. Hence, SOD acts as an efficient enzyme which maintains the physiological conditions and the regulation of intercellular levels of ROS under oxidative stress [44]. In the present study, NaCl stress to watermelon increased the SOD activity with increasing the concentrations. The control leaves showed low level of enzyme activity than the salt treated leaf samples. The SOD activity gradually increased by 9%, 16%, 24%, 29%, 42% and 49% after treated with 50, 100, 150, 200, 250, 300 mM NaCl respectively. The activity of SOD peaked at 300 mM NaCl with increased percentage of 49% ( $142 \text{ U g}^{-1} \text{ min}^{-1} \text{ FW}$ ), than that of control plant ( $95 \text{ U g}^{-1} \text{ min}^{-1} \text{ FW}$ ). Similar to our report, others observed that significant increase in SOD activity due to salt stress, increase in ROS ( $\text{O}_2^-$ ) simultaneously increase SOD activity, because this enzyme acts as a scavenger of  $\text{O}_2^-$  [53].

The effect of NaCl on the APX and POD activity in the leaves of watermelon at various concentrations was presented in Figure 4. Ascorbate peroxidase and peroxidase activity also showed a similar increasing trend up to maximum concentration of 300 mM NaCl. There was a gradual increase in activity of POD by 5%, 15%, 20%, 25%, 35% and 45% after treated with 50, 100, 150, 200, 250, 300 mM NaCl respectively, when compared to control. The higher activity of POD was observed at 300 mM NaCl (45%,  $29 \text{ U g}^{-1} \text{ min}^{-1} \text{ FW}$ ) with respect to control plant ( $20 \text{ U g}^{-1} \text{ min}^{-1} \text{ FW}$ ). Similar to our results, in mangrove such as *Aegiceras corniculatum* there was a significant increase in peroxidase activity under saline stress [54]. During salt stress increase in peroxidase activity indicates that the formation of huge amount of hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) and which could release from membrane structure [55]. The peroxidase enzyme involved in the decomposition of co-substrates like phenolic compounds and antioxidants. The isomers of peroxidase consume  $\text{H}_2\text{O}_2$  and phenolic compounds for the synthesis of secondary metabolites necessary for plant growth, development and differentiation [56]. In the present study, POD activity was increased in all the stressed plants than the control plants, it was agreed with other researchers report that increase in POD activity under salinity stress [57]. Salinity condition induced the production of ROS [6], which can damage cellular proteins, lipids and nucleic acids ([58], [59], [60]). In our experiments, the APX enzyme activity was higher under salt treatment than in the control leaves, this concluded that APX activities showed better relation with salinity treatment. The APX activity increased progressively in a concentration dependent manner with 5%, 13%, 18%, 21%, 26% and 31% after treated with 50, 100, 150, 200, 250, 300 mM NaCl respectively. The activity of APX was higher at 300 mM NaCl (31%,  $50 \text{ U g}^{-1} \text{ min}^{-1} \text{ FW}$ ), compared to control ( $38 \text{ U g}^{-1} \text{ min}^{-1} \text{ FW}$ ). There are several reports which confirmed the increase of antioxidant enzyme activity during salinity treatments. In pea plant, the antioxidants ascorbate peroxidase (APX), dehydroascorbate reductase (DHAR) and monodehydroascorbate reductase (MDHAR) were increased under salt stress conditions. During stress conditions the cellular metabolism was regulated by APX and GR, and the cellular redox state was maintained properly which enables the plants to tolerate against environmental stresses [61].

The GPX and GR activity against different concentrations of NaCl was presented in Figure.5. There was a significant increase in the glutathione peroxidase and glutathione reductase activity up to 300mM of NaCl. The Compared with control, the GR activity dramatically increased by 16%,33%,38%,50%,55% and 66% after treated with 50,100,150,200,250,300 mMNaCl respectively. The activity of GR peaked at 300 mMNaCl with increased percentage of 66% ( $0.30 \text{ U g}^{-1} \text{ min}^{-1} \text{ FW}$ ), than that of control plant ( $0.18 \text{ U g}^{-1} \text{ min}^{-1} \text{ FW}$ ). In the present study, we found that GPX activity significantly increased by 11%,16%,27%,33%,38% and 55% after treated with 50,100,150,200,250,300 mMNaCl respectively. The activity of GPX was higher at 300 mMNaCl (55%,  $2.8 \text{ U g}^{-1} \text{ min}^{-1} \text{ FW}$ ), compared with control ( $1.8 \text{ U g}^{-1} \text{ min}^{-1} \text{ FW}$ ). Our results were in agreement with the reports of other researchers, that the activities of APX and GR were increased in *A. portulacoides*[62], *Salicornia brachiata*[63] and *B. parviflora*[64] under salinity stress. Similarly in *L. stocksii* the activity of antioxidants plays a key role in detoxification of ROS under salt stress [8]. Plants exposed to salinity were prone to oxidative stress because of the formation of ROS such as  $\text{O}_2^-$ ,  $\text{H}_2\text{O}_2$  and  $\text{OH}^\cdot$  ([65],[66]). These ROS can affect the integrity of cellular membranes, enzymes activities and the plant photosynthetic apparatus ([8], [67]). The antioxidants enzymes are abundantly present in plant cells, it act as a scavenger to protect the cells from damage caused by oxidative stress. The plant defense mechanism plays a vital role in plant stress tolerance, it majorly depends on the activity of SOD, CAT and POD antioxidant enzymes [68],[69].

#### IV. CONCLUSION

In conclusion, watermelon growth was decreased at higher concentration of NaCl and the levels of photosynthetic pigments also reduced. This study also indicate that the increase in antioxidant enzyme activity provides the detoxification mechanism, which may improves the salinity tolerance in watermelon. Salt-induced enhancement of antioxidant enzymes was the most important mechanisms for its stress tolerance. Further investigation on analyzing the expression level of antioxidant genes to salt stress will provide basic regulatory mechanism behind the salt tolerance. These candidate genes will be highly useful to develop plants with field salt tolerance.

#### V. ACKNOWLEDGEMENTS

The first author is thankful to Bharathidasan University, Tiruchirappalli, for the award of University Research Fellowship.

#### REFERENCES

- [1] S.Vij and AK. Tyagi, "Emerging trends in the functional genomics of the abiotic stress response in crop plants," *Plant Biotechnology Journal*, vol.5, pp. 361–380, 2007
- [2] [http://www.fao.org/worldfoodsituation/en/?utm\\_source=faohomepage&utm\\_medium=web&utm\\_campaign=featurebar](http://www.fao.org/worldfoodsituation/en/?utm_source=faohomepage&utm_medium=web&utm_campaign=featurebar)
- [3] JK. Zhu, "Plant salt tolerance," *Trends in Plant Science*, vol. 6, pp. 66–71. 200
- [4] R.Munns and M. Tester, "Mechanisms of salinity tolerance," *Annual Review of Plant Biology*, vol.59, pp. 651–681,2008.
- [5] S. Yu, W. Wang and B.Wang, "Recent progress of salinity tolerance research in plants," *Russian Journal of Genetics*, vol.48, pp.497–505, 2012
- [6] R. Ozgur, B.Uzilday, AH. Sekmen and I. Turkan, "Reactive oxygen species regulation and antioxidant defence in halophytes" *Functional Plant Biology*, vol.40, pp.832–847, 201
- [7] KB. Hamed, F. Chibani, C. Abdely and C. Magne, "Growth, sodium uptake and antioxidant responses of coastal plants differing in their ecological status under increasing salinity," *Biologia*. Vol.69, pp.193–201, 201
- [8] MN. Jithesh, SR. Prashanth, KR. Sivaprakash and AK. Parida, "Antioxidative response mechanisms in halophytes: their role in stress defence," *J Genet.*, 85(3) 237-54, 200
- [9] J.Bose, A.Rodrigo-Moreno and SJ. Shabala, "ROS homeostasis in halophytes in the context of salinity stress tolerance," *Exp Bot.*, 65(5):1241-57, Mar 2014
- [10] A. Hameed and MA. Khan, "Halophytes: biology and economic potentials," *Karachi University Journal of Science*, vol. 39, pp.40–44, 2011
- [11] R.M. Balal, MA. Shahid, MM. Javaid, "Foliar treatment with *Lolium perenne* (Poaceae) leaf extract alleviates salinity and nickel-induced growth inhibition in pea," *Braz. J. Bot* vol. 39, pp. 453, 2016
- [12] H. Yetisir and V. Uygun "Plant growth and mineral element content of different gourd species and watermelon under salinity stress," *Turkish Journal of Agriculture and Forestry*, vol. 33, pp.65-67, 2009
- [13] H. Zhang, G.Gong, S. Guo, Y. Ren, Y. Xu and KS. Ling, "Screening the USDA watermelon germplasm collection for drought tolerance at the seedling stage," *Hortscience* vol.46 9, pp. 1245-1248, 2011
- [14] S. Guo, J.Liu, Y. Zheng, M. Huang, H. Zhang and G. Gong, "Characterization of transcriptome dynamics during watermelon fruit development: sequencing, assembly, annotation and gene expression profiles," *BMC Genomics*. Vol. 12, pp.454,2011
- [15] C. Kaya, D. Higgs, H. Kirnak and I. Tas, "Mycorrhizal colonisation improves fruit yield and water use efficiency in watermelon (*Citrullus lanatus* Thunb.) grown under well-watered and water-stressed conditions," *Plant Soil* vol.253, pp. 287–292, 2003.
- [16] <http://faostat3.fao.org/download/Q/QC/E> (FAO of the United Nations (2015). Food and Agricultural Commodities Production.
- [17] HA. Schreiber and CO. Stanbery "Barley production as influenced by timing of soil moisture and timing of N application," *Agron.J*.pp. 442-445, 1965
- [18] AD. Day and P.K Thompson "Effect of soil moisture regimes on growth of barley" *Agron.J*. vol. 67, pp.430-433, 1975
- [19] B. Etherton, "Relationship of cell transmembrane potential to potassium and sodium accumulation ratios in oats and pea seedlings," *Plant physiol*. Vol. 38, pp.581-585, 1963

- [20] HK. Lichtenthaler and AR. Wellburn, "Determinations of total carotenoids and chlorophylls a and b of leaf extracts in different solvents," *BiochemSoc Trans*, vol.11, pp.591-592, 1983
- [21] AO. Aremu, MW. Bairu, L. Szucova, JF.Finnie and J. Van Staden, "The role of meta-topolins on the photosynthetic pigment profiles and foliar structures of micropropagated 'Williams' bananas," *J.PlantPhysiol*, vol. 169(15), pp.1530-1541, 2012
- [22] W.Beyer and I. Fridovich, "Assaying for superoxide dismutase activity: some large consequences of minor changes in conditions" *Anal Biochem*, vol. 161, pp.559-566, 1987
- [23] H. Aebi, "Catalase in vitro," *Methods Enzymol*, Vol.105, pp. 121-126, 1984.
- [24] Y. Nakano and K. Asada, "Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts." *Plant Cell Physiol* vol. 22, pp. 867- 880, 1981
- [25] JP. Connell, and JE Mullet, "Pea chloroplast glutathione reductase: purification and Characterization" *Plant Physiol*, vol. 82, 351-356, 1986
- [26] H. Rahnama, and H. Ebrahimzadeh, "The effect of NaCl on antioxidant enzyme activities in potato seedlings," *BiologiaPlantarum*, vol. 49,
- [27] CH. Foyer, N. Souriau, S. Perret, M. Lelandais, KJ. Kunert, C. Pruvost and L. Jouanin, "Overexpression of glutathione reductase but not synthetase leads to increases in antioxidant capacity and resistance to photoinhibition on poplar trees," *Plant Physiol* vol. 109, pp. 1047–1057, 1995
- [28] MMF. Mansour and KHA. Salama, "Cellular basis of salinity tolerance in plants," *Environmental and Experimental Botany*, vol. 52, pp.113-122, 2004
- [29] V. Chinnusamy, A. Jagendorf and JK. Zhu, "Understanding and improving salt tolerance in plants," *Crop Science*, vol.45, pp. 437-448,2005
- [30] GK. Genc, M. Mcdonald and Tester, "Reassessment of tissue Na<sup>+</sup> concentration as a criterion for salinity tolerance in bread wheat," *Plant, Cell & Environment*, vol. 30, pp. 1486-1498, 2007
- [31] Mathur N, Singh J, Bohra A, Vyas A "Biomass production, productivity and physiological changes in moth bean genotypes at different salinity levels". *Am.J. Plant Physiol*. 1(2): 210-213. 2006
- [32] Jamil M, REhman S, Rha E.S "Salinity effect on plant growth, psII photochemistry and chlorophyll content in sugar beet (*Beta vulgaris* L.) and cabbage (*Brassica oleracea capitata* L.)". *Pak. J.Bot.*39(3): 753-760, 2007
- [33] Kapoor K, Srivastava A "Assessment of salinity tolerance of Vignamungo var.Pu-19 using ex vitro and in vitro methods". *Asian J.Biotechnol*. 2(2): 73-85, 2010
- [34] Hamada A.M. "Alleviation of the adverse effects of Nacl on germination, seedling growth and metabolic activities of maize plants by calcium salts". *Bull.Fac.Sci.AssiutUinv*. 24:211-220, 1995
- [35] Misra A, Sahu A.N, Misra M, Singh P, Meera I, Das N, Kar M, Sahu P "Sodium chloride induced changes in leaf growth and pigment and protein contents in two rice cultivars". *Biol.Plantarum*. 39(2): 257-262, 1997
- [36] Dantus B.F, Riberio L, Aragao C.A "Physiological response of cowpea seeds to salinity stress. *Rev.Bras.Sementes*". 27(1): 144-148, 2005
- [37] Memon S.A, Hou X, Wang L.J "Morphological analysis of salt stress response of pakchoi". *EJEAF che*. 9(1): 248-254, 2010
- [38] Tort, N., Turkyilmaz, B., "A physiological investigation on the mechanisms of salinity tolerance in some barley culture forms". *J.F.S*. 27, 1–16, 2004
- [39] Lee, G., Carrow, R.N., Duncan, R.R., "Photosynthetic responses to salinity stress of halophytic seashore paspalum ecotypes". *Plant Sci*. 166 (6), 1417–1425, 2000
- [40] Siler, B., Mistic, D., Filipovic, B., Popovic, Z., Cvetic, T., Mijovic, A., "Effects of salinity on in vitro growth and photosynthesis of common centaury (*Centaurium erythraea Rafn.*)". *Arch. Biol. Sci*. 59 (2), 129–134, 2007.
- [41] Soussi M, Ocana A, Lluch C. "Effect of salt stress on growth, photosynthesis and nitrogen fixation in chickpea (*Cicer arietinum* L.)". *J Exp Bot.*;49: 1329–1337, 1998.
- [42] Santos CV. Regulation of chlorophyll biosynthesis and degradation by salt stress in sunflower leaves. *Sci Hort.*;103: 93–99., 2004
- [43] Mittler R., "Oxidative stress, antioxidants and stress tolerance". *Trends Plant Sci.*, 7: 405–410, 2002
- [44] Willekens H., Inzé D., Van Montagu M., Van Camp W. "Catalases in plants". *Mol. Breed.*, 1: 207–228, 1995
- [45] Sairam R.K, Rao K.V, Srivastava G.C "Differential response of wheat genotypes to long term salinity stress in relation to oxidative stress, antioxidant activity and osmolyte concentration". *Plant sci*. 163: 1037-1046, 2002
- [46] Willekens H, Inze D, Van Montague M, Van camp W., "Catalase in plants. *Mol Breed*.1: 207-228, 1995
- [47] Dionisio-Sese M.L, Tobita S., "Antioxidant response of rice seedlings to salinity stress". *Plant sci*. 135:1-9, 1998.
- [48] Venkatesan A, Chellappan KP., "Salinity effect on the activities of certain anti-oxidant enzymes in *Ipomoea pes-caprae* sweet- a halophyte". *Ind J Plant Physiol*.4:40-42, 1999.
- [49] Agarwal S, Pandey V., "Antioxidant enzyme responses to NaCl stress in *Cassia angustifolia*". *Biol Plant*. 48: 555-560, 2004.
- [50] Azevedo Neto A.D, Prisco J.T, Eneas-Filho J, Braga de Abreu C.E, Gomes-Filho "Effect of salt stress on anti-oxidant enzymes and lipid peroxidation in leaves and roots of salt-tolerant and salt-sensitive maize genotypes". *Environ Exp Bot*. 56: 235-241, 2006
- [51] Mandhania S, Madan S, Sawhney V . "Antioxidant defence mechanism under salt stress in wheat seedlings". *Biol Plant*. 50: 227-231, 2006
- [52] Koca H, Bor M, Ozdemir F, Turkan I., "The effect of salt stress on lipid peroxidation , anti-oxidant enzymes and proline content of sesame
- [52] Esfandiari E, Shekari F, Shekari F, Esfandiari M., "The effect of salt stress on antioxidant enzymes' activity and lipid peroxidation on the wheat seedling" *Not Bot HortAgrobotCluj*. 35: 48-56, 2007
- [53] Manikandan T, Venkatesan A., "Influence of NaCl on growth, organic constituents and certain antioxidant enzymes of *Aegicerascorniculatum* Blanco". *Geobios*.31:30-3, 2004
- [54] Zhang, J. and M.B. Krikham., "Drought stress induced changes in activities of SOD, catalase and peroxidase in wheat spp. *Plant Cell Physiol.*, 35: 785-79, 1994
- [55] Gasper T.H., Penel C., Hagega D. and Greppin H., "Peroxidases in plant growth, differentiation and development process". *Biochemical, molecular and physiological aspects of plant peroxidases*. pp. 249-280, 1991
- [56] Jung S., "Variation in antioxidant metabolism of young and mature leaves of *Arabidopsis thaliana* subjected to drought". *Plant Science*, **166**: 459–466, 2004.
- [57] Blokhina O, Virolainen E, Fagerstedt K.V., "Antioxidants, oxidative damage and oxygen deprivation stress: a review", *Annals of Botany*. 91:179-194, 2003
- [58] Apel K, Hirt H., "Reactive oxygen species: metabolism, oxidative stress, and signal transduction", *Annual Review of Plant Biology*.55: 373-399, 2004
- [59] Gill SS, Tudeja N. "Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants". *Plant Physiology and Biochemistry*. 48: 909-930, 2010

[60] Saed-Moucheshi A, Shekoofa A, Pessarakli M. "Reactive oxygen species (ROS) generation and detoxifying in plants". J Plant Nutr. 37:1573-1585, 2014

[61] Benzarti M, RejebK.B,Debez A, Messedi D, Abdelly C., "Photosynthetic activity and leaf antioxidant responses of Atriplexportulacoidessubjected to extreme salinity". ActaPhysiologia Plantarum.34: 1679-1688, 2012

[62] Parida A.K, Jha B., "Salt tolerance mechanisms in mangroves: a review," Trees. 24: 199-217, 2010.

[63] Parida A.K, Das AB, Mohanty P., "Defence potentials to NaCl in a mangrove, Bruguieraparviflora: differential changes of isoforms of some antioxidant enzymes". Journal of Plant Physiology. 161: 531-542, 2004

[64] Van Breusegem F, Vranova E, Dat J.F, InzeD.,"The role of active oxygen species in plant signal transduction", Plant science. 161: 405-414, 2001

[65] Parida A.K, Das A.B., "Salt tolerance and salinity effects on plants: a review", Ecotoxicology and Environmental safety. 60: 324-349, 2005

[66] Serrano R, Mulet J, Rios G, Marquez J, De Larrinoa I, Leube M, Mendizabal I, Pascual-Ahuir A, Proft M, Ros R, Montesinos., "A glimpse of the mechanisms of ion homeostasis during salt stress". Journal of Experimental Botany. 50: 1023-1036, 1999

[67] Matamoros M.A, Dalton D.A, Ramos J, Clementa M.R, Rubio M.C, BecanaM.,"Biochemistry and molecular biology of antioxidants in the rhizobia-legume symbiosis". Plant Physiology.133: 499-509, 2003

[68] Xue F.F, Liu L, Liu Z.P, Mehta S.K, Zhao G.M., "Protective role of Ca against NaCl toxicity in Jerusalem Artichoke by up-regulation of antioxidant enzyme". Pedosphere.18: 766-774, 2008.




Figure 1. Effect of different concentrations of NaCl on growth parameters of watermelon cv. Arkamanik.




Figure 2. Effect of different concentrations of NaCl on photosynthetic pigments in leaves of watermelon cv. Arkamanik.






Figure 3. Effect of different concentrations of NaCl on CAT and SOD activity in leaves of watermelon cv. Arkamanik.




Figure 4. Effect of different concentrations of NaCl on APX and POD activity in leaves of watermelon cv. Arkamanik.




Figure 5. Effect of different concentrations of NaCl on GPX activity in leaves of watermelon cv. Arkamanik.




Figure 6. Effect of different concentrations of NaCl on GR activity in leaves of watermelon cv. Arkamanik.



10.22214/IJRASET



45.98



IMPACT FACTOR:  
7.129



IMPACT FACTOR:  
7.429



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24\*7 Support on Whatsapp)