



IJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 6

Issue: II

Month of publication: February 2018

DOI:

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

Geodetic Polynomial and Detour Geodetic Polynomial of Radial Graphs

Dr.G.Ramesh¹, A.Sathiya²

¹Associate Professor, Department of Mathematics, Government Arts College(Autonomous), Kumbakonam, Tamilnadu, India.

²Assistant Professor, Department of Mathematics, Thanthai Hans Roever College(Autonomous), Perambalur, Tamilnadu, India.

Abstract: Geodetic polynomials of paths, complete graphs and complete bipartite graphs, radial graphs are given. The detour geodetic polynomials of complete, complete bipartite and radial graphs are also obtained.

Keywords: Distance, Detour, Geodetic Polynomial, Radial Graphs. **AMS Classification:** 05C12, 05C60, 05C75

I. PRELIMINARIES

In this section we present some definitions and results which are essential in the study of succeeding sections.

A. Introduction

The standard distance $d(u,v)$ between two vertices u and v in a connected graph G is the length of a shortest $u-v$ path in G . Although this concept has been known for a very long time, it is only in recent decades that received considerable attention as a subject of its own. In 1990 Buckley and Harary wrote the book Distance in graphs. In 2004 Handbook of graph theory edited by Groes and Yellen[4] contains a section devoted exclusively to distance in graphs. A number of results on distance come from the fact that two vertices u and v are adjacent if and only if $d(u,v)=1$ and two distinct vertices u and v are non adjacent if and only if $d(u,v) \geq 2$. Hence any concept whose definition relies on the adjacency or non adjacency of two vertices in a graph can be restated in term of distance. In this paper we find the geodetic polynomials of some graphs, also we have found the detour geodetic polynomial of some graphs.


1) *Definition 1.1.1:* Let $G(V,E)$ be a simple graph. Let $u, v \in V$ be any two vertices of G . A $u-v$ path of length $d(u, v)$ is called a $u-v$ geodesic.

Let $I[u,v]$ be the set of all vertices lying on some $u-v$ geodesic of G . The set $I(S)$ is defined by

$$I[S] = \bigcup_{u,v \in S} I[u,v]$$

2) *Definition 1.1.2:* A set S of vertices of G is called a geodetic set in G if $I[S] = V[G]$. A geodetic set of minimum cardinality is a *minimum geodetic set*. The cardinality of a minimum geodetic set in G is called *geodetic number* and is denoted by $g(G)$.

3) *Example 1.1.3.* consider the following graph



Here $S_1 = \{v_1, v_5\}$ and $S_2 = \{v_1, v_2, v_3, v_5\}$ are geodetic sets.

Hence the Geodetic Number is 2

4) *Definition 1.1.4* The eccentricity $e(u)$ of a vertex u is the distance to a vertex farthest from u . The radius $r(G)$ of G is defined by $r(G) = \min \{e(u): u \in V(G)\}$


B. Geodetic Polynomial

Let $\mathcal{G}(G,i)$ be the family of geodetic sets of the graph G with cardinality I and $\mathcal{G}_e(G,i) = |\mathcal{G}(G,i)|$. Then the geodetic polynomial $\mathcal{G}(G,x)$ of G is defined as

$$\mathcal{G}(G,x) = \sum_{i=\mathcal{G}(G)}^{|V(G)|} \mathcal{G}_e(G,i)x^i$$

Where $\mathcal{G}(G)$ is the geodetic number of G .

1) *Example 1.2.1:* Consider The Following Graph.



$$d(v_1, v_5) = 3; v_1 - v_2 - v_4 - v_5, v_1 - v_3 - v_4 - v_5, I(u,v) = \{v_1, v_2, v_3, v_4, v_5\}$$

$$|\mathcal{G}_e(G,2)| = 1, |\mathcal{G}_e(G,3)| = 2, |\mathcal{G}_e(G,4)| = 3, |\mathcal{G}_e(G,5)| = 1$$


$$\mathcal{G}(G,x) = \sum_{i=\mathcal{G}(G)}^{|V(G)|} \mathcal{G}_e(G,i)x^i = \sum_2^5 \mathcal{G}_e(G,i)x^i$$

$$\mathcal{G}(G,x) = x^2 + 2x^3 + 3x^4 + x^5$$

2) *Geodetic Polynomial of Complete Graph*

3) *Result 1.3.1.[5]* $\mathcal{G}(K_n,x) = x^n$ where K_n is the complete graph.

4) *Example 1.3.2.* Consider the graph K_5




$$\mathcal{G}(K_5, x) = \sum \mathcal{G}_e(K_5, i)x^i = 1.x^5 = x^5$$

C. Geodetic Polynomial of complete bipartite Graph

1) *Result 1.3.1[5]:* $\mathcal{G}(K_{m,n}, x) = x^n(1+x)^m + x^m(1+x)^n - x^{m+n}, m \geq 2, n \leq 4$

2) *Example 1.3.2.* Consider the graph $K_{2,3}$



$$\mathcal{G}(K_{2,3}, x) = 1.x^2 + 4.x^3 + 5.x^4 + x^5$$


3) *Result 1.3.2[5]*

The geodetic polynomial of $K_{n,n}$ is

a) $\mathcal{G}(K_{n,n}, x) = x^n(2(1+x)n - x^n), 2 \leq n \leq 4.$

b) $\mathcal{G}(K_{n,n}, x) = ((1+x)^n - 1)^2 - 2nx(1+x)^n + 2x^n(1+nx) + 2nx + n^2x^2, n > 4.$

4) *Example 1.3.3* consider the graph $K_{3,3}$



$$\mathcal{G}(K_{3,3}, x) = 2x^3 + 6x^4 + 6x^5 + x^6$$

5) *Result 1.3.4[5]*

$$\mathcal{G}(K_{1,n}, x) = x^n + x^{n+1}, n > 1$$

From the Figure, it is easy to observe that $\mathcal{G}(G_1, x) = (x^2 + x^3)$, $\mathcal{G}(G_2, x) = x^3$

$$\mathcal{G}(G, x) = (x^2 + x^3).x^3$$


$$\mathcal{G}(G, x) = x^5 + x^6.$$

D. *Geodetic Polynomial of Path*

1) *Result 1.4.1[5]*: Let P_n be a path with n vertices. The geodetic polynomial of P_n is

$$\mathcal{G}(P_n, x) = x^2(1 + x)^{n-2}.$$

2) *Example 1.4.2*



$$\mathcal{G}(P_4, x) = x^2 + 2x^3 + x^4.$$

II. GEODETIC POLYNOMIAL OF RADIAL GRAPHS


In this section we find geodetic polynomial of some radial graphs

A. *Definition 2.1*

Two vertices of a graph are said to be radial to each other if the distance between them is equal to the radius of the graph.

B. *Definition 2.2*

The radial graph of a graph G , denoted by $R(G)$ has the vertex set as in G and two vertices are adjacent in $R(G)$ if and only if they are radial in G . consider the following graph




$$e(u) : e(v_1) = 3, e(v_2) = 2, e(v_3) = 2, e(v_4) = 2, e(v_5) = 3$$

The following vertex pairs are radial to each other.

$$(v_1, v_4), (v_2, v_3), (v_2, v_5), (v_3, v_5)$$

Radial graph :



C. Result 2.3[4]

Let P_n be any path on $n \geq 5$ vertices then

$$R(P_n) = \begin{cases} \binom{n}{2} K_2 & \text{if } n \text{ is even} \\ P_3 \cup \binom{n-3}{2} K_2 & \text{if } n \text{ is odd} \end{cases}$$

D. Result 2.5[4] $R(K_{m,n}) = K_m \cup K_n$

E. Result 2.6[4]

Let C_n be any cycle on $n \geq 4$ vertices then

$$R(C_n) = \begin{cases} \binom{n}{2} K_2 & \text{if } n \text{ is even} \\ \cong C_n & \text{if } n \text{ is odd} \end{cases}$$

F. Result 2.8 [4] Every path P_n , $n \neq 4$ is a radial graph, (i.e) $R(\bar{P}_n) = P_n$

G. Theorem 2.9

$$\mathcal{G}(R(P_n), x) = x^n \text{ if } n \text{ is even, } n \geq 5$$

$$\mathcal{G}(R(P_n), x) = x^{n-1} + x^n \text{ if } n \text{ is odd, } n \geq 5$$

Proof:

$$\text{Since } R(P_n) = \binom{n}{2} K_2, n \text{ is even and } n \geq 5 \text{ and } \mathcal{G}(K_2) = x^2$$

Therefore the geodetic polynomial of $R(P_n)$ is x^n

$G(R(P_n), x)$ is x^n , if n is even and $n \geq 5$.

Since $R(P_n) = P_3 \cup \left(\frac{n-3}{2}\right) K_2$, If n is odd, $n \geq 5$ and the result 2.3[3]

the geodetic polynomial of $R(K_n)$ is

$$G(R(P_n), x) = G(P_3) \cdot \left\{ G(K_2, x) \cdot G(K_2, x) \dots \dots \frac{n-3}{2} \text{ times} \right\}$$


$$= (x^2 + x^3) \cdot x^2 \cdot x^2 \cdot \dots \dots \frac{n-3}{2} \text{ times}$$

$$G(R(P_n), x) = (x^2 + x^3) \cdot x^{2\left(\frac{n-3}{2}\right)}$$

$$= (x^2 + x^3) \cdot x^{n-3}$$

$$G(R(P_n), x) = x^{n-1} + x^n$$

H. Example 2.10



$$G(G_1, x) = x^2 + x^3, G(G_2, x) = x^2, G(G_3, x) = x^2$$

$$G(R(P_n), x) = (x^2 + x^3) \cdot x^2 \cdot x^2$$

$$G(R(P_n), x) = x^6 + x^7$$

I. Theorem 2.11

$$G(R(K_{m,n}), x) = x^{m+n}$$


Proof:

Since $R(K_{m,n}) = K_m \cup K_n$ by the result 2.5[3] the geodetic polynomial of $R(K_{m,n})$

$$G(R(K_{m,n}), x) = G((K_m \cup K_n), x)$$

$$= G(K_m, x) \cdot G(K_n, x) = x^m \cdot x^n = x^{m+n}$$

J. Example 2.12 Consider the graph $R(K_{2,3})$



$$G(G_1, x) = x^2, G(G_2, x) = x^3$$

$$G(R(K_{2,3}), x) = x^2 \cdot x^3 = x^5$$

III. DETOUR GEODETIC POLYNOMIAL OF GRAPHS


In this section we find detour geodesic polynomial of some graphs

A. Definition 3.1

Let G be a connected graph. The length of a longest u - v path between two vertices u and v in G is called the detour distance $D(u, v)$ between u and v . A u - v path of length $D(u, v)$ is a u - v detour.

B. Example 3.2

For a graph G given in the following figure, $D(u, v) = 4$.



C. Definition 3.3

A vertex w is said to lie on a u - v path P if w is a vertex of P . The closed interval $I_D[u, v]$ consists of u , v and all vertices lying in some u - v detour of G . For any $S \subseteq V$, $I_D[S] = \bigcup_{u, v \in S} I_D[u, v]$.

D. Definition 3.4


A set S of vertices is called a detour geodesic set of G if $I_D[S] = V$. The minimum cardinality of a detour geodesic set is the detour number and is denoted by $d_g(G)$. A detour set of minimum cardinality $d_g(G)$ is called a minimum detour set of G or detour geodesic basis of G . Equivalently, a set S of vertices is called a detour geodesic set of G if every vertex of G lies on some detour joining a pair of vertices of G .

E. Definition 3.5

The upper detour geodesic number $dg^+(G)$ of G is the maximum cardinality of a minimum detour geodesic set of G .

F. Example 3.6

For the graph G given below



$S = \{x, u\}$, $S_1 = \{x, v\}$, $S_2 = \{x, y\}$ are three minimum detour geodesic set so that $d_g(G) = 2$.

G. Definition 3.7

Let $D_i(G)$ be the family of detour geodesic sets of the graph G with cardinality i and let $D_e(G, i) = |D_i(G)|$. Then the detour geodesic polynomial $D(G, x)$ of G is defined as

$$D(G, x) = \sum_{i=d_g(G)}^{dg^+(G)} D_e(G, i)x^i$$


Where $d_g(G)$ is the Detour number of G .

H. Example 3.8

For the graph G , given in the above Figure $S = \{x, u\}$, $S1 = \{x, v\}$, $S2 = \{x, y\}$, $dg(G) = dg+(G)$ the detour geodetic polynomial of the graph G is Defined by $D_{\square}(\square, x) = \sum_{\square=2}^{\square} \square e(G, i) \cdot \square^{\square} = D_{\square}(\square, x) = 3x^2$.

I. Example 3.9

For the Graph G , Given in the Figure



$$D(u,v) = 4$$

IV. DETOUR GEODETIC POLYNOMIAL OF RADIAL GRAPH

In this section we find detour geodetic polynomial of some radial graphs

A. Theorem 4.1

Detour Geodetic Polynomial of $R(P_n)$ is \square^{\square}

(i.e) $D_{\square}(R(P_n), x) = \square^{\square}$ if n is even

Proof $D_{\square}(R(P_n), x) = D_{\square}(\left(\frac{\square}{2}\right) K_2, x)$
 $= D_{\square}(K_2, x) \cdot D_{\square}(K_2, x) \cdot D_{\square}(K_2, x) \dots \dots \left(\frac{\square}{2}\right) \text{times}$
 $= \square^2 \cdot \square^2 \cdot \square^2 \dots \dots \left(\frac{\square}{2}\right) \text{times} = (\square^2)^{\frac{\square}{2}} D_{\square}(R(P_n), x) = \square^{\square}$.

B. Theorem 4.2

Detour geodetic polynomial of $R(P_n)$ is $D_{\square}(R(P_n), x) = x^{n-1}$ if n is odd

1) Proof : $D_{\square}(R(P_n), x) = D_{\square}(P_3, x) \cdot \{D_{\square}(\left(\frac{(n-3)}{2}\right) K_2, x)\}$

$$= D_{\square}(P_3, x) \cdot \{D_{\square}(\left(\frac{(n-3)}{2}\right) K_2, x)\} \dots \dots \frac{(n-3)}{2} \text{times.}$$

$$= x^2 \cdot x^2 \cdot x^2 \dots \dots \frac{(n-3)}{2} \text{times.} = x^{2(n/2)} \cdot (x^{2(n-3)/2}) = x^{2+n-3} = x^{n-1}$$

C. Theorem 4.3

Detour geodetic Polynomial of $R(\bar{P}_n)$ is $D_{\square}(R(\bar{P}_n), x) = x^2$

1) Proof : Since $D_{\square}(R(\bar{P}_n), x) = D_{\square}(\bar{P}_n, x)$,

And P_n is a tree with 2 end vertices, $d_g(G) = d_g+(G) = 2$ and

There is only one detour set with cardinality 2

$$D_{\square}(\bar{P}_n, x) = x^2$$

Hence $D(R(\bar{P}_n), x) = x^2$.

D. Theorem 4.4

The detour geodetic polynomial of $R(K_{m,n})$ is $D(R(K_{m,n}), x) = (mx^{m-1} + x^m) \cdot (nx^{n-1} + x^n)$

1) *Proof* : Since $R(K_{m,n}) = K_m \cup K_n$

and $D(R(K_{m,n}), x) = D(K_m, x) \cdot D(K_n, x)$

Now $D(K_m, x) = mx^{m-1} + x^m$, $D(K_n, x) = nx^{n-1} + x^n$

Hence $D(R(K_{m,n}), x) = (mx^{m-1} + x^m) \cdot (nx^{n-1} + x^n)$.

V. CONCLUSION

Here Geodetic polynomial of path, complete graph, complete bipartite graph, radial graphs and detour geodetic polynomial of complete, complete bipartite and radial graphs have been studied. Further we can find the detour geodetic polynomial of other important graphs.

REFERENCES

- [1] Buckley. F and Hararay, F. "Distance in Graphs", Addison Wesley, Reading (1990)
- [2] Chartrand, G, and Zhang. P. "Distance in graphs". In Handbook of graph Theory(ed.by J.Gross and J.Yellen),CRC Press,Bocaron,FL(2004) 873-888
- [3] Chartrand, G, Harary. F and Zhang. P. (2000) "Geodetic Sets in Graphs", Discuss. Math.Graph Theory, 20 : 129-13
- [4] Kathiresan. K.M and G. Marimuthu, "A study on radial graphs", ARS COMBINATORIA 96(2010) PP 353-36
- [5] Vijayan. A and T. Binuselin, "An Introduction to Geodetic Polynomial of a Graph", Bulletin Pure and Applied Sciences, Volume 31E (Math and Stat.) Issue (No.1) 2012 : P 25-32.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)