



# IJRASET

International Journal For Research in  
Applied Science and Engineering Technology



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

**Volume:** TPAM-2018 **Issue:** conference **Month of publication:** March 2018

**DOI:**

[www.ijraset.com](http://www.ijraset.com)

Call:  08813907089

E-mail ID: [ijraset@gmail.com](mailto:ijraset@gmail.com)

# Contra (gsp)\*-Continuous Function in Topological Spaces

Kalpana.M<sup>1</sup>,

<sup>1</sup>Assistant Professor, St.Joseph's College Of Arts And Science College For Women, Hosur.

**Abstract:** In this paper we have introduced a new function of contra (gsp)\*-continuous in topological spaces which is properly placed in between the class of closed sets and gsp-closed sets.

**Key Words:** contra g-continuous, contra gs-continuous,  $\alpha$ g-continuous, contra rg-continuous.

## I. INTRODUCTION

Levine [10] introduced the class of  $g$ -closed sets in 1970. Maki.et.al [12] defined  $\alpha g$ -closed sets in 1994. Arya and Tour [3] defined  $gs$ -closed sets in 1990. Dontchev [8], Gnanambal [9] Palaniappan and Rao[17] introduced gsp-closed set,  $rg$ -closed sets respectively. Veerakumar [18] introduced  $g^*$ -closed sets in 1991. J.Dontchev [8] introduced gsp-closed sets in 1995. The purpose of this paper is to introduce the concepts contra (gsp)\*-continuous function.

## II. PRELIMINARIES

A. *Definition 2.1:* A subset  $A$  of topological space  $(X, \tau)$  is called

- 1) a pre-open set [14] if  $A \subseteq \text{int}(cl(A))$  and a pre-closed set if  $cl(\text{int}(A)) \subseteq A$
- 2) a semi-open set [11] if  $A \subseteq cl(\text{int}(A))$  and a semi-closed set if  $\text{int}(cl(A)) \subseteq A$
- 3) a semi-preopen set [1] if  $A \subseteq cl(\text{int}(cl(A)))$  and a semi-preclosed set [1] if
- 4) an  $\alpha$ -open set [15] if  $A \subseteq \text{int}(cl(\text{int}(A)))$  and an  $\alpha$ -closed set [15] if  $cl(\text{int}(cl(A))) \subseteq A$
- 5) a *regular-open* set [14] if  $\text{int}cl(A)=A$  and an *regular-closed* set [14] if  $A = \text{int}cl(A)$

B. *Definition 2.2:* A subset  $A$  of topological space  $(X, \tau)$  is called

- 1) a generalized closed set (briefly  $g$ -closed) [10] if  $cl(A) \subseteq U$  whenever  $A \subseteq U$
- 2) generalized semi-closed set (briefly  $gs$ -closed) [3] if  $scl(A) \subseteq U$  whenever  $A \subseteq U$  and  $U$  is open in  $(X, \tau)$ .
- 3) an  $\alpha$ -generalized closed
- 4) set (briefly  $\alpha g$ -closed) [12] if  $\alpha cl(A) \subseteq U$  whenever  $A \subseteq U$  and  $U$  is open in  $(X, \tau)$
- 5) a regular generalized closed set (briefly  $rg$ -closed) [17] if  $spcl(A) \subseteq U$  whenever  $A \subseteq U$  and  $U$  is regular open in  $(X, \tau)$

C. *Definition 2.3:* A function  $f: (X, \tau) \rightarrow (Y, \sigma)$  is called

- 1) Contra  $g$ -continuous [4] if  $f^{-1}(v)$  is a  $g$ -closed set of  $(x, \tau)$  for every open set  $v$  of  $(y, \sigma)$
- 2) Contra  $\alpha g$ -continuous [9] if  $f^{-1}(v)$  is a  $\alpha g$ -closed set of  $(x, \tau)$  for every open set  $v$  of  $(y, \sigma)$
- 3) Contra  $gs$ -continuous [7] if  $f^{-1}(v)$  is a  $gs$ -closed set of  $(x, \tau)$  for every open set  $v$  of  $(y, \sigma)$
- 4) Contra  $rg$ -continuous [17] if  $f^{-1}(v)$  is a  $rg$ -closed set of  $(x, \tau)$  for every open set  $v$  of  $(y, \sigma)$

D. *Definition 2.4 :* A function  $f: (X, \tau) \rightarrow (Y, \sigma)$  from a topological space  $X$  into a topological space  $Y$  is said to be contra – continuous if  $f^1(V)$  is closed in  $X$  for each open set  $V$  of  $Y$ .

*E. Definition 2.5:* A function  $f: (X, \tau) \rightarrow (Y, \sigma)$  is called contra(gsp)\*-continuous if  $f^{-1}(V)$  is (gsp)\*-closed in  $(X, \tau)$  for each open set  $V$  of  $(Y, \sigma)$ .

1) *Theorem:* Let a function  $f: (X, \tau) \rightarrow (Y, \sigma)$  be a map, where  $(X, \tau), (Y, \sigma)$  are spaces then  $f$  is contra(gsp)\*-continuous iff the inverse image of every closed subset of  $(Y, \sigma)$  is gsp-open in  $(X, \tau)$

2) *Proof*

Let  $F$  be a closed subset in  $(Y, \sigma)$ .

Then  $Y-F$  is open in  $(Y, \sigma)$

Since  $f$  is contra(gsp)\*-continuous,  $f^{-1}(Y-F)$  is (gsp)\*-closed.

But  $f^{-1}(Y-F) = X - f^{-1}(F)$

Thus  $f^{-1}(F)$  is gsp open in  $(X, \tau)$

Conversely,

Let  $G$  be an open subset in  $(Y, \sigma)$

Then  $Y-G$  is closed in  $(Y, \sigma)$

Since the inverse image of every closed subset in  $(Y, \sigma)$  is gsp-open in  $(X, \tau)$

$f^{-1}(Y-G)$  is gsp open in  $(X, \tau)$

But  $f^{-1}(Y-G) = X - f^{-1}(G)$

Thus  $f^{-1}(G)$  is (gsp)\*-closed.

Therefore  $f$  is contra(gsp)\*-continuous.

*F. Theorem:* Every contra-continuous function is contra(gsp)\*-continuous.

1) *Proof:* Let  $f: (X, \tau) \rightarrow (Y, \sigma)$  be contra-continuous.

Let  $V$  be any open set in  $Y$ .

Then the inverse image  $f^{-1}(V)$  is closed in  $X$ .

Since every closed set is (gsp)\*-closed.

$f^{-1}(V)$  is (gsp)\*-closed in  $X$ .

Therefore  $f$  is contra(gsp)\*-continuous.

*G. Theorem:* Every contra(gsp)\*-continuous map is contra g-continuous.

But the converse is not true.

1) *Proof*

Let  $f: (X, \tau) \rightarrow (Y, \sigma)$  be contra(gsp)\*-continuous.

Let  $V$  be any open set in  $Y$ .

Then the inverse image  $f^{-1}(V)$  is (gsp)\*-closed in  $X$ .

Since every gsp\*-closed set is g-closed.  $f^{-1}(V)$  is g-closed in  $X$ . hence  $f$  is contra g-continuous.

2) *Example*

Let  $X=Y = \{a, b, c\}$

$\tau = \{\emptyset, X, \{a\}, \{a, b\}\}$

$\sigma = \{\emptyset, Y, \{b\}\}$  Let  $f: (X, \tau) \rightarrow (Y, \sigma)$  be the identity map. Let us prove that  $f$  is contra g-continuous.

But not contra(gsp)\*-continuous.

We have proved that the g-closed sets are

$\emptyset, X, \{c\}, \{a, c\}, \{b, c\}$

And the gsp\*-closed sets are

$\emptyset, X, \{c\}, \{b, c\}$

$f^{-1}(\{a, c\}) = \{a, c\}$  is g-closed in  $(X, \tau)$

Thus the inverse of every closed set of  $(Y, \sigma)$  is g-closed in  $(X, \tau)$  but not (gsp)\*-closed in  $(X, \tau)$ .

Hence  $f$  is contra g-continuous but not contra(gsp)\*-continuous.

*H. Theorem:* Every contra(gsp)\*-continuous map is contra  $\alpha$ g-continuous.

1) *Proof*

Let  $f: (X, \tau) \rightarrow (Y, \sigma)$  be contra(gsp)\*-continuous.

To prove that,  $f$  is contra  $\alpha g$  -continuous.

$V$  be any open set in  $Y$ . Then the inverse image  $f^{-1}(V)$  is  $(gsp)^*$ -closed in  $X$ . Since every  $(gsp)^*$ -closed set is  $\alpha g$  -closed.  $f^{-1}(V)$  is  $\alpha g$  -closed in  $X$ .

Therefore  $f$  is contra  $\alpha g$  -continuous.

2) *Example*

Let  $X=Y= \{a, b, c\}$

$\tau= \{\varphi, X, \{a\}, \{a,b\}\}$

$\sigma =\{ \phi, Y, \{b\}\}$

Let  $f: (X,\tau) \rightarrow (Y,\sigma)$  be the identity map.

Let us prove that  $f$  is contra  $\alpha g$  -continuous

But not contra  $(gsp)^*$ -continuous We have proved that the  $\alpha g$  -closed sets are  $\varphi, X, \{b\}, \{c\}, \{a,c\}, \{b,c\}, \{a,c\}, \{b,c\}$

$f^{-1}(\{a,c\})=\{a,c\}$  is  $\alpha g$  -closed in  $(X,\tau)$  but not  $(gsp)^*$ -closed in  $(X,\tau)$ . Hence  $f$  is contra  $\alpha g$ -continuous but not contra  $(gsp)^*$ -continuous.

## REFERENCES

- [1] D. Andrijevic, semi- preopen sets, Mat. Vesnik, 38(1)(1986),24-32.
- [2] I. Arokiarani, K. Balachandran and J. Dontchev, some characterizations of  $gp$ -irresolute and  $gp$ -continuous maps between topological spaces, Mem. Fac. Sci. Kochi. Univ. Ser. A. Math., 20 (1999), 93-104
- [3] S.P. Arya and T. Nour, Characterization of  $s$ -normal spaces, Indian J.Pure.Appl.Math.,21 (1990),717-719.
- [4] K. Balachandran, P. Sundaram and H. Maki, On generalized continuous maps in topological spaces, Mem. Fac. Kochi. Univ. Ser.A.Math.,12 (1991),5-13.
- [5] R. Devi, K. Balachandran and H. Maki, generalized  $\alpha$ -closed maps and  $\alpha$ -generalized closed maps, Indian J.Pure.Appl.Math.,29(1)(1998),37-49.
- [6] R. Devi, H. Maki and K. Balachandran, Semi-generalized closed maps and generalized closed maps, Mem.Fac.Sci.Kochi.Univ.Ser.A.Math.,14(1993),41-5
- [7] R. Devi, H. Maki and K. Balachandran, Semi-generalized homeomorphisms and generalized semi-homeomorphisms in topological spaces, Indian J.Pure.Appl.Math.,26(3)(1995),271-284
- [8] J. Dontchev, on generalizing semi-preopen sets, Mem. Fac. Sci. Kochi. Ser. A, Math., 16 (1995), 35-48.
- [9] Y. Gnanambal, on generalized preregular closed sets in topological spaces, Indian J. Pure. Appl. Math., 28 (3)(1997),351-360
- [10] N. Levine, generalized closed sets in topology, Rend. Circ. Math. Palermo, 19 (2) (1970), 89-96.
- [11] N. Levine, semi-open sets and semi-continuity in topological spaces, Amer. Math. Monthly, 70 (1963), 36-41
- [12] H. Maki, R. Devi and K. Balachandran, Associated topologies of generalized  $\alpha$ -closed sets and  $\alpha$ -generalized closed sets, Mem. Fac. Sci. Kochi Univ. Ser. A, Math., 15 (1994), 51-63.
- [13] H. Maki, J. Umehara and T. Noiri, Every topological spaces is pre- $T_{1/2}$ , Mem. Fac. Sci. Kochi. Univ. Ser. A, Math., 17 (1996), 33-42
- [14] A.S. Mashhour, M.E. Abd EI-Monsef and S.N.EI-Deeb, on pre-continuous and weak pre-continuous mappings proc.Math. and Phys.Soc.Egypt,53(1982),47-53.
- [15] O. Njastad, on some classes of nearly open sets, pacific J.Math.,15(1965),961-970.
- [16] N. Nagaveni, studies on generalizations of homeomorphisms in topological spaces, Ph.D, thesis, Bharathiar University, Coimbatore, 1999.
- [17] N. Palaniappan and K.C. Rao, Regular generalized closed sets, Kyungpook Math.J.,33(2)(1993),211-219.
- [18] M.K.R.S. Veerakumar, Between closed sets and  $g$ -closed sets, Mem. Fac. Sci. Kochi. Univ. Ser.A, Math., 17 (1996), 33-42.



10.22214/IJRASET



45.98



IMPACT FACTOR:  
7.129



IMPACT FACTOR:  
7.429



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24\*7 Support on Whatsapp)