



# IJRASET

International Journal For Research in  
Applied Science and Engineering Technology



---

# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

---

**Volume:** TPAM-2018 **Issue:** conference **Month of publication:** March 2018

**DOI:**

[www.ijraset.com](http://www.ijraset.com)

Call:  08813907089

E-mail ID: [ijraset@gmail.com](mailto:ijraset@gmail.com)

# Hausdorff Dimension on Fuzzy Space

S. Pavana Swetha<sup>1</sup>

<sup>1</sup>Assistant Professor Department of Mathematics St. Joseph's College of Arts and Science for Women- Hosur.

**Abstract:** Fuzzy mathematics forms a branch of mathematics related to fuzzy set theory and fuzzy logic. Hausdorff dimension is a concept in measure theory. It serves as a measure of a local size of a space taking into account the distance between its points. In Hausdorff dimension distance between all members of that set are defined and dimension drawn from the real number  $R$ . In this paper we study about the Hausdorff dimension is drawn from fuzzy space. Also we discuss Frostman's lemma on fuzzy space.

**Keywords:** Fuzzy space, Fuzzy logic, Fuzzy Hausdorff function, Hausdorff dimension, level-n dyadic cubes, Frostman's lemma.

## I. INTRODUCTION

Fuzzy sets can be regarded as a generalization of characteristic functions taking values between 0 and 1 (including 0 and 1) and the characteristic function on a set  $X$  is the constant mapping taking the whole of  $X$  to 1. The point of  $X$  posses the same statuses relative to the characteristic function on  $X$ , (i.e.), are not distinguishable by characteristic function on  $X$  while they are distinguishable by a fuzzy set on  $X$ . Hausdorff dimension is a concept in mathematics introduced in 1918 by mathematician Felix Hausdorff. The dimension generalizes the notion of the dimension of a real vector space. That is, the Hausdorff dimension of an  $n$ -dimensional inner product space equals  $n$ . Hausdorff dimension of a single point is zero, of a line is 1, of a square is 2, and of a cube is 3. For set of points that define a smooth shape or a shape that has a small number of corners, the shapes of traditional geometry and science, the Hausdorff dimension is an integer agreeing with a dimension corresponding to its topology. Usually we find the Hausdorff dimension on the metric space or general space. In this paper we find the Hausdorff dimension on the fuzzy space.

## II. PRELIMINARIES

Fuzzy logic provides an inference morphology that enables approximate human reasoning capabilities to be applied to knowledge based systems. The theory of fuzzy logic provides a mathematical strength to capture the uncertainties associated with human cognitive process, such as thinking and reasoning.

A. *Some essential characteristics of fuzzy logic:*

- 1) In fuzzy logic, exact reasoning is viewed as a limiting case of approximate reasoning.
- 2) Everything is a matter of degree.
- 3) Knowledge is interpreted a collection of elastic or equivalently, fuzzy constraints on a collection of variables.
- 4) Inference is viewed as a process of propagation of elastic constraints.
- 5) Anylogic system can be fuzzified.

There are two main characteristics of fuzzy systems that gives them better performance for specific application:

B. *Definition: 1*

Let  $X$  be a non empty set. A fuzzy set  $A$  in  $X$  is characterized by its membership function  $\mu_A: X \rightarrow [0,1]$  and  $\mu_A(X)$  is interpreted as the degree of membership of elements  $x$  in fuzzy set  $A$  for each  $x \in X$ .

C. *Definition: 2*

Let  $A$  and  $B$  be fuzzy set in  $X$ , then

$$A = B \Leftrightarrow \mu_A(x) = \mu_B(x) \quad \forall x \in X.$$

$$A \subseteq B \Leftrightarrow \mu_A(x) \leq \mu_B(x) \quad \forall x \in X.$$

$$C = A \cup B \Leftrightarrow \mu_C(x) = \max(\mu_A(x), \mu_B(x)) \quad \forall x \in X.$$

$$D = A \cap B \Leftrightarrow \mu_D(x) = \min(\mu_A(x), \mu_B(x)) \quad \forall x \in X. \quad E = A' \Leftrightarrow \mu_E(x) = 1 - \mu_A(x), \quad \forall x \in X.$$

$$A = \{A_i | i \in I\} \text{ the union } C = \bigcup_I A_i \text{ and intersection } D = \bigcap_I A_i \text{ defined by}$$

$$\mu_C(x) = \sup\{\mu_{A_i}(x)\} \quad \forall x \in X \text{ and } \mu_D(x) = \inf\{\mu_{A_i}(x)\} \quad \forall x \in X$$

D. Definition: 3

A fuzzy space X is said to be fuzzy Hausdorff if each  $\lambda$  – diagonal  $d(\lambda)$  of X is a fuzzy closed set of  $X \times X$ .

E. Definition: 4

For each integer  $k$  let  $\Delta_k$  be the set of cubes in  $R^n$  of side length  $2^{-k}$  and corners in the set  $2^{-k}Z^n = \{2^{-k}(v_1, v_2, \dots, v_n) : v_j \in Z\}$  and let  $\Delta$  be the union of all the  $\Delta_k$ .

F. Definition: 5

Let  $(X, \rho)$  be a metric space. For any subset  $U \subset X$ , let  $diam U$  denote its diameter, that is  $diam U = \sup\{\rho(x, y) | x, y \in U\}$ , Let S be any subset of X, and  $\delta > 0$  a real number. Define  $H_\delta^d(X) = \inf \{ \sum_{i=1}^\infty (diam U_i)^d : \cup_{i=1}^\infty U_i \supseteq S, diam U_i < \delta \}$

G. Definition: 6

The Hausdorff dimension of X is defined by  $dim_H(X) = \inf\{d \geq 0 : H^d(X) = 0\}$

Where  $H^d(X) = \lim_{\delta \rightarrow 0} H_\delta^d(X)$ .

Equivalently  $dim_H(X)$  may be defined as the infimum of the set of  $d \in [0, \infty)$  such that the d- dimensional Hausdorff measure of x is zer.

H. THEOREM : 6.1

Let X be fuzzy set and A be any subset of X. If  $H^p(A) < 1$ , then  $H^q(A) = 0$  for all  $q > p$ .

$$\begin{aligned} H^q(A) &= \lim_{\delta \rightarrow 0} H_\delta^q(A) \\ &= \lim_{\delta \rightarrow 0} \inf \{ \sum_{\alpha \in I} (diam U_\alpha)^q : \cup_{\alpha \in I} U_\alpha \supseteq A, diam U_\alpha < \delta \} \\ &\leq \lim_{\delta \rightarrow 0} \inf \{ \sum_{\alpha \in I} (diam U_\alpha)^p \delta^{q-p} : \cup_{\alpha \in I} U_\alpha \supseteq A, diam U_\alpha < \delta \} \\ &= \lim_{\delta \rightarrow 0} \delta^{q-p} \inf \{ \sum_{\alpha \in I} (diam U_\alpha)^p : \cup_{\alpha \in I} U_\alpha \supseteq A, diam U_\alpha < \delta \} \\ &= \lim_{\delta \rightarrow 0} \delta^{q-p} H^p(A) \end{aligned}$$

$$H^q(A) = 0 \quad (\text{let } \delta \rightarrow 0). \text{ for all } q > p.$$

Hausdorff dimension of an interval on fuzzy space:

I. THEOREM : 6.2

Let X be the fuzzy space. Let  $A = [a, b]$  show that  $m(A) = b - a$  which leads to  $dim_H(A) = 1$ .

1) Proof: Let X be any fuzzy space and  $A = [a, b]$

First we prove  $m(A) \leq b - a$ .

Let  $\epsilon > 0$  and N be an integer such that  $h = \frac{b-a}{N} < \frac{\epsilon}{2}$ .

Let  $x_i = a + ih$  where  $i = 1, 2, 3 \dots N$ .

We define  $C = \{C_i, i = 1, 2, 3 \dots N\}$  where C is an open cover for A.

Assume  $\eta < b - a$   $C_1 = [a, x_1 + \frac{\eta}{N}]$

$$C_i = [x_i - \frac{\eta}{2N}, x_{i+1} + \frac{\eta}{2N}]$$

$$C_i = [x_{N-1} - \frac{\eta}{N}, b]$$

For each  $i$ ,  $diam(C_i) = h + \frac{\eta}{N} < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$ .

Thus  $m(C) < \epsilon$  and  $\sum diam(C_i) = h + \frac{\eta}{N} + (N-2)(h + \frac{\eta}{N}) + (h + \frac{\eta}{N})$

$$= hN + \eta$$

$$= b - a + \eta$$

$$m(A) \leq b - a + \eta \quad \text{for every } \eta > 0$$

$$\Rightarrow m(A) \leq b - a.$$

Now show that  $m(A) \geq b - a$ .

Let  $C = \{C_j : j \in Z_+\}$  be an open cover of  $A \cap \eta$

Since A is a compact and  $\epsilon > 0$

$$|x - y| < \epsilon \quad \text{for every } x, y \in A$$

Then there exist  $C_i \in \mathcal{C}$  such that  $x, y \in C_i$ .

Let N be a positive integer  $h = \frac{b-a}{N} < \epsilon$ .

Define  $x_i = a + ih$  for  $i = 1, 2, \dots, N$

$$x_i - x_{i-1} = h \leq \text{diam}(C_j(i))$$

Hence  $b - a = \sum_{i=1}^N x_i - x_{i-1} < \sum \text{diam}(C_j(i))$

$$b - a \leq \text{diam}(C_j(i))$$

$$b - a \leq m(A)$$

Thus  $m(A) \geq b - a$ .

Hence  $m(A) = b - a$ .

### J. Elementary Properties of Hausdorff Dimension on Fuzzy Space

It has the following properties

- 1) Let X and Y be any two fuzzy space, If  $X \subset Y$  then  $\dim_H(X) \leq \dim_H(Y)$
- 2) If  $X_i$  is a countable collection of fuzzy sets with  $\dim_H(X_i) \leq d$  then  $\dim_H(\cup_i X_i) \leq d$ .
- 3) If a fuzzy space X is countable then  $\dim_H(X) = 0$ .
- 4) Let X be a fuzzy space, if  $\dim_H(X) = d$  and  $\dim_H(Y) = d'$ ,  
then  $\dim_H(X \times Y) \geq d + d'$ .

Let X be a fuzzy space, if  $f: X \rightarrow f(x)$  is a lipschitz map,

$$\text{then } \dim_H(f(x)) \leq \dim_H(X).$$

### K. Theorem: 6.3 (Frostman's Lemma)

Let X be a fuzzy space. If  $X \subseteq \mathbb{R}^d$  is closed and  $H_\infty^\alpha(X) > 0$ , then there is a regular probability measure supported on X.

1) Proof:

Let X be a fuzzy space and  $X \subseteq \mathbb{R}^d$  is closed.

Without loss of generality we assume that  $X \subseteq [0, 1]^d$ .

Consider X as intersection with each of the level-0dyadic cubes.

$$X = \bigcup_{D \in D_0} X \cap \bar{D}$$

By Hausdorff Property

If  $H_\infty^\alpha(X \cap \bar{D}) = 0$ , for each D in the union then  $H_\infty^\alpha = 0$ .

Thus  $H_\infty^\alpha(X \cap \bar{D}) > 0$  for  $D \in D_0$

By translating X we may assume that  $\bar{D} = [0, 1]^d$ .

Let  $A = [0, 1]^d$  and let  $\Pi: A^N \rightarrow [0, 1]^d$

Denote the map  $\Pi(\omega) = \sum_{i=1}^\infty 2^{-n} \omega_n$ .

For  $d=1$

$$\Pi: \{0,1\}^N \rightarrow [0,1]$$

This map  $\Pi_0$  gives the binary representation.

For  $d > 1$ ,  $\Pi$  maps each components of  $\{0, 1\}^N$  in  $(\{0,1\}^N)^d$ .

Thus  $(\{0,1\}^d)^N \cong (\{0,1\}^N)^d$

Therefore the map  $\Pi$  is onto, since each co-ordinates  $x_i$  in binary representation as

$$x_i = 0.x_{1n}x_{2n}x_{3n} \dots x_{dn} \text{ and set } \omega_n = \{x_{1n}, x_{2n}, x_{3n}, \dots, x_{dn}\}$$

But  $\Pi$  is not one-to-one if x has co-ordinates which are dyadic rational. Because there will be multiple pre-images.

The space  $A^N$  can be given the metric

$$d(\omega, \eta) = 2^{-n} \text{ for } n = \min\{k \geq 0 : \omega_{k+1} \neq \eta_{k+1}\}$$

By lipschitz map  $\Pi$  is compact and continuous. Thus every closed subset  $X \subseteq [0, 1]^d$  transforms into a closed subset  $\Pi^{-1}(X)$  of  $A^N$ .

Conversely every closed subset of  $Y \subseteq A^N$  represents as  $X = \Pi(Y)$  it is closed subset of  $[0, 1]^d$ .

For  $\omega_1, \omega_2, \dots, \omega_n \in A$ , the cylinder set  $[\omega_1, \omega_2, \dots, \omega_n] \subseteq A^N$  is

$$[\omega_1, \omega_2, \dots, \omega_n] = \{\eta \in A^N : \eta_1, \eta_2, \dots, \eta_n = \omega_1, \omega_2, \dots, \omega_n\}$$

Choose arbitrary  $\varepsilon$ , thus  $[\varepsilon] = A^N$ .

The metric  $d$  is defined, so that  $[\omega_1, \omega_2, \dots, \omega_n] = B_{2^{-n}}(\eta)$  for every  $\eta = [\omega_1, \omega_2, \dots, \omega_n]$  and the diameter of this ball is  $2^{-n}$ . so cylinder set is closed.

For each family of sets  $C_n = \{[a] : a \in A^N\}$  forms a finite partition of  $A^N$ .

Complement of  $[a]$  is the union of finitely many closed sets, so cylinder sets are also open.

The image of  $\Pi[\omega_1, \omega_2, \dots, \omega_n]$  is the closure of the dyadic cube  $D \in D_0$  containing

$\sum_{i=1}^n 2^{-i} \omega_i = 0. \omega_1. \omega_2 \dots \omega_n$  which is a set of diameter  $\sqrt{d} \cdot 2^{-n}$ . the pre-image  $\Pi^{-1}(D)$  of any level- $n$  dyadic cell  $D \in D_n$  intersects at most  $2^d$  level- $n$  cylinder sets.

### III. CONCLUSION

Fuzzification of mathematical concepts is based on a generalization of these concepts from characteristic functions to membership functions. Fuzzification is the process of changing a real scalar values into a fuzzy values. In fuzzy space the Hausdorff dimension value is normalized to unity. Fuzzification gives more clarity on finding dimension.

### REFERENCES

- [1] L.A. Zadeh, Fuzzy sets Information and Control 8(1965) 338-353.
- [2] C.K. Wong, Covering properties of Fuzzy topological spaces, Journal of mathematical analysis and application 43, 697-704(1973).
- [3] K.K. Azad, Fuzzy Hausdorff space and Fuzzy perfect mappings, of mathematical analysis and application, 82, 297-305(1981).
- [4] Yuval Kohavi, Hadar Davdovich, Topological dimension, Hausdorff dimension and fractals, May 2006.
- [5] Jimmy Briggs and Tim Tyree, Hausdorff Measure.
- [6] Micheal Hochman, Lecture on fractal geometry and dynamics.
- [7] Dierk Schleicher, Hausdorff dimension, Its properties and its surprises.
- [8] Jay Shah, Hausdorff dimension and its applications.
- [9] Walter Rudin, Real and Complex analysis.
- [10] C. A. Rogers, Hausdorff measure.



10.22214/IJRASET



45.98



IMPACT FACTOR:  
7.129



IMPACT FACTOR:  
7.429



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24\*7 Support on Whatsapp)