



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 6 Issue: IV Month of publication: April 2018

DOI: <http://doi.org/10.22214/ijraset.2018.4713>

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

Colletotrichum Diseases of Forest Nurseries and their Biological Management in vitro using Rhizoplane Mycoflora of Grasses

Mallikarjunaswamy G. E.¹, Bharath Nair²

^{1,2}Department of Forest Pathology, Forest Health Division, KFRI, Thrissur, Kerala, India

Abstract: Microbes are the promising source for the biological management of many plant pathogenic organisms. Biocontrol agents are the viable alternative to chemical fungicides and also reduces developing fungicidal resistance in pathogens. Grasses and their roots are liable sources for microbial activity. The aim of the study was to isolate and identify major saprophytic rhizoplane mycoflora of selected perennial grass species *Alloteropsis cimicina*, *Ischeamum indicum*, *Opplismenus compositus*, *Panicum repens* and *Perotis indica*. Fungal species were isolated from rhizoplane regions by agar plate method in different seasons. Total 55 fungal isolates belongs to six genera, nine species of fungi and two were non-sporulating fungal isolates. The fungal isolates were identified and tested for their hypersensitive activity in sensitive plant species. Thirty six isolates out of 55, exhibited in vitro antagonism to phytopathogenic fungal species – *Colletotrichum gloeosporioides*. The fungus *C. gloeosporioides* is the major foliar pathogen cause anthracnose, leaf spot and blight are the major disease symptoms observed in nurseries. Among fungi tested *Trichoderma harzianum* was found to be more effective antagonistic activity against the pathogen in vitro.

Key words: Perennial grasses, Rhizoplane fungi, *Colletotrichum gloeosporioides*, Biological control, In vitro

I. INTRODUCTION

Forest nurseries established with the aim of providing healthy plant stocks is hindered by a number of factors. Diseases pose a major threat of which fungi forms a major pathogen for the successful production of seedlings and their by hindering planting programmes [1], [2]. Rots, Wilts, damping off and various foliar diseases are the major disease symptoms found in seedlings. Foliar diseases in plants are caused by a number of fungal pathogens among which *Colletotrichum* sp. are the major ones. *Colletotrichum gloeosporioides* Penz. causes anthracnose, leaf spot and leaf blight diseases on a wide variety of plants. *Colletotrichum gloeosporioides* is a common pathogenic fungi found more abundantly distributed in tropical and subtropical regions of the world than in temperate regions. A number of fungicides are available and indiscriminate usage resulting in environmental pollution has led in search of an alternative and eco-friendly methods.

Plant-microbe interactions especially between roots and microbes provide a wide array of opportunities to explore the complexities in association as well as their interaction in the growth and development. Rhizosphere and Rhizoplane regions supports a large number of microorganisms. The role of soil microbes is very complex, they help in nutrient cycling, provide nutrients to plants and stimulate plant growth [3], [4], [5].

Grasses form an important component of ecosystem which keeps rejuvenating in every growing season. They produce fibrous roots which homes abundance of diverse microbes. Rhizosphere and Rhizoplane regions supported populations of fungal communities of anamorphic ascomycetes, teleomorphic ascomycetes, zygomycetes and certain non-sporulating fungi [6]. A great diversity of rhizosphere and rhizoplane microorganisms have been described and also in many cases been used as bio-controlling agents. The present work has been carried out to study the antagonistic activity of rhizoplane fungi against *C. gloeosporioides* for their effectiveness in vitro.

II. MATERIALS AND METHODS

A. Isolation of pathogenic fungi

Disease survey was carried out in Central nurseries of Kerala located at Chettikulam, Kannavam, Kulathupuzha and Nilambur. Infected seedling samples were collected and were brought to the laboratory. The samples were washed thoroughly, blotted and were inoculated on antibiotic amended PDA medium. The associated fungal species were isolated and identified by referring standard manuals [7], [8], [9], [10], [11].

B. Isolation Of Rhizoplane Fungi

Rhizoplane fungi were isolated from roots of grass species *Alloteropsis cimicina*, *Ischeamum indicum*, *Opplismenus compositus*, *Panicum repens* and *Perotis indica*. Root samples were washed thoroughly in slow running tap water, blotted and were fragmented into 1 cm long segments. Root segments were inoculated on antibiotic amended PDA medium. Fungal colonies were isolated and identified by referring to standard manuals as described earlier.

C. Dual Inoculation Of Colletotrichum Gloeosporioides And Potential Antagonist On Pda

Dual inoculation of pathogen and an antagonist was set up. A 5 mm disc of pathogen with similar size of each potential antagonist was taken from the growing edge of five day-old pure culture using a cork borer. The control plates were inoculated for pathogen and antagonists separately. Three replications per treatment were set up for each pathogen and antagonist combinations. Inoculated petri-dishes were incubated at room temperature. Daily growth measurements of fungal colonies were taken for 7 days. The percentage inhibition of radial growth of pathogen was calculated using formula [12].

$$\text{Percentage of Inhibition} = \frac{R_1 - R_2}{R_1} \times 100$$

R₁ – Test organism in Control

R₂ – Test organism in Dual culture

D. Statistical Analysis

Antagonistic ability of fungal isolates were statistically analysed and compared by Duncan's Multiple Range Test (DMRT) using SPSS (ver. 21) software developed by IBM Corporation.

III. RESULTS AND DISCUSSION

Disease survey conducted in the Central Nurseries of Kerala resulted in the observation of seedling diseases, leaf spots and blights. A number of fungal species were isolated and among the isolates *Colletotrichum gloeosporioides* (Fig 1 & 2) was found to be a major pathogen producing symptoms appeared as irregular spot, light to dark brown in colour surrounded by necrotic margin (Table 1) (Fig 3 - 8). Rhizoplane fungi isolated from grasses - *Alloteropsis cimicina*, *Ischeamum indicum*, *Opplismenus compositus*, *Panicum repens* and *Perotis indica* resulted a total of 55 fungal isolates belongs to six genera, nine species of fungi and two were non-sporulating fungal isolates. The fungal isolates were identified and tested for their hypersensitive activity in sensitive plant species (Chilli, Tomato and Tobacco). Thirty six isolates out of 55, exhibited in vitro antagonism to phytopathogenic fungal species – *Colletotrichum gloeosporioides*. The fungal organisms namely *Gliocladium*, *Chaetomium* and *Trichoderma* have been known for their antagonistic activity [13], [14], [15]. The result showed the inhibition percentage of *Trichoderma harzianum* (Fig 3) to be highest (73%-78%). Other tested isolates were seen to be showing moderate antagonistic activity against the pathogen (Table 2) (Fig 9). *Trichoderma* sp. have been known for their antagonism interaction and various species of *Trichoderma* namely *T. koningii*, *T. harzianum* and *T. viride*, respectively have been studied for their antagonistic activity in-vitro [16] [17]. *Colletotrichum* sp. are known to cause various foliar diseases and their management with natural agents have been practised [18], [19], [20], [21], [22]. Biological agents against forest nursery diseases are being practised to some extent. [23] reported that *Trichoderma viride*, *T. harzianum* and *Pseudomonas fluorescens* were effective against damping-off pathogens *Rhizoctonia solani* and *Cylindrocladium quinquesetatum*.




Fig 1 & 2. 1. Pure culture of *Colletotrichum gloeosporioides* & 2. Conidia of *C. gloeosporioides*




Fig 3-8. Leaf spot blight diseases of *Colletotrichum gloeosporioides* of 3. *Cassia fistula*, 4. *Gmelina arborea*, 5. *Swetinia macrophylla*, 6. *Tectona grandis*, 7. *Terminalia bellirica* and 8. *T. arjuna*.

Table 1. Colletotrichum gloeosporioides and associated fungi isolated from various seedlings from Central Nurseries of Kerala

Sl. No.	Tree species	Fungal diseases	Fungal organisms	Central nurseries
1	Acacia auriculiformis	Leaf spot / blight	Cladosporium Cladosporioides Colletotrichum gloeosporioides* Curvularia lunata Myrothecium roridum Pestalotiopsis sp.	KPZ
2	Acacia mangium	Leaf spot / blight	Cladosporium cladosporioides Colletotrichum gloeosporioides* Curvularia lunata Myrothecium roridum Pestalotiopsis sp.	KPZ
3	Aegle marmelos	Leaf spot	Alternaria alternata Colletotrichum gloeosporioides* Phoma sp.	CKM
4	Cassia fistula	Leaf spot	Colletotrichum gloeosporioides* Curvularia lunata Pestalotiopsis sp.	NBR
5	Gmelina arborea	Leaf spot	Colletotrichum gloeosporioides* Curvularia lunata	KNM, NBR, KPZ
6	Pongamia pinnata	Leaf spot	Bipolaris sp. Colletotrichum gloeosporioides*	KNM, NBR
7	Saraca asoka	Leaf spot	Colletotrichum gloeosporioides*	CKM
8	Swetinia macrophylla	Leaf spot / blight	Alternaria alternata Cladosporium cladosporioides Colletotrichum gloeosporioides* Curvularia lunata Pestalotiopsis sp.	KNM, CKM
9	Tectona grandis	Leaf spot / blight	Alternaria alternata Cladosporium cladosporioides Colletotrichum gloeosporioides* Curvularia lunata Pestalotiopsis sp.	KNM, NBR, CKM
10	Terminalia arjuna	Leaf spot	Cladosporium cladosporioides Colletotrichum gloeosporioides*	CKM
11	Terminalia bellirica	Leaf spot	Colletotrichum gloeosporioides* Pestalotiopsis sp.	KNM, KPZ
12	Thespesia populinea	Leaf spot	Colletotrichum gloeosporioides* Fusarium oxysporum	KNM

*Dominating Fungus associated with naturally infected plant materials

CKM- Chettikulam, KNM- Kannavam, KPZ- Kulathupuzha, NBR- Nilambur

Table2. Antagonistic activity of fungal species from rhizoplane regions of grasses against *Colletotrichum gloeosporioides* in vitro

Sl. No.	Fungal organisms	Antagonistic activity of fungal species (%) ¹				
		A. c.	I. i.	O. c.	P. r.	P. i.
1	<i>Aspergillus flavus</i>	0.00±0.00 ² a ³	0.00±0.00 a	0.00±0.00 a	33.45±0.33 e	0.00±0.00 a
2	<i>Aspergillus niger</i>	0.00±0.00 a	30.67±0.56 d	0.00±0.00 a	35.77±0.24 f	39.67±0.34 f
3	<i>Curvularia lunata</i>	26.67±0.45 b	0.00±0.00 a	16.73±0.08 b	0.00±0.00 a	0.00±0.00 a
4	<i>Fusarium oxysporum</i>	0.00±0.00 a	21.43±0.11 c	40.48±0.75 g	21.43±0.14 b	42.67±0.61 g
5	<i>Fusarium</i> sp.	0.00±0.00 a	0.00±0.00 a	26.91±0.08 c	23.81±0.35 c	31.43±0.54 c
6	<i>Penicillium chrysogenum</i>	0.00±0.00 a	32.36±0.77 e	0.00±0.00 a	0.00±0.00 a	0.00±0.00 a
7	<i>Penicillium</i> sp.	29.43±0.74 cd	31.42±1.44 de	38.76±0.38 f	40.06±1.72 g	37.16±1.08 e
8	<i>Phomopsis</i> sp.	30.33±0.73 d	0.00±0.00 a	30.38±1.59 e	40.44±1.23 g	35.76±0.87 d
9	<i>Trichoderma harzianum</i>	73.30±1.35 e	75.78±0.85 f	75.77±0.77 h	77.87±1.30 h	74.76±1.05 h
10	NSF-1	30.13±1.09 d	19.05±1.06 b	28.75±0.40 d	29.76±0.91 d	25.68±1.10 b
11	NSF-2	28.57±1.05 c	19.05±0.63 b	27.75±0.39 cd	0.00±0.00 a	0.00±0.00 a

¹Data is an average of three replicates, ² Standard deviation and ³Experiment was conducted in a factorial design. Means carrying same letters in a row are not significantly different (DMRT, P_{0.05})

A. c. - *Alloteropsis cimicina*, I. i.- *Ischaemum indicum*, O. c.- *Opplismenus compositus*, P. r.- *Panicum repens*, P. i. - *Perotis indica*




Fig 9. Antagonistic activity of *Trichoderma harzianum* against *Colletotrichum gloeosporioides*

IV. CONCLUSION

Shallow fibrous roots of grasses harbours a variety of microbes which are important for plants growth and development and can be possibly applied as an alternative for the management of plant diseases. The desirable demand of plant stock has led to the over dependence on chemicals for time bound supply of materials but the negative effects that they imply has not been stressed so far. The alternative strategy for environmental friendly applications now a days need more serious attention for sustainable management of natural resources. Natural biocontrol agents give an insight into the eco-friendly management of diseases thereby helping in growth and development. Among various natural agents, the fungus *Trichoderma harzianum* proves their antagonistic potential for their effective application as an alternative to the chemical control against a wide set of fungal plant pathogens [24]. Thus, the potentialities of such agents can also applied not only in agricultural field but in forestry sector also.

V. ACKNOWLEDGEMENTS

The authors are grateful to the Director, KFRI and to the field staffs of Kerala Forest Department at various Central Nurseries in the State.

REFERENCES

- [1] B. K. Bakshi. Forest pathology, Principles and practises in forestry, Controller of Publications, Delhi. 400, 1976.
- [2] W. J. Bloomberg. The epidemiology of forest nursery diseases. *Ann. Rev. Phytopathol.*, 23: 83-96, 1985.
- [3] G. Thorn. The fungi in soil. In: *Modern Soil Microbiology* (eds JD van Elsas, JTTrevors, EMH Wellington). New York, Marcel Dekker 63–127, 1997.
- [4] P. Bridge and B.M. Spooner. Soil fungi: diversity and detection. *Plant and Soil* 232, 147–154, 2001.
- [5] F.M. Martin, S. Perotto and P. Bonfante. Mycorrhizal fungi. In: *The Rhizosphere – a fungal community at the interphase between soil and roots* (eds R Pinton, Z Varanini, P Nannipieri). New York, Marcel Dekker 263–296, 2001.
- [6] M. M. Vasanthakumari, G. E. Mallikarjunaswamy and M. B. Shivanna. Root mycoflora of certain grass species of Eragrosteae tribe in forests near Bhadra reservoir of Shimoga dist, Karnataka. Paper presented at 2nd Asian Congress of Mycology and Plant Pathology, Osmania University, Hyderabad, A. P., India, 2007.
- [7] B.C. Sutton. *The Coelomycetes*. CMI, Kew, 1980.
- [8] V. J. A. Arx VJA. The genera of fungi sporulating in pure culture, J. Cramer. In der A.R. Gantner Verlag Kommanditgesellschaft. FL – 9490, Vaduz. 424, 1981.
- [9] C. V. Subramanian. *Hyphomycetes. Taxonomy and Biology*. Academic Press, London, Vol. I and II. 930, 1983.
- [10] P. Ramarao and C. Manoharachary. Soil fungi from Andhra Pradesh. Department of Botany, Osmania University. 164, 1990.
- [11] M. B. Ellis and J. P. Ellis. *Microfungi on land plants an identification handbook*. Croom helm Australia Pvt. Ltd., Sydney, Australia. 818, 2001.
- [12] J. H. Vincent. Distortion of fungal hyphae in the presence of certain inhibitors. *Nature* 15: 850, 1947.
- [13] A. K. Mathew and S. K. Gupta. Biological control of root rot of french bean caused by *Rhizoctonia solani*. *J. Mycol. Pl. Pathol.* 28:202-205, 1998.
- [14] R. D. Prasad, R. Rangeshwaran and S. P. Kumar. Biological control of root and collar rot of sunflower. *J. Mycol. Pl. pathol.*29:184-188, 1999.
- [15] K. K. Pandey, P. K. Pandey and J. P. Upadhyay. Mycoparasitism of *Trichoderma* spp. on *Fusarium* and *Rhizoctonia*. *J. Mycol. Pl. Pathol.* 35:174-176, 2005.
- [16] R. N. Bunker and K. Mathur. Antagonism of local biocontrol agents to *Rhizoctonia solani* inciting dry root rot of chilli. *J. Mycol. Pl. Pathol.*31:50-53, 2001.
- [17] R. Grosch, J. Lottmann, V. N. C. Rehn, K.G. Rehn, L. Mendonca-Hagler, K. Smalla and G. Berg. Analysis of antagonistic interactions between *Trichoderma* isolates from Brazilian weeds and the soil-borne pathogen *Rhizoctonia solani*. *J. Plant Dis. Protec.* 114:167-175, 2007.
- [18] J. W. Klopper, R. Rodriguez-Ubana, G. W. Zehnder, J. F. Murphy, E. Sikora and C. Fernández. Plant root-bacterial interactions in biological control of soil borne diseases and potential extension to systemic and foliar diseases, ASDS Keynote Address Session 3. *Australasian Plant Pathology* 28: 21-26, 1999.
- [19] S. A. Palaniyandi, S. H. Yang, J.H. Cheng, L. Meng and J. W. Suh. Biological control of anthracnose (*Colletotrichum gloeosporioides*) in yam by *Streptomyces* sp. MJM5763. *Journal of Applied Microbiology* 111, 443–455, 2011.
- [20] M. A. Rahman, M. A. Razvy and M. F. Alam. Antagonistic activities of *Trichoderma* strains against chilli anthracnose pathogen. *International Journal of Microbiology and Mycology (IJMM)*, vol. 1, No. 1, 7-22, 2013.
- [21] J. L. S. Heng, U. K. Shah, N. A. A. Rahman, K. Shaari and H. Hamzah. *Streptomyces ambofaciens* S2 - A Potential Biological Control Agent for *Colletotrichum gloeosporioides* the Causal Agent for Anthracnose in Red Chilli Fruits. *J Plant Pathol Microbiol* S1: 006, 2015.
- [22] S. Narmadhavathy, K. Nayar, D. C. V. Rani and M. Ranjith. In vitro screening of nutrients, fungicides and bio-control agents against *Colletotrichum fructicola* causing anthracnose leaf spot of culinary melon. *International Journal of Applied and Pure Science and Agriculture (IJAPSA)* Volume 02, Issue 05, 173-176, 2016.
- [23] C. Mohanan. Biological control of seedling diseases in forest nurseries in Kerala. *J. Biol. Control.* 21(2):189-195, 2007.
- [24] G. E. Harman, C. R. Howell, A. Viterbo, I. Chet .and M. Lorito. *Trichoderma* species – opportunistic, avirulent plant symbionts. *Nature Rev. Microbiol.*, 2:43-56, 2004a.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)