



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 6 Issue: VI Month of publication: June 2018

DOI: <http://doi.org/10.22214/ijraset.2018.6180>

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

Synthesis and Characterization of Novel Compound derived from α -Benzilmonoximehydrazone with 4, 4-dimethylaminobenzaldehyde and its Fe(II), Ni(II) and Pd(II) Complexes

Suryabhan Chaugule¹, Raj Badekar², Pradip Shimpi³, Rama Lokhande⁴

^{1,2,4}School of Basic Sciences, Jaipur National University, Jaipur-302017, INDIA.

³Dept. of Chemistry, K. M. Agrawal College, Kalyan-421301, INDIA.

Abstract: *L* The α -benzilmonoximehydrazone-4, 4-dimethylaminobenzaldehyde (HBMHBB) was prepared in Methanol from the reaction of α -benzilmonoximehydrazone with 4, 4-dimethylaminobenzaldehyde in the presence of hydrochloric acid and also prepared its Fe(II), Ni(II) and Pd(II) complexes. The UV spectrum, PMR spectral data elucidate the structure of synthesized compounds. All prepared compounds are non-electrolyte in nature. On the basis of spectral analysis suggests that Fe(II) and Ni(II) complexes are high spin octahedral and Pd(II) is square planar geometries in environment.

Keywords: α -benzilmonoximehydrazone, 4,4-dimethylaminobenzaldehyde, Ni(II), Pd(II) and Fe(II)

I. INTRODUCTION

Schiff base derivatives attract significant interest and occupy an important role in the development of coordination chemistry. Moreover, Schiff bases complexes containing transition metals have been studied in several research areas such as structural chemistry¹. The Schiff base ligands and their corresponding metal complexes have expanded enormously and include a vast area of organometallic compounds and various aspects of bioinorganic chemistry²⁻⁴. Schiff bases ligands have been reported to show a variety of biological actions by virtue of the azomethine linkage, which is responsible for various antibacterial, antifungal, herbicidal, clinical and analytical activities⁵⁻⁷. Transition metal complexes with nitrogen donor atom of Schiff bases are of particular interest⁸⁻¹⁰, because of their ability to possess unusual configuration¹¹. On the other hand, Schiff bases containing α -benzilmonoxime known to better coordinating ligand¹²⁻¹³. Quite recently, the Schiff base ligand derived from the condensation of α -benzilmonoximehydrazone was previously prepared and characterized¹⁴⁻¹⁷. As a continuation to our work, a new Schiff base such as α -benzilmonoximehydrazone-4,4-dimethylaminobenzaldehyde derived from α -benzilmonoximehydrazone condense with 4,4-dimethylaminobenzaldehyde in the presence of hydrochloric acid and also prepared its Fe(II), Ni(II) and Pd(II) complexes. IUPAC name of the prepared ligand is 4-[(E)-(2E)-[(2E)-2-(hydroxyimino)-1, 2-diphenylethylidene] hydrazinylidene]methyl]-N,N-dimethylaniline, abbreviated as HBMHDAB. The title ligand and its synthesized metal complexes were characterized on the basis of elemental analysis, ¹H NMR, FT(IR), Electronic absorption spectra, Mass and EPR spectra, Magnetic susceptibility measurement.

II. EXPERIMENTAL

All the chemicals used were of AR grade. The solvents were dried and distilled before use according to standard procedure¹⁸. FT(IR) spectra were recorded in KBr medium on a 'Perkin- Elmer spectrum 100'. The pmr spectra recorded on a 'Brucker AV300 NMR Spectrometer' instrument in d₆ DMSO solvent using TMS as internal standard. The UV-visible spectra of ligand and its all metal complexes were recorded on JASCO spectrophotometer V650. Magnetic susceptibility measurements of the complexes in the solid state were determined by Gouy balance using [Ni(en)₃S₂O₃] as callibrants. Molar conductance of the complexes was measured in nitrobenzene at room temperature using Synchronic conductivity bridge type 305.

A. Synthesis of Schiff base Ligand:

α -Benzilmonoximehydrazone was prepared by reported method¹⁹. The title ligand was synthesized by the condensation of α -benzilmonoximehydrazone and 4, 4-dimethylaminobenzaldehyde (1:1 molar ratio) dissolved in ethanol. The resulting reaction

mixture was refluxed for 5h the yellow solid precipitate was obtained through filtered, wash with hot distilled water dried, recrystallised from ethanol. Yield: 78% (M.P. 191°C).

B. Synthesis of Metal Complexes

An ethanolic solution of title ligand (0.02mol) was mixed with aqueous solution of metal (II) chloride (0.01mol), pH was adjusted to 7.5-8.0 by using 0.1N NaOH, colored precipitated was separate out. Filtered and washed with hot distilled water and recrystallized from methanol.

III. RESULTS AND DISCUSSIONS

The analytical data along with some physical properties of the ligand and its various metal complexes are summarized in Table-1. The ligand on interaction with Fe(II), Ni(II) and Pd(II) chlorides, yields complexes corresponding to the general formula ML_2 . The analytical data show that the metal to ligand ratio is 1:2. They are insoluble in water, soluble in common organic solvents. The low molecular conductance value of the complexes reveals their non-electrolytic nature²⁰. High melting points of all metal complexes suggests strong metal-ligand bond.

Table-1: Analytical and physical data for HBMHDAB and its Fe(II), Ni(II), Pd(II) metal complexes

Compound	Color (M.P. in °C)	% Yield	Percentage Expected (Found)					Conductance Ω^{-1}	Magnetic Moment in BM
			C	H	N	O	M		
HBMHDAB	Yellow (191)	79.03	74.59 (73.88)	5.95 (5.98)	15.40 (15.00)	4.32 (4.39)	-	-	-
Fe(BMHDAB) ₂	Blue (238)	83.12	69.53 (67.99)	5.30 (4.81)	14.11 (14.29)	4.03 (4.01)	7.04 (6.98)	1.80	5.17
Ni(BMHDAB) ₂	Brown (243)	79.63	71.71 (70.91)	5.46 (4.99)	14.55 (14.03)	4.16 (4.28)	7.63 (7.62)	2.00	3.08
Pd(BMHDAB) ₂	Brown (260)	79.99	65.40 (64.75)	4.98 (4.92)	13.27 (12.90)	3.79 (3.18)	12.56 (12.02)	1.30	-

A. FT(IR) Spectra

Absence of any band between 3300-3400 cm^{-1} in present ligand, in α -benzilmonoximehydrazone reported¹⁹ at 3287 cm^{-1} for $-NH_2$ group indicating successfully replacement of amino group during condensation. In order to study the binding modes of Schiff base to metal in the complexes, IR spectrum of the free ligand was compared with the spectra of the metal complexes. The band at 3267 cm^{-1} in the ligand was disappeared in all complexes, indicating this hydroxyl group deprotonated during complex formation. The bands at 1645 cm^{-1} and 1745 cm^{-1} due to the azomethine and oximino group respectively of the ligand underwent to Shift to lower frequency (Table-2) after complexation, indicating the coordination of azomethine and oximino nitrogen to metal ion and this can be explained by the donation of electrons from nitrogen to the empty *d*-orbitals of the metal atom. The nature of metal-ligand bonding is confirmed by the newly formed bands at the region 490-555 cm^{-1} in the spectra of the complexes which is tentatively assigned to M-N vibration. On the basis of FT(IR) spectra of ligand and its all complexes, suggested that HBMHDAB ligand coordinated to metal ion through nitrogen atoms only. The formation of metal to ligand is M-N₄ type of composition.

Table: - 2: FT(IR) spectral data for HBMOH and its metal complexes

Compound	$\nu(O-H)$	Ar. C-H	C=NO	C=NN	>CH-	N - O	N-N	M-N
HBMHDAB	3267	3058	1645	1745	1690	1000	1072	-
Fe(BMHDAB) ₂	-	3069	1600	1690	1691	1025	1093	492, 458
Ni(BMHDAB)	-	3025	1603	1710	1695	1017	1099	482, 555
Pd(BMHDAB) ₂	-	3012	1608	1705	1690	1018	1100	515, 490

B. ¹H NMR spectra:

The ¹H NMR spectra of the ligand and its Pd(II) complex done in d₆ DMSO solvent. The ¹H NMR spectrum of the prepared ligand exhibits a multiplet signal at 6.7-8.50δ (m, 14H) suggests that the aromatic groups present in ligand. This band is unshifted in Pd(II) complexes, indicate non-involvement of aromatic group in the coordination and also two methane group observed in the region 1.2-1.6δ are unshifted in complex, suggests they are not involve in coordination. A sharp and singlet band observed at 10.20δ (s, 1H) in title ligand is absent in the palladium (II) complex, confirm the oximino group is deprotonated during complex formation. Other bands are unchanged in complex, suggests non-involvement of these group in coordination.

Table -3: PMR spectral data of the HBMOH

Compound	-OH	Phenyl Rings	N-CH ₃	-CH=
HBMHDAB	10.28	6.70-8.50	1.20-1.60	0.38
Pd(BMHDAB) ₂	-	6.80-8.40	1.20-1.60	0.39

C. Magnetic Moment

Room temperature magnetic susceptibility measurements indicate that the Pd(II) complex is diamagnetic suggesting absence of unpaired electron in this complex as expected for square planar (*d*⁸) geometry of Pd(II) complex. Ink blue colored Fe(BMHDAB)₂ exhibits paramagnetic moment of 5.17 BM at room temperature. The Fe(II) ion shows little tendency to form four coordinate complexes²¹. Most ferrous complexes are found to be octahedral showing either diamagnetism or paramagnetism corresponding to four unpaired electrons. The observed magnetic moment for Fe(BMHDAB)₂ complex, suggests high spin octahedral environment for this complex. The brown colored Ni(BMHDAB)₂ complex shows room temperature magnetic moment of 3.08BM. High spin Ni(II) complexes expected to be paramagnetic properties due to two unpaired electrons. Octahedral Ni(II) complexes have magnetic moments between 3.0-3.3BM, which indicates a small but definite orbital contribution to the magnetic moment. Tetrahedral complexes, however, range from 3.45-4.0BM, the larger the distortion from a regular tetrahedron, the lower the magnetic moment. Square planar Ni(II) complexes usually have no unpaired electrons, are diamagnetic and thus magnetic moment of zero. The observed magnetic moment for Ni(BMHDAB)₂ complex is in the range expected for octahedral Ni(II) complexes.

D. Electronic Absorption Spectra

The electron absorption spectra of the ink blue [Fe(BMHDAB)₂] in chloroform solution shows a symmetrical broad band around 646nm, along with a small band at around 555nm. The spectrum also shows a high intensity band at 380nm. The position of the last band along with the high intensity shown, suggests that it originated from a charge transfer phenomenon. While, the earlier two bands could be said originate from the intra metal transitions. For Fe(II) high spin case, the ground state (⁵D) is supported to split into ⁵E_g and ⁵T_{2g} states, as such are d-d transition corresponding to ⁵T_{2g}→⁵E_g may be expected. However, the ⁵E_g state is expected to further split into ⁵B_{1g} and ⁵A_{1g} owing perhaps, to Jahn-Teller distortion. As such Fe(II) high spin octahedral complexes could show two, fairly closely spectral bands. The electronic spectrum of the light green Ni(II) complex in chloroform solution exhibits a peak at 980nm which assigned to ³A_{2g}→³T_{2g}(ν₁) transition, this value assigned to 10Dq of the Ni(II) complex. Another broad band is observed at 620nm which assigned to the ³A_{2g}→³T_{1g}(F)(ν₂). The ratio ν₂/ν₁, is found to be 1.55, which falls in the range the octahedral complexes of Ni(II)²². From the observed position of these two transitions, the frequency of the third d-d transitions ³A_{2g}→³T_{1g}(P)(ν₃) and B values has been calculated employing the following equations²³.

$$B = [(\nu_2 - 2\nu_1)(\nu_2 - \nu_1)] / (15\nu_2 - 27\nu_1)$$

$$\nu_3 = (225B^2 + 100Dq^2 + 180DqB)^{1/2}$$

The third band, ν₃ is expected to occur at 16160cm⁻¹. However, this band is not observed in the solution spectrum of the complex probably because, it is masked by the tail-end of the strong charge transfer band around 25970cm⁻¹ (ε = 5668 dm³/mol/cm). The value of the Racah interelectronic repulsion parameter B is 0.7420cm⁻¹, calculated with the help of the appropriate equations²⁴. The expression $B = 15B'$ is found to be 11143cm⁻¹ and the ratio $\beta = \frac{B(\text{found})}{B'(\text{free ion})}$, the covalency factor is found to be 0.8216cm⁻¹ and is in close agreement octahedral Ni(II) complexes²⁵.


The green Pd(II) complex in chloroform shows charge transfer bands at 440nm. The ligand field transitions are not clearly observed.

Table -4: Electronic spectral data for HBMHDAB and its metal complexes

No.	Compound	Solvent	Band position in nm	Intensity ϵ	Assignment
1	HBMHDAB	Methanol	339	12252	Oximino $\pi \rightarrow \pi^*$ transition
			249	11181	Azomethine $\pi \rightarrow \pi^*$ transition
		0.1N NaOH	356	12116	Oximino $\pi \rightarrow \pi^*$ transition
			256	11288	Azomethine $\pi \rightarrow \pi^*$ transition
2	Fe(BMHDAB) ₂	Methanol	221	23600	Charge transfer M \rightarrow L transition
			646	271	$^5T_{2g} \rightarrow ^5E_g$ transition
		Chloroform	555	5444	Charge transfer M \rightarrow L transition
			380	27899	Charge transfer M \rightarrow L transition
3	Ni(BMHDAB) ₂	Methanol	328	29100	Charge transfer M \rightarrow L transition
			225	18190	Charge transfer M \rightarrow L transition
		Chloroform	980	10	$^3A_{2g} \rightarrow ^3T_{1g}$ transition
			620	328	$^3A_{2g} \rightarrow ^3T_{1g}(F)$ transition
			523	14371	Charge transfer M \rightarrow L transition
5	Pd(BMHDAB) ₂	Methanol	333	1811	Charge transfer M \rightarrow L transition
		Chloroform	578	1910	Charge transfer M \rightarrow L transition
			449	13530	Charge transfer M \rightarrow L transition

IV. CONCLUSION

In the present paper, coordination chemistry of a Schiff base ligand obtained from the condensation reaction of α -benzilmonoximehydrazone and 4,4-dimethylaminobenzaldehyde is described. Fe(II), Ni(II) and Pd(II) metal complexes have been synthesized using the title schiff base ligand and characterized on the basis of analytical, magnetic and spectral data. The ligand coordinates through its oximino and azomethine nitrogen atoms to the metal ions and acts as a neutral bidentate ligand. All complexes are non-electrolyte, high thermal stability and strong metal-ligand bond. On the basis of spectral and magnetic moment analysis, Fe(II) and Ni(II) complexes are high spin octahedral and Pd(II) is square planar geometry. On the basis of FT(IR) spectra of ligand and its all complexes, suggested that HBMHDAB ligand coordinated to metal ion through nitrogen atoms only. The proposed structure of metal complexes of HBMHDAB follows;



Where M = Fe(II), Ni(II) and Pd(II)

V. ACKNOWLEDGMENTS

We thank to SAIF IIT Bombay for use of instrument facilities.

REFERENCES

- [1] Hine J. and Yeh C. Y.; "Chemical and Biological Studies of Nakiterpiosin and Nakiterpiosinone"; Journal American Chemical Society, 89(1967)1669.
- [2] Spichiger-Keller U.; "Chemical Sensors and Biosensors for Medical and Biological Applications"; Wiley-VCH, Weinheim, (1998).
- [3] Lawrence J. F. and Frei R. W.; "Chemical Derivation in Chromatography"; Elsevier, Amsterdam,(1976).
- [4] Anacona J. R., Bastardo E. and Camus J.; "Manganese (II) and palladium (II) complexes containing a new macrocyclic Schiff base ligand: antibacterial properties"; Transition Metal Chemistry, 24(1999) 478.
- [5] Elmali A., Kabak M. and Elerman Y.; "Synthesis and Antimicrobial Activities of Some New Synthesized Imide and Schiff's Base Derivatives"; Journal Molecular Structure, 477 (2000) 151.
- [6] Anacona J. R., Bastardo E. and Camus J.; "Manganese (II) and palladium (II) complexes containing a new macrocyclic Schiff base ligand: antibacterial properties"; Transition Metal Chemistry, 24(1999) 478.
- [7] Trujillo A., Sinbandhit S., Toupet L., Carrillo D., Manzu C. and Hamon J.R.; "Organometallic-inorganic conjugated unsymmetrical Schiff-base hybrids structures of functionalized; Journal of Inorganic Organometallic Polymer., 18 (2008) 81.
- [8] Clarke B., Clarke N., Cunningham D., Higgins T., McArdle P., Ni Chlolu M. and O'Gara M.; "Transition-metal Schiff-base complexes as ligands in tin chemistry"; Journal of Organometallic Chemistry, 559 (1998) 55.
- [9] Pandeya S. N., Sriram D., Nath G. and De Clercq E.; "Synthesis, antibacterial, antifungal and anti-HIV evaluation of Schiff and Mannich bases of isatin derivatives with 3-amino-2-methylmercapto quinazolin-4 (3H)-one"; Pharma. Acta Helv, 74 (1999) 11.
- [10] Pandeya S. N., Sriram D., Nath G. and De Clercq E.; "Synthesis, antibacterial, antifungal and anti-HIV activities of norfloxacin Mannich bases"; Arzneimittel Forsch., 50 (2000) 11.
- [11] Singh W. M. and Dash B. C.; "Synthesis, characterization and biological studies of Co (II), Ni (II), Cu (II) and Zn (II) complexes with bidentate Schiff bases derived by heterocyclic ketone"; Pesticides, 22 (1988) 33.
- [12] Soleimany E.; "Novel complexes of Mn (II), Co (II), and Cu (II) with ligand derived from dibromobenziloxime"; Journal of the Chinese Chemical Society; 57, (2010), 653- 658.
- [13] Soleimany E.; "Novel complexes of Mn (II), Co (II), and Cu (II) with ligand derived from dibromobenziloxime"; Journal of the Chinese Chemical Society; 57, (2010), 653- 658.
- [14] Badekar R. R.; "Studies on some metal complexes with -(1E,2E)-1,2-diphenylethane-1,2-dione hydrazine oxime" M.Sc. Thesis, University of Mumbai,(2012)
- [15] Badekar R. R., Kulkarni S. W., Lokhande R. S. and Thawkar B. S.; "Synthesis, characterization and antibacterial activity of hydrazone derivative of α -Benzilmonoxime ligand and its Iron (III), chromium (III) and ruthenium (III) metal complexes"; International Journal of Applied Research; 2(9), (2016), 175-179.
- [16] Badekar R. R., Kulkarni S. W., Lokhande R. S. and Patil R. M.; "Synthesis and Characterization of α - Benzilmonoximehydrazone ligand and its Zn(II), Cd(II) and Hg(II) metal complexes"; International Journal of Advanced Research; 4(7), (2016), 1093-1097.
- [17] Badekar R. R., Kulkarni S. W., Lokhande R. S. and Patil R. M.; "Synthesis and Characterization of (1E, 2E)-1, 2-Diphenylethane-1, 2-Diene Hydrazone Oxime ligand and its Fe(II) and Pd(II) metal complexes"; International Journal of Advanced Research; 2(9), (2016), 796-800.
- [18] Perkin D.D., Armarego W.L.F. and Perrin D.R.; "Purification of Lab Chemicals"; 2nd edition, Pergamon Press Ltd (1980).
- [19] Badekar R. R.; "Synthesis, characterization and biological activities of some metal complexes with -(1E,2E)-1,2-diphenylethane-1,2-dione hydrazine oxime" PhD. Thesis, Jaipur National University,(2017)
- [20] Greay W.J.; Coord. Chem. Rev.; 7 (1971) 81.
- [21] Cotton F.A. and Wilkinson; Advanced Inorganic Chemistry; 6th ed., Jahn-Wiley Inc. (1999).
- [22] Drago R.S.; "Physical Methods in Inorganic Chemistry"; EWP, New Delhi; P-135-181, (1965).
- [23] Geary W.J.; "The use of conductivity measurements in organic solvents for the characterization of coordination compounds"; Coordination Chemistry Review; 7 (1971) 81.
- [24] Koing E.; "Structure and Bonding"; V-9 (1971) R175.
- [25] Lewis J and Wilkinson R.J.; "Modern Coordination Chemistry; Interscience, N.Y.; P-427 (1960).



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)