



IJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 6

Issue: X

Month of publication: October 2018

DOI:

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

On Cubic Equation With Four Unknowns

$$4(x^3 + y^3) + 12(s^2 - 1)(x + y)z^2 = (3s^2 + 1)w^3, (s \neq \pm 1)$$

J. Shanthi¹, M.A. Gopalan²

¹Assistant Professor, Department of Mathematics, SIGC, Trichy-620002, Tamilnadu, India.

²Professor, Department of Mathematics, SIGC, Trichy-620002, Tamilnadu, India.

Abstract: In this paper, the cubic equation with four unknown given by $4(x^3 + y^3) + 12(s^2 - 1)(x + y)z^2 = (3s^2 + 1)w^3, (s \neq \pm 1)$ is considered for determining its non-zero distinct integer solutions.

Keywords: Cubic with four unknowns, homogeneous cubic, integer solutions.

I. INTRODUCTION

It is well-known that there are varieties of cubic equations with four unknowns to obtain integer solutions satisfying them [1-3]. In particular, different choices of cubic equations with four unknowns are presented in [4-12]. This paper has a different choice of cubic equation with four unknowns given by $4(x^3 + y^3) + 12(s^2 - 1)(x + y)z^2 = (3s^2 + 1)w^3, (s \neq \pm 1)$ to obtain its infinitely many non-zero distinct integer solutions.

II. METHOD OF ANALYSIS

The cubic equation with four unknowns to be solved for its non-zero distinct integer solutions is given by

$$4(x^3 + y^3) + 12(s^2 - 1)(x + y)z^2 = (3s^2 + 1)w^3 \tag{1}$$

Introduction of the linear transformations

$$x = u + v, y = u - v, w = 2u \quad (u \neq v \neq 0) \tag{2}$$

in (1) leads to

$$v^2 = s^2u^2 - (s^2 - 1)z^2 \tag{3}$$

Again, considering the linear transformations

$$u = X + (s^2 - 1)T \tag{4}$$

$$z = X + s^2T \tag{5}$$

in (3), it gives

$$X^2 = (s^4 - s^2)T^2 + v^2 \tag{6}$$

The fundamental solution of (6) is

$$T_0 = 2v, X_0 = (2s^2 - 1)v$$

To obtain the other solutions of (6), consider its pellian equation

$$X^2 = (s^4 - s^2)T^2 + 1$$

whose general solution $(\tilde{T}_n, \tilde{X}_n)$ is given by

$$\tilde{X}_n = \frac{1}{2}f_n, \tilde{T}_n = \frac{1}{2\sqrt{s^4 - s^2}}g_n$$

where

$$f_n = \left(2s^2 - 1 + 2\sqrt{s^4 - s^2}\right)^{n+1} + \left(2s^2 - 1 - 2\sqrt{s^4 - s^2}\right)^{n+1}$$

$$g_n = \left(2s^2 - 1 + 2\sqrt{s^4 - s^2}\right)^{n+1} - \left(2s^2 - 1 - 2\sqrt{s^4 - s^2}\right)^{n+1}$$

Applying the lemma of Brahmagupta between the solutions (T_0, X_0) and $(\tilde{T}_n, \tilde{X}_n)$, the other solutions to (6) are given by

$$\left. \begin{aligned} T_{n+1} &= v f_n + \frac{(2s^2 - 1)v}{2\sqrt{s^4 - s^2}} g_n, \\ X_{n+1} &= \frac{(2s^2 - 1)v}{2} f_n + v\sqrt{s^4 - s^2} g_n, n = 0, 1, 2, \dots \end{aligned} \right\} \quad (7)$$

Employing (4), (5) and (2), the sequence of solutions to (1) is represented as below:

$$\begin{aligned} x_{n+1} &= \frac{(4s^2 - 3)v}{2} f_n + \frac{(4s^4 - 5s^2 + 1)v}{2\sqrt{s^4 - s^2}} g_n + v, \\ y_{n+1} &= \frac{(4s^2 - 3)v}{2} f_n + \frac{(4s^4 - 5s^2 + 1)v}{2\sqrt{s^4 - s^2}} g_n - v, \\ z_{n+1} &= \frac{(4s^2 - 1)v}{2} f_n + \frac{(4s^4 - 3s^2)v}{2\sqrt{s^4 - s^2}} g_n, \\ w_{n+1} &= (4s^2 - 3)v f_n + \frac{(4s^4 - 5s^2 + 1)v}{\sqrt{s^4 - s^2}} g_n, n = -1, 0, 1, \dots \end{aligned}$$

In addition to the above solutions, there are other choices of solutions to (1) that are illustrated below:

It is worth to note that (6) can be represented as the system of double equations as presented below in Table 1:

TABLE 1
System Of Double Equations

System	1	2	3
$X + v$	$s(s + 1)T$	$s(s^2 - 1)T$	$s^2(s + 1)T$
$X - v$	$s(s - 1)T$	sT	$(s - 1)T$

Solving each of the above systems, one obtains the values to X, v, T .

Substituting these values in (4), (5) and (2), the corresponding solutions to (1) are obtained and they are exhibited below:

Solutions to system 1:

$$x = (2s^2 + s - 1)T, y = (2s^2 - s - 1)T, z = 2s^2T, w = (4s^2 - 2)T$$

Solutions to system 2:

$$x = 2k(s^3 + s^2 - s - 1), y = 2k(s^2 + s - 1), z = k(s^3 + 2s^2), w = 2k(s^3 + 2s^2 - 2)$$

Solutions to system 3:

$$x = k(2s^3 + 4s^2 - 2), y = k(2s^2 + 2s - 4), z = k(s^3 + 3s^2 + s - 1), w = 2k(s^3 + 3s^2 + s - 3)$$

Further, (6) is written as

$$v^2 + (s^4 - s^2)T^2 = X^2 \quad (8)$$

Assume

$$X = s^2A^2 + (s^4 - s^2)s^2B^2 \quad (9)$$

write 1 as

$$1 = \frac{(s + i\sqrt{s^4 - s^2})(s - i\sqrt{s^4 - s^2})}{s^4} \quad (10)$$

Using (9) and (10) in (8) and employing the method of factorization, define

$$v + i\sqrt{s^4 - s^2}T = \left(sA + i\sqrt{s^4 - s^2} sB \right)^2 \frac{\left(s + i\sqrt{s^4 - s^2} \right)}{s^2}$$

On equating the real and imaginary parts, one obtains

$$\left. \begin{aligned} v &= s(A^2 - (s^4 - s^2)B^2) - 2AB(s^4 - s^2) \\ T &= A^2 - (s^4 - s^2)B^2 + 2ABs \end{aligned} \right\} \quad (11)$$

Substituting (9) and (11) in (4), (5) and (2), the corresponding values of x, y, z and w satisfying (1) are given by

$$\begin{aligned} x &= (2s^2 + s - 1)A^2 + (1 - s)(s^4 - s^2)B^2 + 2AB(s^3 - s - s^4 + s^2) \\ y &= (2s^2 - s - 1)A^2 + (1 + s)(s^4 - s^2)B^2 + 2AB(s^3 - s + s^4 - s^2) \\ z &= 2s^2A^2 + 2ABs^3 \\ w &= (4s^2 - 2)A^2 + 2(s^4 - s^2)B^2 + 4s(s^2 - 1)AB \end{aligned}$$

III.CONCLUSIONS

In this paper, an attempt has been made to find non-zero distinct integer solutions to the cubic equation with four unknowns given by $4(x^3 + y^3) + 12(s^2 - 1)(x + y)z^2 = (3s^2 + 1)w^3, (s \neq \pm 1)$ in conclusion one may search for other sets of integer solutions to the considered cubic equation.

REFERENCES

- [1] Dickson. L.E., "History of the Theory of Numbers", Vol 2, Diophantine analysis, New York, Dover, 2005.
- [2] Mordell. L.J., "Diophantine Equations, Academic Press, New York, 1969.
- [3] Carmichael.R.D., "The Theory of numbers and Diophantine Analysis", New York, Dover, 1959.
- [4] Gopalan M.A., and Premalatha S., "Integral Solutions of $(x + y)(xy + w^2) = 2(k^2 + 1)z^3$ ", Bulletin of Pure and Applied Sciences, vol.29E N0.2, Pp.197-202, 2009.
- [5] Gopalan M.A., and Pandichelvi V., "Remarkable solutions of cubic equation with four unknowns $x^3 + y^3 + z^3 = 28(x + y + z)w^2$ ", Antarctica J.Math, vol.7,No.4, Pp.393-401,2010.
- [6] Gopalan M.A., and Sivagami B., "Integral solutions of homogeneous cubic equation with four unknowns $x^3 + y^3 + z^3 = 3xyz + 2(x + y)w^3$ ", Impact J.sci.Tech, Vol.4, No.3, Pp 53-60, 2010.
- [7] Gopalan M.A., and Premalatha S., "On the cubic Diophantine with four unknowns $(x - y)(xy - w^2) = 2(n^2 + 2n)z^3$ ", International Journal of Mathematical Sciences, Vol.9, N0.1-2, Pp.171-175, 2010.
- [8] Gopalan M.A., and Kaligarani J., "Integral solutions of $x^3 + y^3 + (x + y)xy = z^3 + w^3 + (z + w)zw$ ", Bulletin of Pure and Applied Sciences, Vol.29E, No.1, Pp.169-173, 2010.
- [9] Gopalan M.A., and Premalatha S., "Integral solutions of $(x + y)(xy + w^2) = 2(k + 1)z^3$ ", The Global Journal of Applied mathematics and Mathematical Sciences, Vol.3, No.1-2, Pp.51-55, 2010.
- [10] Goplalan M.A., Vidhyalakshmi S., and Mallika S., "Observations on cubic equation with four unknowns $xy + 2z^2 = w^3$ ", The Global Journal of Mathematics and Mathematical Sciences, Vol.2, No.1, Pp.69-74, 2012.
- [11] M.A.Gopalan and K.Geetha, "Observation on Cubic Equation with Four Unknowns $X^3 + Y^3 + XY(X + Y) = Z^3 + 2(X + Y)W^2$ ", International Journal of Pure and Applied Mathematical Sciences, Vol.6, No.1, Pp.25-30, 2013.
- [12] Goplalan M.A., Manju Somanath and Sangeetha V., "Lattice Points on the homogeneous cubic equation with four unknowns $(x + y)(xy + w^2) = (k^2 - 1)z^3, k > 1$ ", Indian Journal of Science, Vol.2, No.4, 97-99, 2013.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)