



IJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 6 Issue: XII Month of publication: December 2018

DOI:

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

Synthesis and Antibacterial Activity of Novel Vanillic Acid Hybrid Derivatives [Part III]

Maruti S. Satpute¹, Vijay D. Gangan², Indu Shastri^{3*},

^{1,3}Department of Chemistry, R. D. National College and W. A. Science College, Linking Road, Bandra (W), Mumbai 400050.

²Department of Chemistry, Loba R & D Centre, Loba Chemie Pvt. Ltd., Plot No. D - 22, Tarapur MIDC, Boisar, Palghar – 401506, India.

Abstract: Vanillic acid (4-hydroxy-3-methoxybenzoic acid) is a dihydroxybenzoic acid derivative used as a flavoring agent. It is used in the synthesis of various active pharmaceutical ingredients such as Etamivan, Modecainide, Brovanexine, Vanitolidide, Vanyldisulfamide etc.

In this paper, novel ester / hybrid derivatives of vanillic acid were synthesized and tested for potential antibacterial activity. This combinatorial synthesis of novel vanillic ester / hybrid derivatives can be a useful approach to generate potent chemotherapeutic agents in developing new drug candidates.

Keywords: Vanillic acid, IR, ¹H NMR, TOF MS, DCC, DMAP, antibacterial, ditch-plate method.

I. INTRODUCTION

Phenolic phytochemicals are known to exhibit anti-inflammatory, antioxidant, anticarcinogenic, antidiabetic, antiatherosclerosis and immunomodulatory activities in animals^{1,2}.

These are mostly polyphenols known as secondary plant metabolites³, present in plants and trees. Polyphenols are commonly divided into flavonoids and the hydroxyl cinnamic acids. Vanillic acid is a naturally occurring active compound having antimicrobial, anti-inflammatory and antioxidant / anticancer properties⁴⁻⁹.

In continuation to our earlier work^{10,11}, we thought of synthesizing compounds with novel ether, ester and hybrid derivatives of Vanillic acid wherein Vanillic acid would be etherified, esterified and hybridized with various other compounds and to check whether these compounds possess above biological activities.

The objective of this study is to condense two molecules of the same disease domain to produce more potent candidate in the same disease domain or to condense two molecules of different disease domain to produce mixed variety of those disease domain or to have drug candidate with entirely different biological activity.

II. MATERIALS AND METHODS

A. Materials

Chemicals used were of a laboratory grade. The reactions were monitored by TLC on aluminium-backed silica plate visualized by UV-light.


B. Experimental

Melting points were determined on a Thomas Hoover capillary melting point apparatus using digital thermometer. IR spectra were recorded on a Shimadzu FTIR Prestige model as KBr pellet. ¹H NMR spectra were recorded on a Varian 400 MHz spectrometer in CDCl₃. Chemical shifts were recorded in parts per million down field from tetramethyl silane. Mass spectra were recorded on a TOF MS ES mass spectrometer. Elemental analysis were carried out as a percentage on a Thermo finnigan, Flash EA 1112 series, Italy.

III. RESULTS AND DISCUSSION

Preparation of 3,4-dialkoxy benzoic acids :- Vanillic acid was subjected to esterification (EtOH / Conc. H₂SO₄) followed by etherification (K₂CO₃ / Acetone / Alkyl halide) to yield crude ether derivatives which were purified by column chromatography. These purified ether derivatives were subjected to hydrolysis (Aq. KOH / EtOH and then Conc. HCl) to yield 3,4-dialkoxy vanillic acids respectively.

Reaction Scheme 1 :




Above 4-alkoxy-3-methoxy benzoic acids were then condensed with ethyl vanillate under DCC/DMAP/Pyridine condition to yield hybrid derivatives whose structures were unambiguously confirmed by IR, $^1\text{H NMR}$, Mass spectroscopy and elemental analysis and tested for their potential antimicrobial activity.

Synthesis of Fused Molecules using compound (1) and (2) :- These were prepared by following general method as depicted below.

To a stirred solution of ethyl vanillate [A] (1 eq.) in 30 ml dichloromethane was added DCC [C] (1.3 eq.), DMAP [D] (0.05 eq.), pyridine [E] (0.5 eq.) and the reaction mixture stirred at room temperature for 5 min. Clear solution of reaction mixture was obtained. To this, compound [B] aromatic / substituted aromatic acid (1.3 eq.) was added and stirring continued at room temperature for next 24 hrs. As the reaction proceeds, urea derivative precipitates out as by product. The progress of the reaction was monitored by TLC for completion of reaction.


Work up :-The reaction mixture filtered through celite bed which get rids of by product urea derivative. The filtrate was concentrated to minimum, preadsorbed on silica gel (100 – 200 mesh) and purified by column chromatography with increase in concentration of ethyl acetate in petroleum ether. The general yields of these reactions ranges between 70 – 90 %. This is another method of preparing esters and follows green chemistry parameters.

Reaction scheme 2 :



Sr. No.	A	B	Product (R)
1	Ethyl vanillate	3-methoxy-4-heptoxy benzoic acid	-C ₇ H ₁₅
2	Ethyl vanillate	Benzoic acid	-

Probable mechanism for fused / hybrid molecules :



Compound 1 :- (4-ethoxycarbonyl-2-methoxy-phenyl) 4-heptoxy-3-methoxy-benzoate

¹H NMR (CDCl₃, 200 MHz) δ ppm : 0.85 (t, J = 7.2 Hz, 3H, terminal methyl from heptyl bromide moiety), 1.4 (t, J = 7.0 Hz, 3H, -CH₃ from -COOCH₂CH₃ group), 1.0 – 1.5 (m, 8H, 4 x -CH₂ from heptyl bromide moiety), 1.7 – 2.0 (m, 2H, 1 x -CH₂ from heptyl bromide moiety), 3.93 (s, 6H, 2 x Ar-OCH₃ group), 4.1 (t, J = 6.6 Hz, 2H, 1 x -OCH₂ from heptyl bromide moiety), 4.4 (q, J = 7.0 Hz, 14 Hz, 2H, -CH₂ from COOCH₂CH₃ group), 6.8 – 7.9 (m, 6H, ArH); TOF MS ES: 467 (M + Na); IR (KBr) cm⁻¹: 2929, 2854, 2873 (methyl, methylenes, methines), 1732 - 1708 (2 x ester carbonyl), 1597 (aromatic); Molecular Formula C₂₅H₃₂O₇; Pale pinkish solid; Melting range 136 – 140°C; Elemental Analysis, Calcd.: C 67.52 %, H 7.28 %, O 25.20 %. Found C 67.49 %, H 7.31 %, O 25.21 %;

Compound 2 :- ethyl 4-benzoyloxy-3-methoxy-benzoate

¹H NMR (CDCl₃, 200 MHz) δ ppm : 1.4 (t, J = 7.0 Hz, 3H, -CH₃ from -COOCH₂CH₃ group), 3.93 (s, 3H, Ar x -OCH₃), 4.4 (q, J = 7.0 Hz, 14 Hz, 2H, -CH₂ from COOCH₂CH₃ group), 6.8 – 8.0 (m, 8H, ArH); TOF MS ES: 323 (M + Na); IR (KBr) cm⁻¹: 2981, 2937, 2850 (Methyl, methines), 1742 - 1708 (2 x ester carbonyl), 1602 (aromatic); Molecular Formula C₁₇H₁₆O₅; Pale reddish solid; Melting range 86 – 90°C; Elemental Analysis, Calcd.: C 68.12 %, H 5.28 %, O 26.60 %. Found C 68.10 %, H 5.31 %, O 26.59 %;

A. Chromatographic System

- 1) Column Chromatography:** For column chromatography 100 – 200 mesh Acme grade silica gel is used. The crude reaction mixture is concentrated under reduced pressure to yield crude mass which is preadsorbed on silica gel and purified by column chromatography with increase in concentration of Ethyl acetate in Petroleum ether. The fractions having similar 'rf' values were pooled together, concentrated and subjected for characterization using various spectroscopic techniques.
- 2) Thin Layer Chromatography:** TLC plates were prepared using silica gel G (ACME, BOMBAY). Pet. ether: EtOAc (85 : 15) was used as the solvent system.
- 3) Radial Chromatography:** The circular glass plates of thickness 1 mm, were prepared by using silica gel (PF254, E. MERCK, 50 g) in cold distilled water (105 ml). For elution, gradually increasing concentrations of EtOAc in pet ether were employed.

B. Biological Activity

Antibacterial Activity using ditch plate method¹² :-

The synthesized molecules were screened for their antibacterial activity at 100 µg/ml concentration using ditch plate method against Gram + ve (Staphylococcus aureus, Corynebacterium diphtheria) and Gram negative (Escherichia coli, Klebsiella pneumonia, Salmonella typhi) bacterial species qualitatively. The results of the antibacterial activities are summarized in Table 1.

- 1) Theory:** Ditch plate method is the method of chosen to test the anti-bacterial activity of compounds. It is a preliminary method to screen the anti-microbial potential of compounds / drugs, which are insoluble or partially soluble in aqueous phase. In this method, the test compound is seeded in an agar plate and the test organisms are streaked across to test the inhibition of the growth as a marker of anti-microbial activity.
- 2) Procedure:** A ditch (10 mm x 70 mm) is cut into sterile MH agar plate. The test drug / compound is added to 5 ml molten MH agar butt at 40°C and this mixture is poured into the ditch and allowed to solidify. The ditch should be made in level with the rest of the agar by pouring the mixture. The different bacterial cultures are streaked perpendicular to the ditch using nichrome wire loop. The plate is then incubated at 37°C for 24 hours. The results are observed as inhibition of bacterial growth on the ditch as well as adjacent to the ditch.

Table 1 : Antibacterial Activity Results

SAMPLE NO.	ACTIVE AGAINST
Base molecule (Ethyl vanillate)	Staphylococcus aureus [Gram positive] Escherichia coli [Gram negative]
1	Staphylococcus aureus [Gram positive] Salmonella typhi [Gram negative] Corynebacterium diphtheriae [Gram positive] Escherichia coli [Gram negative]
2	Staphylococcus aureus [Gram positive] Klebsiella pneumoniae [Gram negative] Corynebacterium diphtheriae [Gram positive] Escherichia coli [Gram negative]

IV. CONCLUSION

The novel hybrid derivatives of vanillic acid were synthesized by cost effective industry viable process following the principle of green chemistry. The synthesis of hybrid derivatives is another way to prepare ester derivatives using DCC as dehydrating agent in a reasonably good yield. The probable mechanism for the formation of hybrid derivative was also discussed.

The biological activity suggest that the base molecule ethyl vanillate have anti-bacterial activity against both the bacterial cultures. Its derivatives *viz.* 1 and 2 were also active against certain Gram + ve and Gram – ve cultures. Thus, fused molecules of vanillic acid (1, 2) having long alkyl side chain were potential antibacterial candidates. In depth analysis of these compounds through structure activity relationship studies would provide further insight and can be an interesting topic of future studies.

The structural diversity and the pronounced biological activities encountered in the vanillic acid derivatives suggests that this class of compounds is worthy for further studies that may lead to derivatives by using combinatorial chemistry approach is an alternative strategy to new therapeutic discovery. In other words the generation of diverse vanillic acid derivatives develop new therapeutic molecules that might result in candidates having better activity.

All synthesized hybrid derivatives were chemically new and confirmed by Scifinder search.

V. ACKNOWLEDGEMENT

One of the authors Mr. Maruti S. Satpute thanks Dr. Indu Shastri, Research Guide and HOD, National College of Science and Commerce for suggesting a problem and constant valuable guidance through out the course and providing laboratory facilities.

REFERENCES

- [1] Wattenberg L. W., Coccia J. B. and Lam L. K. Inhibitory effects of phenolic compounds on benzo[a]pyrene induced neoplasia. *Cancer Res.* 1980, 40 : 2820 – 2823.
- [2] Talalay P, De Long M. J. and Prochaska H. J. Identification of a common chemical signal regulating the induction of enzymes that protect against chemical carcinogens. *Proc. Natl. Acad. Sci.* 1988, 85 : 8261 – 8265.
- [3] Macheix J. J., Fleuriot A. and Billot J. *Fruit phenolics.* CRC press Inc., Boca Raton 1990, FL.
- [4] Natarajan K, Singh S, Burke T. R. Jr., Grunberger D. and Aggarwal B. B. Caffeic acid phenethyl ester is a potent and specific inhibitor of activation of nuclear transcription factor NF-kappa B. *Proc. Natl. Acad. Sci.* 1996, 93 : 9090 – 9095.
- [5] Chiao C., Carothers A. M., Grunberger D., Solomon G., Preston G. A. and Barrett J. C. Apoptosis and altered redox state induced by caffeic acid phenethyl ester (CAPE) in transformed rat fibroblast cells. *Cancer Res.* 1995, 55 : 3576 – 3583.
- [6] Burke T. R. Jr, Fesen M. R., Mazumder A., Wang J., Carothers A. M., Grunberger D., Driscoll J., Kohn K. and Pommier Y. Hydroxylated aromatic inhibitors of HIV-1 integrase. *J. Med. Chem.* 1995, 38 (21) : 4171 – 4178.
- [7] Bose J. S., Gangan V. D., Jain S. K. and Manna S. K. Novel caffeic acid ester derivative induces apoptosis by expressing FasL and downregulating NF-KappaB : Potentiation of cell death mediated by chemotherapeutic agents. *J. Cell Physiol.* 2009, 218 (3) : 653 – 662.
- [8] Bose J. S., Gangan V. D., Jain S. K. and Manna S. K. Downregulation of inflammatory responses by novel caffeine acid ester derivative by inhibiting NF-kappaB. *Journal of Clinical Immunology* 2009, 29, pp. 90 - 98.
- [9] Bose J. S., Gangan V. D. (2007). Applied for US patent on “NOVEL CHEMOTHERAPEUTIC AGENTS AGAINST INFLAMMATION AND CANCER” from Reliance Life Sciences [INDIA 1696/MUM/2006, Publication No. WO/2008/062466, [International Application No. PCT/IN2007/000488].
- [10] Satpute M. S., Shastri, I. and Gangan V. D. (2018); *Rasayan Journal of Chemistry* (In Press).
- [11] Satpute M. S., Shastri, I. and Gangan V. D. (2018); Communicated to *International Journal of Scientific Research in Science and Technology*.
- [12] a) Mwambete K. D. and Lyombe F. (2011). Antimicrobial activity of medicated soaps commonly used by Dar es Salaam residents in Tanzania. *Indian J. Pharm. Sci.* 2011, 73 (1), pp. 92 – 98. b) Al lafi T et. al.. The effect of miswak used in Jordan and Middle East on oral bacteria. *International Dental Journal*, 1995, 45 (3), pp. 218 – 222.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)