



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 7 Issue: III Month of publication: March 2019

DOI: <http://doi.org/10.22214/ijraset.2019.3376>

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

Consequence of Insecticide Cypermethrin on Pollen Viability of *Capsicum annum*, L.; (Solanaceae)

Dr. M. Arul Sheeba Rani¹, Arya K

¹Department of Botany, Nirmala College for Women, Coimbatore- 18

Abstract: Viability has been defined as having the capacity to live, grow, germinate or develop. Pollen is a fine to coarse powdery substance comprising pollen grains which are male macro gametophytes of seed plants, which produce male gamete. The effect of each factor affects pollen viability in a species-specific way, depending on the physiology of the pollen grains and the presence or absence of specific structural modifications. Pollen viability was found to be decreased in several dosages in both recommended and double recommended dosages. Pollen Morphology of these species varies. Shape of the grain found to be useful in spore or pollen identification. The function of the pollen grain in the life cycle of a plant, one way to test pollen viability would be to use the pollen for pollination. The study aims to find the viability by the treatment of insecticide in *Capsicum annum*, L.; it is large, shrubby perennial herb. Chemical that used as plant protectants and atmospheric pollutants affect pollen viability of a large number of crop plants. Cypermethrin is a synthetic pyreniod used as an insecticide in large scale commercial agriculture application.

Keywords: Pollen, Viability, Cypermethrin, *Capsicum annum*.


I. INTRODUCTION

The study of pollen and spores is the subject matter of Palynology. The terms viability, stainability, vigour, germinability, fertility, and fertilization ability indicate different aspects of the pollen potential. Palynology is a study of the various aspects of pollen grains and spores. Pollen viability is important for growers and breeders. Pollen viability assessment is critical in the study of the following aspects: monitoring pollen state during storage; genetics and pollen-stigma interactions; crop improvement and breeding programs; gene bank maintenance; incompatibility and fertility studies; evaluation of pollen germination probability after exposure to certain conditions, and in evaluation of dispersal and gene flow Using different terms for different testing criteria based on the developmental stage of the pollen grain being tested (Dafni and Firmage, 2000). Cypermethrin is a synthetic pyreniod used as an insecticide in large scale commercial agriculture application. Viability is defined as the ability to live, develop, or in the case of plants, to germinate when condition are favorable to the plant exist. Studies of pollen viability and morphology are of high importance in relation to genetic breeding programmers.

II. MATERIALS AND METHODS

Location Map - (Plate -1)

Study area - (Plate-2)



Tamilnadu is one of the 28 states of India. Coimbatore is a city in the South Indian State of Tamilnadu. It is lies in South India at 411 metres (1349ft) above sea level. Nirmala College is located in southern parts of Western Ghats. Temperature varies between 28°C to 34°C.

A. Selected Sample

For the present study three types of pollen is selected in the Nirmala College campus. Chilli plants are grown in college garden. The pollen grains of *Capsicum annuum*, L.; is collected from the month of November. Morphology of the pollen grains were observed. And the percentage of pollen viability observed and the data were then represented in tables (Table -1 & 2).

Sample: 1- *Capsicum annuum*, L.;

Systematic Position

Kingdom : Plantae

Class : Dicotyledons

Order : Tubiflorae

Sub order : Solanales

Family : Solanaceae

Genus : *Capsicum*

Species : *C. annuum*, L.;

B. Plant Description

- 1) *Capsicum annuum*, L.; it is native to Southern North America and Northern Southern America. The plant is not an annual but is frost tender. The single flowers are off-white in colour, while the stem is densely branched and up to 60 cm tall.
- 2) *Use*: Fruit are berries that may be green, yellow, orange or red when ripe. Due to its universal use, it is known as “queen of spices”. Chilli peppers are primarily used for culinary purposes, as a spice added to various dishes added and sauces.

C. Methodology: Acetolysis (Erdtman's, 1960)

Acetolysis is the popular method to stain, identify and photograph the pollen It is transferred the pollen grain to 70% alcohol into a centrifuge tube.

Centrifuged the contents for five minutes at 1000rpm and decanted the alcohol. The sediments were washed with glacial acetic acid. 5ml of concentrated sulphuric acid is added into the tube and placed in water bath and heated at 70 °C for 5 minutes. The sediments washed with distilled water.

2ml of glycerin added into the suspension. A drop of suspension placed on a clean slide and cover slip was placed over it. Microphotographs were taken.

D. Application of Insecticide

Healthy *Capsicum annuum* seedlings were grown in an area. The insecticide was applied on the seedling in the field conditions. In total three applications were made.

2ml/L water as the recommended dosage and 4ml/L as double the recommended dosage. A total thee groups one is untreated and two treated groups for insecticide applications were used during this study. This application given by sprayer between 8.00 am and 9.00 am in the morning.

E. Pollen Viability Study

Mature flower buds from three plants were collected in the morning before anthesis. Evaluation of pollen viability followed the method reported by Alexander (1969). Pollen viability was determined by acetocarmine stainability. Flower buds were collected daily at anthesis at about 8.30 am and squashed in drops of acetocarmine stain on a clean glass slide and covered with clean cover slide.


The fully or deeply stained pollen grains with spherical shape were considered viable. Noted the total number of pollen and red stained pollen and calculated the percentage of viable pollen using the formula (Dempsey, 1962).

$$\text{Percentage of viable pollen} = \frac{\text{No of pollen having proper red stain} \times 100}{\text{Total number of pollen}}$$


III. RESULTS AND DISCUSSION

A. Plates Showing Pollen Viability In Recommended Dosages

Pollen viability control Pollen viability of dose -1 Pollen viability of dose -2






Pollen viability of Dose-3



Pollen viability of Dose-4



Pollen viability of Dose-5




B. Showing Viability In Double Recommended Dosages


Pollen viability of Dose-3




Pollen viability of Dose-4




Pollen viability of Dose-5



Pollen viability of Dose-4



Pollen viability of Dose-5



Acetolysed Pollen of *Capsicum*




Table: 1- percentage in recommended dosages: 2ml/100ml water


Control	Dosage	Recommended dosage in percentage
86.7%	Dose 1	86.6%
	Dose 2	64%
	Dose 3	55.5%
	Dose 4	45%
	Dose 5	23.8%

Table: 2 – percentage in double the recommended dosages: 4ml/100ml

Control	Dosage	Double recommended dosage in percentage
86.7%	Dose 1	73.3%
	Dose 2	45%
	Dose 3	34.3%
	Dose 4	26.6%
	Dose 5	13.8%

The study shows the insecticide application on pollen viability of *Capsicum annuum* were varied as shown in (Table1). The percentage of pollen viability in control was 86.7 %. In the first dosage of recommended and double the recommended dosage, the percentage of pollen viability in the recommended dosage and double the recommended after the first treatment were 86.6% (Table- 1) and 73.3% (Table- 2) respectively. The percentage of pollen viability after the second treatment in recommended dose was 64%, The pollen viability obtained for recommended dosages in third, fourth and fifth treatment were 55.5%, 45% and 23.8% (Table- 1) respectively. The rate of decrease in pollen viability shown in double the recommended dosages than the recommended. Whereas double the recommended dose it decreased to 45%. In them the percentage of pollen viability obtained in double recommended dosages in third, fourth and fifth was 34.3%, 26.6% and 13.8% respectively, (Table -2).

Chart: 1 - Percentage of recommended and double recommended dosages



The pollen viability of pepper is high. Carlos *et al.*, (2001) reported high pollen viability in some cultivars of *Capsicum annuum*. The finding indicate than the non-formation of fruit or the formation of seedless fruit observed in some inter specific crosses established and described in literature cannot be attributed to pollen viability alone. The chart- 1 showing viability is decreasing from dose 1 to dose 5.

Blue line that shows the decreasing viability in recommended dosages and the red line show the viability decrease in double recommended dosages. Here the dosages represented in percentage. The graph shows the viability in recommended dosages and double the recommended dosages. Viability is decreased in the recommended dosages from dose 1-dose 5. The dosage level represented in blue line. When it compared to that double recommended dosages viability greatly decreased from dose 1- dose 5. It is represented in red line, (Chart- 1).

Pollen of *capsicum* is white coloured and medium sized. When it is in dry, shape is to become prolate. Pollen triangular, elliptic, suborbicula from equatorial view and three lobed polar views. The aperture were three colpate.

IV. CONCLUSION

The distinction of viable (fertile) from non viable (in fertile) pollen grains was performed with high accuracy. In *Capsicum*, though Cypermethrin is used as an insecticide but now here it seems to act as a growth promoter in the first treatment but have negative effect on further treatment. Pollen viability was found to be decreased in further dosages in both recommended and double recommended dosages. The decrease in pollen viability in double the recommended dose was drastic as compared to the recommended dosages. These treatments resulted in a high degree of plant damage, ovular sterility and reduction in yield. Pollen viability is recorded for several doses. Pollen Morphology of these species varies. Shape of the grain found to be useful in spore or pollen identification. The study of pollen and spores is the subject matter of Palynology. It has immense application in basic sciences, such as taxonomy, and is applied field like geopalynology, aeropalynology, latropalynology, criminology and mellitology.

REFERENCES

- [1] Abo-Hassan, A.A., Nasr T. A and Elshuks. H.A (1982). Effect of type and storage of Pollen on fruiting of Khudrawi dates. The 1st Symp. Date Palm in Saudi Arabia, Pakistan journal of Botany, ISSN; 0556-3321, Vol- 47(1), pp - 102-106.
- [2] Acharya. p, and Rajput (2003). Variability and correlation studies for different traits in *Capsicum* with respect of leaf curl complex . Indian journal of Horticulture, Vol-60(4), ISSN; 2376- 0354, pp - 381 -387.
- [3] Aizen. M, Rovere. A (1995). Does pollen viability decrease with aging. A cross Population examination in *Austrocedrus chilensis* (Cupresaceae). International journal of plant science, Vol-156, ISSN; 1058-5893, pp-227-231.
- [4] Aksoy ozlem, Deveci Asuman, Gonaca. A.L (2013). The affects of some pesticides on pollen germination of *Bignonis semperflorens*. Journal of functional and Environmental Botany, Vol- 3(2), ISSN; 2231-1742, pp-114-119.
- [5] Antonium G.F (2004). Residues and half -lives of pyrethrins of field grown pepper and tomato. Part B. Pesticides, food contaminants and agricultural waste, Journal of Environmental science and health, Vol- 39(4), ISSN-1093-4529; pp- 491-503.
- [6] Arya, P.S. Saini S.S (1976). Variability and correlation studies in chillies (*Capsicum frutescens* L.) Haryana journal of Horticultural science, Vol -10, ISSN - 0970-2873; pp- 236-244.
- [7] Cruz. D, Acosta. J and Negron. F (1984). Evaluation of various pesticides for the control of aphids, mites and nematodes in peppers. The journal of Agriculture of the University of Puerto Rico, Vol- 6, ISSN-0041-994X, pp- 457-465.
- [8] Einhardt et al., (2006) and Nycine and Pilley (2007). Pollen viability and germination in *Jatropha ribifolia* and *Jatropha mollissima*. African journal of Biotechnology. Vol 10 (3). ISSN- 1684-5315; pp- 368-374.
- [9] Tort I.N, Zturk Guvensen (2005). Effects of same fungicides on pollen morphology and anatomy of Tomato (*Lycopersicon esculentum* mill) Pakistan Journal of Botany, Vol-37 (1), ISSN; 0556-3321, pp-23-30.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)