



IJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 7 Issue: III Month of publication: March 2019

DOI: <http://doi.org/10.22214/ijraset.2019.3442>

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

Fibonacci Product Cordial Labelling of some Graphs

Tesssymol Abraham¹, Shiny Jose²

¹Department of Mathematics, Baselius College, Kottayam-686001

²St. George's College, Aruvithura, Kottayam-686112

^{1,2}Mahatma Gandhi University, Kottayam, Kerala

Abstract— An injective function $\varphi: V(G) \rightarrow \{F_1, F_2, \dots, F_n\}$, where F_j is the j^{th} Fibonacci number ($j = 1, \dots, n$), is said to be Fibonacci product cordial labeling if the induced function $\varphi^*: E(G) \rightarrow \{0, 1\}$ defined by $\varphi^*(uv) = (\varphi(u)\varphi(v)) \bmod 2$ satisfies the condition $|e_{\varphi^*}(0) - e_{\varphi^*}(1)| \leq 1$. A graph which admits Fibonacci product cordial labeling is called Fibonacci product cordial graph. In this paper we investigated the Fibonacci product cordial labeling of Shell graph, Comb, Crown, Star graph and Complete bipartite graphs $K_{2,n}$.

Keywords— Fibonacci Product Cordial Labeling; Fibonacci Product Cordial graph, Shell, Comb, Crown

I. INTRODUCTION

In graph labeling vertices or edges or both assigned values subject to certain conditions. There are many types of labeling. More number of labeling techniques found their starting point from graceful labeling introduced by Rosa[7]. A Fibonacci cordial labeling have been introduced by Rokad, A.K., and G.V.Ghodasara[6]. Also Fibonacci product cordial labeling have introduced by Tesssymol Abraham and Shiny Jose [8].

We begin with simple, finite, connected and undirected graph $G = (V(G), E(G))$ with p vertices and q edges. In this paper we discussed Fibonacci product cordial labeling of Shell graph, Comb, Crown, Star graph and Complete bipartite graphs $K_{2,n}$. We will provide brief summary of definitions and other information which are necessary for present investigations

Notations:

$v_\varphi(0)$: Number of vertices with label 0 under the map φ

$v_\varphi(1)$: Number of vertices with label 1 under the map φ

$e_{\varphi^*}(0)$: Number of edges with label 0 under the map φ^*

$e_{\varphi^*}(1)$: Number of edges with label 1 under the map φ^*

FPCL: Fibonacci product cordial labeling

II. PRELIMINARIES

A. Definition 2.1 Fibonacci Product Cordial Labeling [8]

An injective function $\varphi: V(G) \rightarrow \{F_1, F_2, F_3, \dots, F_n\}$ where F_j is the j^{th} Fibonacci number ($j = 1, \dots, n$), is said to be Fibonacci product cordial labeling if the induced function $\varphi^*: E(G) \rightarrow \{0, 1\}$ defined by $\varphi^*(uv) = (\varphi(u)\varphi(v)) \bmod 2$ satisfies the condition $|e_{\varphi^*}(0) - e_{\varphi^*}(1)| \leq 1$.

B. Definition 2.2 Fibonacci Product Cordial Graph [8]

A graph which admits Fibonacci product cordial labeling is called Fibonacci product cordial graph

C. Definition 2.3 Comb Graph [3]

The comb is a graph obtained by joining a single pendent edge to each vertex of a path. It is denoted by $P_n AK_1$

D. Definition 2.4 Shell Graph [5]

A shell graph of size n , denoted by S_n is the graph obtained from the cycle $C_n(v_1 v_2 \dots v_n)$ by adding $(n - 3)$ consecutive chords incident with a common vertex. The vertex at which all the chords are concurrent is called the apex vertex. The shell is also called fan $fn-1$. Thus, $S_n = fn-1 = P_{n-1} + K_1$.

E. Definition 2.5 Crown Graph [3]

A crown graph is the graph obtained from the cycle $C_n(v_1 v_2 \dots v_n)$ with pendant edge attached at each vertex of the cycle. It is denoted by $C_n \odot K_1$

III. PROPERTIES OF FIBONACCI PRODUCT CORDIAL LABELING

A. Theorem 3.1

Shell graph S_n is a Fibonacci product cordial graph for $n \geq 3, n \in N$.

1) *Proof:* Let w_1 be the common end vertex (apex) and w_2, w_3, \dots, w_n be the other vertices of a Shell Graph S_n .

Then each edge is of the form $w_1 w_j$ where $2 \leq j \leq n$.

We define labeling function $\varphi: V(G) \rightarrow \{F_1, F_2, F_3, \dots, F_n\}$ as follows.

$$\varphi(w_i) = F_i \text{ if } 1 \leq i \leq n$$

Define $\varphi^*: E(G) \rightarrow \{0, 1\}$ by $\varphi^*(w_1 w_j) = (\varphi(w_1) \varphi(w_j)) \pmod 2$ for $2 \leq j \leq n$

Then we have $|\epsilon_{\varphi^*}(0) - \epsilon_{\varphi^*}(1)| \leq 1$. Hence Shell graph admits Fibonacci product cordial labeling and hence it is a Fibonacci product cordial graph.

2) Illustration 3.1:




Figure 3.1. FPCL Of Shell Graph S_{11}

B. Theorem 3.2

The Comb is Fibonacci product cordial graph for all $n \in N$

1) *Proof:* Let G be a comb graph $P_n AK_1$ with $2n$ vertices and $2n-1$ edges. Let

$V(G) = \{v_1, v_2, \dots, v_n, w_1, w_2, \dots, w_n\}$ where v_1, v_2, \dots, v_n are the vertices of the path P_n and

w_1, w_2, \dots, w_n are the pendant vertices attaching by the path P_n . Then $E(G) = \{e_i, e_{ij}\}$ where $e_i = (v_i, w_i)$ and

$$e_{ij} = (v_i, v_j)$$

Case 1: $n \equiv 0, 1 \pmod 3$

Define the function $\varphi: V(G) \rightarrow \{F_1, F_2, F_3, \dots, F_{2n}\}$ as

$$\varphi(v_i) = \begin{cases} F_{2i-2} & \text{if } i \equiv 0 \pmod 3 \\ F_{2i} & \text{if } i \equiv 1 \pmod 3 \\ F_{2i+2} & \text{if } i \equiv 2 \pmod 3 \end{cases}$$

$$\varphi(w_i) = F_{2i-1} \text{ if } 1 \leq i \leq n$$

Then $\varphi^*: E(G) \rightarrow \{0, 1\}$ by $\varphi^*(e_i) = (\varphi(v_i)\varphi(w_i)) \bmod 2$ and $\varphi^*(e_{ij}) = (\varphi(v_i)\varphi(v_j)) \bmod 2$ admits Fibonacci product cordial labeling. Hence Combs are Fibonacci product cordial graphs.

2) *Illustration 3.2:*

..




Figure 3.2. FPCL Of Comb P_7AK_1

Case 2: $n \equiv 2 \pmod 3$

Define the function $\varphi: V(G) \rightarrow \{F_1, F_2, F_3, \dots, F_{2n}\}$ as

For $1 \leq i \leq n - 1$

$$\varphi(v_i) = \begin{cases} F_{2i-2} & \text{if } i \equiv 0 \pmod 3 \\ F_{2i} & \text{if } i \equiv 1 \pmod 3 \\ F_{2i+2} & \text{if } i \equiv 2 \pmod 3 \end{cases}$$

$$\varphi(v_i) = F_{2i} \quad \text{if } i = n$$

$$\varphi(w_i) = F_{2i-1} \quad \text{if } 1 \leq i \leq n$$

Then $\varphi^*: E(G) \rightarrow \{0, 1\}$ by $\varphi^*(e_i) = (\varphi(v_i)\varphi(w_i)) \bmod 2$ and $\varphi^*(e_{ij}) = (\varphi(v_i)\varphi(v_j)) \bmod 2$ admits Fibonacci product cordial labeling. Hence Combs are Fibonacci product cordial graphs.

3) *Illustration 3.3:*




Figure 3.3. FPCL Of Path $P_{11}AK_1$

C. Theorem 3.3

The Crown $C_n \odot K_1$ is a Fibonacci product cordial graph for all $n \geq 3, n \in \mathbb{N}$

1) *Proof:* Let $C_n \odot K_1$ be a Crown graph with $2n$ vertices and $2n$ edges. Let $V(G) = \{v_1, v_2, \dots, v_n, w_1, w_2, \dots, w_n\}$ where v_1, v_2, \dots, v_n are the vertices of the cycle C_n and w_1, w_2, \dots, w_n are the pendant vertices attaching by the cycle C_n . Then $E(G) = \{e_i, e_{ij}\}$ where $e_i = (v_i, w_i)$ and $e_{ij} = (v_i, v_j)$

Case 1: $n \equiv 0 \pmod 3$

Define the function $\varphi: V(C_n \odot K_1) \rightarrow \{F_1, F_2, F_3, \dots, F_{2n}\}$ as

$$\varphi(v_i) = \begin{cases} F_{2i-2} & \text{if } i \equiv 0 \pmod 3 \\ F_{2i} & \text{if } i \equiv 1 \pmod 3 \\ F_{2i+2} & \text{if } i \equiv 2 \pmod 3 \end{cases}$$

$$\varphi(w_i) = F_{2i-1} \quad \text{if } 1 \leq i \leq n$$

Then $\varphi^*: E(C_n \odot K_1) \rightarrow \{0, 1\}$ by $\varphi^*(e_i) = (\varphi(v_i)\varphi(w_i)) \bmod 2$ and $\varphi^*(e_{ij}) = (\varphi(v_i)\varphi(v_j)) \bmod 2$ admits Fibonacci product cordial labeling. Hence Crowns are Fibonacci product cordial graphs.

2) Illustration 3.4 :




Figure 3.4. FPCL Of Crown $C_9 \odot K_1$

Case 2: $n \equiv 1, 2 \pmod 3$

Define the function $\varphi: V(C_n \odot K_1) \rightarrow \{F_1, F_2, F_3, \dots, F_{2n}\}$ as

For $1 \leq i \leq n - 5$

$$\varphi(v_i) = \begin{cases} F_{2i-2} & \text{if } i \equiv 0 \pmod 3 \\ F_{2i} & \text{if } i \equiv 1 \pmod 3 \\ F_{2i+2} & \text{if } i \equiv 2 \pmod 3 \end{cases}$$

For $n - 4 \leq i \leq n$

$$\varphi(v_i) = F_{2i}$$

$$\varphi(w_i) = F_{2i-1} \quad \text{if } 1 \leq i \leq n$$

Then $\varphi^*: E(C_n \odot K_1) \rightarrow \{0, 1\}$ by $\varphi^*(e_i) = (\varphi(v_i)\varphi(w_i)) \bmod 2$ and $\varphi^*(e_{ij}) = (\varphi(v_i)\varphi(v_j)) \bmod 2$ admits Fibonacci product cordial labeling. Hence Crowns are Fibonacci product cordial graphs.

3) Illustration 3.5:




Figure 3.5. FPCL Of Crown $C_8 \odot K_1$

D. Theorem 3.4:

The Star graphs $K_{1,n}$ are Fibonacci product cordial graphs for $n = 2,3,5$

1) *Proof:*

Let u be the apex vertex and $\{v_1, v_2, \dots, v_n\}$ be the pendant vertices of the star graph $K_{1,n}$. Then $E(K_{1,n}) = \{e_i\}$

where $e_i = (u, v_i)$

Define: $V(K_{1,n}) \rightarrow \{F_1, F_2, F_3, \dots, F_n\}$ as

$$\varphi(u) = F_1 \quad \text{and} \quad \varphi(v_i) = F_{i+1} \quad \text{for } 1 \leq i \leq n$$

Also define $\varphi^*: E(K_{1,n}) \rightarrow \{0, 1\}$ by $\varphi^*(e_i) = (\varphi(u)\varphi(v_i)) \bmod 2$. Then we have $|e_{\varphi^*(0)} - e_{\varphi^*(1)}| \leq 1$.

Hence Star graphs $K_{1,n}$, $1 \leq n \leq 3$ are Fibonacci product cordial graphs.

2) *Illustration 3.6:*




Figure 3.6.FPCL Of Star Graph $K_{1,3}$

E. Theorem 3.5

The Complete bipartite graphs $K_{2,n}$, $1 \leq n \leq 4$ are Fibonacci product cordial graphs

1) *Proof:* Let $X = \{u_1, u_2\}$ and $Y = \{v_1, v_2, \dots, v_n\}$ be the bipartition of the vertices of the complete bipartite graphs $K_{2,n}$

Then $E(G) = \{e_{ij}\}$ where $e_{ij} = (u_i, v_j)$ for $i = 1, 2$, and $1 \leq j \leq n$

Case 1: $n \equiv 0 \pmod 2$

Define: $V(K_{2,n}) \rightarrow \{F_1, F_2, F_3, \dots, F_n\}$ as

$$\varphi(u_i) = F_i \quad \text{for } 1 \leq i \leq 2$$

$$\varphi(v_i) = F_{2+i} \quad \text{for } 1 \leq i \leq n$$

Also define $\varphi^*: E(K_{2,n}) \rightarrow \{0, 1\}$ by $\varphi^*(e_i) = (\varphi(u)\varphi(v_i)) \bmod 2$. Then we have $|e_{\varphi^*(0)} - e_{\varphi^*(1)}| \leq 1$.

Hence Star graphs $K_{1,n}$, $1 \leq n \leq 3$ are Fibonacci product cordial graphs.

Case 1: $n \equiv 1 \pmod 2$

Define: $V(K_{2,n}) \rightarrow \{F_1, F_2, F_3, \dots, F_n\}$ as

$$\varphi(u_i) = F_{2i-1} \quad \text{for } 1 \leq i \leq 2$$

$$\varphi(v_i) = F_{2i} \quad \text{for } 1 \leq i \leq n$$

Then $\varphi^*: E(K_{2,n}) \rightarrow \{0, 1\}$ by $\varphi^*(e_i) = (\varphi(v_i)\varphi(w_i)) \bmod 2$ and $\varphi^*(e_{ij}) = (\varphi(v_i)\varphi(v_j)) \bmod 2$ admits

Fibonacci product cordial labeling. Hence Complete bipartite graphs $K_{2,n}$ are Fibonacci product cordial graphs.

2) *Illustration 3.7:*




Figure 3.7. FPCL Of Complete Graph $K_{2,4}$



IV. CONCLUSION

Here we proved five new theorems corresponding to Fibonacci product cordial labeling. Analogous results can be derived for other graph families as well as for different graph labeling problems..

REFERENCES

- [1] Cahit ,I.,1987. Cordial graphs: a weaker version of graceful and harmonious graphs,Ars Com-bin.,23: 201-207.
- [2] David M. Burton . Elementary Number Theory, Sixth edition, Tata McGraw-Hill Publishing Company Limited, New Delhi.
- [3] Gallian J.A. 2014. A dynamical survey of graph labeling, The Electronic Journal of Combinatorics, 17, DS6.
- [4] Harary, F.1972. Graph Theory, Addison Wesley, Reading Mass
- [5] S. .Meera ,M. Renugha and M.Sivasakthi. Sep 2015. Cordial Labeling For Different Types Of shell Graph, International Journal Of Scientific &Engineering, Vol 6,pp 1282-1288 .
- [6] Rokad A.H. and G.V.Ghudasara.Feb 2016 .Fibonacci Cordial Labeling of Some Special Graphs , Annals of Pure and Applied Mathematics, Vol. 11, PP 133-144
- [7] Rosa, A.1966-1967.On certain valuation of the vertices of a graph, Theory of Graphs, International Symposium, Rome, New York, and Dunod Paris : 349-355.
- [8] Tessymol abraham and Dr.Shiny Jose .Jan 2019.Fibonacci product cordial labeling, International Journal of Emerging Technologies and Innovative Research,Vol.6,PP 58-63.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)