



# **iJRASET**

International Journal For Research in  
Applied Science and Engineering Technology



---

# **INTERNATIONAL JOURNAL FOR RESEARCH**

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

---

**Volume: 3**

**Issue: V**

**Month of publication: May 2015**

**DOI:**

**[www.ijraset.com](http://www.ijraset.com)**

**Call:  08813907089**

**E-mail ID: [ijraset@gmail.com](mailto:ijraset@gmail.com)**

# On $Q^*$ - Homeomorphisms in Topological Spaces

P.Padma<sup>1</sup>, V.Ramya<sup>2</sup>, S.Udayakumar<sup>3</sup>

<sup>1</sup>Department of Mathematics, PRIST University, Thanjavur, INDIA

<sup>2</sup>Department of Mathematics, Sri Akilandeswari Women's college, Vandhavasi, Thiruvannamalai, INDIA.

<sup>3</sup>Department of Mathematics, A.V.V.M Sri Pushpam college, Poondi, Thanjavur, INDIA

**Abstract:** In this paper, we first introduce a new class of closed map called  $Q^*$  - closed map. Moreover, we introduce a new class of homeomorphism called  $Q^*$  - homeomorphism, which are weaker than homeomorphism. We also introduce  $Q^*$  - homeomorphisms and prove that the set of all  $Q^*$ -homeomorphisms forms a group under the operation of composition of maps. We prove that  $Q^*$  - homeomorphism and  $Q^*$  - homeomorphism are independent.

**Subject classification:** 54G12

**Keywords:**  $Q^*$  homeomorphisms,  $Q^*$  - homeomorphisms,  $Q^*$  compact space, pre  $Q^*$  - open,  $Q^*$ -closed map,  $Q^*$  - open map.

## I. INTRODUCTION

The notion homeomorphism plays a very important role in topology. By definition, a homeomorphism between two topological spaces  $X$  and  $Y$  is a bijective map  $f : X \rightarrow Y$  when both  $f$  and  $f^{-1}$  are continuous. It is well known that as Janich [[8], p.13] says correctly: homeomorphisms play the same role in topology that linear isomorphism's play in linear algebra, or that biholomorphic maps play in function theory, or group isomorphism's in group theory, or isometrics in Riemannian geometry. In 1995, Maki, Devi and Balachandran [4] introduced the concepts of semi - generalized homeomorphisms and generalized semi-homeomorphisms and studied some semi topological properties. Devi and Balachandran [3] introduced a generalization of  $\alpha$ -homeomorphism in 2001. Recently, Devi, Vigneshwaran, Vadivel and Vairamanickam [5,23] introduced  $g^*\alpha$ -homeomorphisms and  $rg\alpha$ - homeomorphisms and obtained some topological properties. In this paper, I first introduce a new class of closed map called  $Q^*$  - closed map. Moreover, we introduce a new class of homeomorphism called  $Q^*$  - homeomorphism, which are weaker than homeomorphism. We also introduce  $Q^*$  - homeomorphisms and prove that the set of all  $Q^*$ -homeomorphisms forms a group under the operation of composition of maps. We prove that  $Q^*$  - homeomorphism and  $Q^*$  - homeomorphism are independent.

## II. PRELIMINARIES

Throughout this paper  $(X, \tau)$  and  $(Y, \sigma)$  represent topological spaces on which no separation axioms are assumed unless otherwise mentioned. For a subset  $A$  of a space  $(X, \tau)$ ,  $cl(A)$ ,  $int(A)$  and  $A^c$  denote the closure of  $A$ , the interior of  $A$  and the complement of  $A$  in  $X$ , respectively. We recall the following definitions and some results, which are used in the sequel.

Definition 2.1 A set  $A$  is said to be semi open if there exists an open set  $G$  such that  $G \subset A \subset cl(G)$  or  $A \subset cl(int(A))$ . The set  $A$  is semi closed if  $int(cl(A)) \subset A$ .

Definition 2.2 The function  $f : X \rightarrow Y$  is said to be irresolute if  $f^{-1}(A)$  is a semi open set whenever  $A$  is a semi open set in  $Y$ .

Definition 2.3 A subset  $A$  of a space  $(X, \tau)$  is called pre-closed if  $cl(int(A)) \subset A$ .

Definition 2.4 A space  $X$  is said to be  $Q^*$ -regular if for every  $Q^*$  - closed set  $F$  and a point  $x \notin F$ , there exist disjoint  $Q^*$  - open sets  $U$  and  $V$  such that  $x \in U$  and  $F \subset V$ .

Definition 2.5[15] A subset  $A$  of a topological space  $(X, \tau)$  is called a  $Q^*$  - closed if  $int(A) = \emptyset$  and  $A$  is closed

## III. $Q^*$ HOMEOMORPHISMS

$Q^*$  - closed sets in topological spaces were introduced by M. Murugalingam and N. Lalitha in 2010. In this section we study the properties of  $Q^*$  - homeomorphisms,  $Q^*$  compact spaces are discussed.

Definition 3.1 A function  $f : (X, \tau) \rightarrow (Y, \sigma)$  is pre  $Q^*$  - closed if  $f(U)$  is  $Q^*$  closed in  $Y$  for every  $Q^*$  closed set in  $X$ .

Definition 3.2 A bijection  $f : (X, \tau) \rightarrow (Y, \sigma)$  is a  $Q^*$  - homeomorphism if  $f$  is  $Q^*$  irresolute and pre  $Q^*$  - open.

Example 3.1 Let  $X = Y = \{a, b, c\}$ ,  $\tau = \{\emptyset, X, \{b\}, \{b, c\}\}$  and  $\sigma = \{\emptyset, Y, \{b, c\}\}$ . Let  $f : X \rightarrow Y$  be the identity map. Hence  $f$  is  $Q^*$  - homeomorphism.

Proposition 3.1 Every  $Q^*$  - closed set is  $\alpha$ -closed and hence pre-closed.

## International Journal for Research in Applied Science & Engineering Technology (IJRASET)

Definition 3.3 A space  $X$  is said to be  $Q^*$ -compact if every cover of  $X$  by  $Q^*$ -open sets has a finite sub cover .

Definition 3.4 A map  $f : X \rightarrow Y$  is called  $Q^*$ -closed map if the image of each  $Q^*$ -closed in  $(X, \tau)$  is closed in  $(Y, \sigma)$  .

Example 3.2 (1) Let  $X = Y = \{ a, b, c \}$ ,  $\tau = \{ \phi, X, \{ b \} \}$  and  $\sigma = \{ \phi, Y, \{ b \}, \{ b, c \} \}$ . Then  $\phi, \{ a, c \}$  are  $Q^*$ -closed in  $X$  . Let  $f : X \rightarrow Y$  be the identity map . Then  $f(\phi) = \phi, f(\{ a, c \}) = \{ a, c \}$  . Since  $\phi, \{ a, c \}$  are closed in  $Y$  . Therefore ,  $f$  is -  $Q^*$  closed map .

Let  $X = Y = \{ a, b, c \}$ ,  $\tau = \{ \phi, X, \{ a, b \} \}$  and  $\sigma = \{ \phi, Y, \{ a \}, \{ b \} \}$  . Let  $f : (X, \tau) \rightarrow (Y, \sigma)$  be the identity map . Then  $f$  is not a  $Q^*$ - closed map .

The next theorem shows that normality is preserved under continuous

Definition 3.5 Let  $x$  be a point of  $(X, \tau)$  and  $V$  be a subset of  $X$  . Then  $V$  is called a  $Q^*$ -neighborhood of  $x$  in  $(X, \tau)$  if there exists a  $Q^*$ - open set  $U$  of  $(X, \tau)$  such that  $x \in U \subset V$  .

Theorem3.1 If  $f : (X, \tau) \rightarrow (Y, \sigma)$  is a  $Q^*$ - continuous ,  $Q^*$ - closed map from a  $Q^*$ -normal space  $(X, \tau)$  onto a space  $(Y, \sigma)$  then  $(Y, \sigma)$  is  $Q^*$ - normal.

Proof. Since every  $Q^*$ -closed map is  $Q^*$ -closed.

Analogous to a  $Q^*$ -closed map, we define a  $Q^*$ -open map as follows:

Definition 3.6 A map  $f : (X, \tau) \rightarrow (Y, \sigma)$  is said to a  $Q^*$ -open map if theimage  $f(A)$  is  $Q^*$ -open in  $(X, \tau)$  for each open set  $A$  in  $(Y, \sigma)$  .

Proposition 3.2 For any bijection  $f : (X, \tau) \rightarrow (Y, \sigma)$ , the followingstatements are equivalent:

- (i)  $f^{-1} : (Y, \sigma) \rightarrow (X, \tau)$  is  $Q^*$ -continuous,
- (ii)  $f$  is a  $Q^*$ -open map and
- (iii)  $f$  is a  $Q^*$ -closed map.

Proof.

Step 1 : (a)  $\Rightarrow$  (b)

Let  $V$  be a  $Q^*$ - open set in  $X$  . Then  $X - V$  is  $Q^*$ - closed in  $X$  . Since  $f^{-1}$  is  $Q^*$ - continuous ,  $(f^{-1})^{-1}(X - V) = f(X - V) = Y - f(V)$  is closed in  $Y$  . Then  $f(V)$  is open in  $Y$  . Hence  $f$  is a  $Q^*$ - open map.

Step 2 : (b)  $\Rightarrow$  (c) .

Let  $f$  be a  $Q^*$ - open map . Let  $U$  be a  $Q^*$ - closed set in  $X$  . Then  $X - U$  is  $Q^*$ - open in  $X$  . Since  $f$  is  $Q^*$ - open ,  $f(X - U) = Y - f(U)$  is open in  $Y$  . Then  $f(U)$  is closed in  $Y$  . Hence  $f$  is a  $Q^*$ - closed .

Step 3 : (c)  $\Rightarrow$  (a) .

Let  $V$  be  $Q^*$ - closed set in  $X$  . Since  $f : X \rightarrow Y$  is  $Q^*$ - closed ,  $f(V)$  is closed in  $Y$  . That is  $(f^{-1})^{-1}(V)$  is closed in  $Y$  . Hence  $f^{-1}$  is  $Q^*$ - continuous. ■

Proposition 3.3 Let  $f : X \rightarrow Y$  be a bijective and  $Q^*$ - continuous map. Then the following statements are equivalent.

- (a)  $f$  is a  $Q^*$ - open map.
- (b)  $f$  is a  $Q^*$ - homeomorphism.
- (c)  $f$  is a  $Q^*$ - closed map.

Proof .

Step 1 : (a)  $\Rightarrow$  (b) .

Given  $f$  is bijective ,  $Q^*$ - continuous map and  $Q^*$ - open map . Hence  $f$  is  $Q^*$ - homeomorphism.

Step 2 : (b)  $\Rightarrow$  (c) .

Suppose (b) holds. Let  $F$  be a closed set in  $(X, \tau)$  . By (b) ,  $f^{-1}$  is  $Q^*$ - continuous and so  $(f^{-1})^{-1}(F) = f(F)$  is  $Q^*$ - closed in  $(Y, \sigma)$  . This proves (c).

Step 3 : (c)  $\Rightarrow$  (a) .

Follows from Proposition 3.2 .

Definition 3.7 Let  $A$  be a subset of  $X$ . A mapping  $r : X \rightarrow A$  is called a  $Q^*$ - continuous retraction if  $r$  is  $Q^*$ - continuous and the restriction  $\tau_A$  is the identity mapping on  $A$  .

Definition 3.8 A topological space  $(X, \tau)$  is called a  $Q^*$ - Hausdorff if for each pair  $x, y$  of distinct points of  $X$  , there exists  $Q^*$ - neighborhoods  $U_1$  and  $U_2$  of  $x$  and  $y$ , respectively, that are disjoint .

Lemma 3.1 Let  $A$  be a subset of  $X$ . Then  $p \in Q^* - cl(A)$  if and only if for any  $Q^*$ - neighborhood  $N$  of  $p$  in  $X$ ,  $A \cap N \neq \phi$

Theorem 3.1 Let  $A$  be a subset of  $X$  and  $r : X \rightarrow A$  be a  $Q^*$ - continuous retraction . If  $X$  is  $Q^*$ -Hausdorff , then  $A$  is a  $Q^*$ - closed set of  $X$ .

## International Journal for Research in Applied Science & Engineering Technology (IJRASET)

Proof. Suppose that  $A$  is not  $Q^*$ -closed. Then there exists a point  $x$  in  $X$  such that  $x \in Q^*-\text{cl}(A)$  but  $x \notin A$ . It follows that  $r(x) \neq x$  because  $r$  is  $Q^*$ -continuous retraction. Since  $X$  is  $Q^*$ -Hausdorff, there exists disjoint  $Q^*$ -open sets  $U$  and  $V$  in  $X$  such that  $x \in U$  and  $r(x) \in V$ . Now let  $W$  be an arbitrary  $Q^*$ -neighborhood of  $x$ . Then  $W \cap U$  is a  $Q^*$ -neighborhood of  $x$ .

Since  $x \in Q^*-\text{cl}(A)$ , by Lemma 3.1, we have  $(W \cap U) \cap A \neq \emptyset$ . Therefore there exists a point  $y$  in  $W \cap U \cap A$ . Since  $y \in A$ , we have  $r(y) = y \in U$  and hence  $r(y) \notin V$ . This implies that  $r(W) \not\subset V$  because  $y \in W$ . This is contrary to the  $Q^*$ -continuity of  $r$ . Consequently,  $A$  is a  $Q^*$ -closed set of  $X$ . ■

Theorem 3.1 Let  $f : X \rightarrow Y$  be a bijection. If  $f$  is pre  $Q^*$ -open then  $f^{-1} : X \rightarrow Y$  is irresolute.

Proof. Suppose that the bijection  $f$  is pre  $Q^*$ -open then  $f^{-1}$  is a function from  $Y$  to  $X$ . Let  $U$  be a  $Q^*$ -open set in  $X$ . Since every  $Q^*$ -open set is semi open we have  $U$  is semi open in  $X$ . Then  $f(U)$  is a semi open set in  $Y$ . But  $f(U) = (f^{-1})^{-1}(U)$ . Hence  $(f^{-1})^{-1}(U)$  is a semi open set in  $Y$ . Put  $g = f^{-1}$ . We have  $g^{-1}(U)$  is semi open. Consequently,  $g$  is irresolute. That is,  $f^{-1}$  is irresolute. ■

Theorem 3.2 Let  $f : X \rightarrow Y$  be a bijection. Then  $f$  is pre  $Q^*$ -open if and only if  $f^{-1} : Y \rightarrow X$  is  $Q^*$ -irresolute.

Proof.

Step 1 : Suppose that the bijection  $f$  is pre  $Q^*$ -open then  $f^{-1}$  is a function from  $Y$  to  $X$ . Let  $U$  be a  $Q^*$ -open set in  $X$ . Then  $f(U)$  is a  $Q^*$ -open set in  $Y$ . But  $f(U) = (f^{-1})^{-1}(U)$ . Hence  $(f^{-1})^{-1}(U)$  is a  $Q^*$ -open set in  $Y$ . Put  $g = f^{-1}$ . We have  $g^{-1}(U)$  is  $Q^*$ -open. Consequently,  $g$  is  $Q^*$ -irresolute. That is,  $f^{-1}$  is  $Q^*$ -irresolute.

Step 2 : Suppose that  $f$  is  $Q^*$ -irresolute. Let  $U$  be a  $Q^*$ -open set in  $X$ . Then  $(f^{-1})^{-1}(U)$  is a  $Q^*$ -open set in  $Y$ . But  $(f^{-1})^{-1}(U) = f(U)$ . Therefore  $f(U)$  is a  $Q^*$ -open set in  $Y$ . Hence  $f$  is pre  $Q^*$ -open. ■

Theorem 3.3 If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are both  $Q^*$ -irresolute, then their composition  $g \circ f : X \rightarrow Z$  is an  $Q^*$ -irresolute map.

Proof. Let  $V$  be  $Q^*$ -open set in  $Z$ . Then  $(g \circ f)^{-1}(V) = (f^{-1} \circ g^{-1})(V) = (f^{-1}(g^{-1}(V)))$ . Since  $g$  is  $Q^*$ -irresolute, it follows that  $g^{-1}(V)$  is a  $Q^*$ -open set. Since  $f$  is  $Q^*$ -irresolute, it follows that  $(f^{-1}(g^{-1}(V)))$  is a  $Q^*$ -open set. Thus for each  $Q^*$ -open set  $V$  in  $Z$ ,  $(g \circ f)^{-1}(V)$  is  $Q^*$ -open in  $X$ . Therefore,  $g \circ f$  is an  $Q^*$ -irresolute function. ■

Theorem 3.4  $Q^*$ -homeomorphism is an equivalence relation. We write  $X \sim Y$  whenever two spaces  $X, Y$  are  $Q^*$ -homeomorphic.

Proof.

Step 1 : Let  $i : X \rightarrow X$  be the identity map on  $X$ . Then it is bijective and  $Q^*$ -irresolute. Also  $(i)^{-1}$  is a pre  $Q^*$ -open map. Hence  $i$  is a  $Q^*$ -homeomorphism. Accordingly  $X \sim X$ . The relation is reflexive.

Step 2 : Suppose that  $X \sim Y$ . Then there exists a  $Q^*$ -homeomorphism  $h : X \rightarrow Y$ . But then  $h$  is bijective. Accordingly  $h^{-1} : Y \rightarrow X$  is bijective. Also  $h$  is  $Q^*$ -irresolute. Hence  $h^{-1}$  is a pre  $Q^*$ -open map. Hence  $Y \sim X$ .

Step 3: Suppose that  $X \sim Y$  and  $Y \sim Z$ . Then there is a  $Q^*$ -homeomorphism  $f : X \rightarrow Y$  and there is a  $Q^*$ -homeomorphism  $g$  from  $Y$  to  $Z$ . But then  $f$  and  $g$  are bijective. Accordingly,  $g \circ f$  is bijective and pre  $Q^*$ -open. Thus,  $g \circ f$  is  $Q^*$ -homeomorphism. Therefore,  $X \sim Z$ . Hence  $\sim$  is transitive. From step (1), (2) and (3),  $Q^*$ -homeomorphism is an equivalence relation. ■

Theorem 3.5 Every  $Q^*$ -compact subset of a Hausdorff space is semiclosed.

Proof. Suppose that  $A$  be a  $Q^*$ -compact subset of a Hausdorff space  $X$ . Since every  $Q^*$ -compact space is semi compact we have  $A$  is semi compact. Let  $x \in X - A$ . Then there exists 2 disjoint semi open sets  $U_x$  and  $V_x$  such that  $x \in U_x$  and

## International Journal for Research in Applied Science & Engineering Technology (IJRASET)

$A \subset V_x$ . But then  $X \in U_x \subset X - V_x \subset X - A$ . Therefore,  $X - A$  is semi open. Hence  $A$  is semi-closed in  $Y$ . ■

**Theorem 3.6** Every  $Q^*$ -compact subset of a  $Q^*$ -Hausdorff space is  $Q^*$ -closed.

**Proof.** Suppose that  $A$  be a  $Q^*$ -compact subset of a  $Q^*$ -Hausdorff space  $X$ . Let  $x \in X - A$ . Then there are 2 disjoint  $Q^*$ -open sets  $U_x$  and  $V_x$  such that  $x \in U_x$  and  $A \subset V_x$ . But then  $X \in U_x \subset X - V_x \subset X - A$ . Therefore,  $X - A$  is  $Q^*$ -open. Hence  $A$  is  $Q^*$ -closed in  $Y$ . ■

**Theorem 3.7** Let  $X$  be  $Q^*$ -compact and set  $Y$  be a Hausdorff space. If  $f : X \rightarrow Y$  is  $Q^*$ -continuous,  $Q^*$ -irresolute and bijective, then  $f$  is a  $Q^*$ -homeomorphism.

**Proof.** Let  $A$  be a  $Q^*$ -closed subset of the  $Q^*$ -compact space  $X$ . Then  $A$  is  $Q^*$ -compact. But  $f$  is  $Q^*$ -irresolute. Hence  $f(A)$  is  $Q^*$ -compact. Take  $g = f^{-1}$ . Then  $g^{-1}(A)$  is  $Q^*$ -closed, by theorem 3.4. Consequently  $g$  is a  $Q^*$ -irresolute map. That is,  $f^{-1}$  is  $Q^*$ -irresolute. Therefore,  $f$  is a  $Q^*$ -homeomorphism. ■

**Proposition 3.4** If  $f : (X, \tau) \rightarrow (Y, \sigma)$  and  $g : (Y, \sigma) \rightarrow (Z, \gamma)$  are  $Q^*$ -homeomorphisms, then their composition  $g \circ f : (X, \tau) \rightarrow (Z, \gamma)$  is also  $Q^*$ -homeomorphism.

**Proof.** Let  $U$  be a  $Q^*$ -open set in  $(Z, \gamma)$ . Now,  $(g \circ f)^{-1}(U) = f^{-1}(g^{-1}(U)) = f^{-1}(V)$ , where  $V = g^{-1}(U)$ . By hypothesis,  $V$  is  $Q^*$ -open in  $(Y, \sigma)$  and so again by hypothesis,  $f^{-1}(V)$  is  $Q^*$ -open in  $(X, \tau)$ . Therefore,  $g \circ f$  is  $Q^*$ -irresolute. Also for a  $Q^*$ -open set  $G$  in  $(X, \tau)$ , we have  $(g \circ f)(G) = g(f(G)) = g(W)$ , where  $W = f(G)$ . By hypothesis  $f(G)$  is  $Q^*$ -open in  $(Y, \sigma)$  and so again by hypothesis,  $g(f(G))$  is  $Q^*$ -open in  $(Z, \gamma)$ . i.e.,  $(g \circ f)(G)$  is  $Q^*$ -open in  $(Z, \gamma)$  and therefore  $(g \circ f)^{-1}$  is  $Q^*$ -irresolute. Hence  $g \circ f$  is a  $Q^*$ -homeomorphism. ■

**Example 3.3** Let  $X = Y = Z = \{a, b, c\}$ ,  $\tau = \{\phi, X, \{b\}, \{a, b\}, \{b, c\}\}$  and  $\sigma = \{\phi, Y, \{b\}, \{b, c\}\}$ ,  $\eta = \{\phi, Z, \{b, c\}\}$ . Then  $\phi, \{a\}, \{a, c\}$  are  $Q^*$ -closed in  $Y$ . Let  $f : X \rightarrow Y$  be the identity map. Then  $f$  and  $g$  are  $Q^*$ -homeomorphism. Here  $g \circ f$  is  $Q^*$ -continuous, since  $\{b, c\}$  is  $Q^*$ -open in  $Z$  and  $(g \circ f)^{-1}(\{b, c\}) = \{b, c\}$  is  $Q^*$ -open in  $X$ . Hence  $g \circ f$  is  $Q^*$ -homeomorphism.

**Remark 3.1.** We denote the family of all  $Q^*$ -homeomorphisms from  $(X, \tau)$  onto itself by  $Q^* - h(X, \tau)$ .

**Theorem 3.8** The set  $Q^* - h(X, \tau)$  is a group under the composition of maps.

**Proof.** Define a binary operation  $*$  :  $Q^* - h(X, \tau) \times Q^* - h(X, \tau) \rightarrow Q^* - h(X, \tau)$  by  $f * g = g \circ f$  for all  $f, g \in Q^* - h(X, \tau)$  and  $\circ$  is the usual operation of composition of maps. Then by Proposition 3.4,  $g \circ f \in Q^* - h(X, \tau)$ . We know that the composition of maps is associative and the identity map  $I : (X, \tau) \rightarrow (X, \tau)$  belonging to  $Q^* - h(X, \tau)$  serves as the identity element. If  $f \in Q^* - h(X, \tau)$ , then  $f^{-1} \in Q^* - h(X, \tau)$  such that  $f \circ f^{-1} = f^{-1} \circ f = I$  and so inverse exists for each element of  $Q^* - h(X, \tau)$ . Therefore,  $(Q^* - h(X, \tau), \circ)$  is a group under the operation of composition of maps. ■

**Theorem 3.9** Let  $f : (X, \tau) \rightarrow (Y, \sigma)$  be a  $Q^*$ -homeomorphism. Then  $f$  induces an isomorphism from the group  $Q^* - h(X, \tau)$  onto the group  $Q^* - h(Y, \sigma)$ .

**Proof.** Using the map  $f$ , we define a map  $\theta_f : Q^* - h(X, \tau) \rightarrow Q^* - h(Y, \sigma)$  by  $\theta_f(h) = f \circ h \circ f^{-1}$  for every  $h \in Q^* - h(X, \tau)$ . Then  $\theta_f$  is a bijection. Further, for all  $h_1, h_2 \in Q^* - h(X, \tau)$ ,  $\theta_f(h_1 \circ h_2) = f \circ (h_1 \circ h_2) \circ f^{-1} = (f \circ h_1 \circ f^{-1}) \circ (f \circ h_2 \circ f^{-1}) = \theta_f(h_1) \circ \theta_f(h_2)$ . Therefore,  $\theta_f$  is a homeomorphism and so it is an isomorphism induced by  $f$ . ■

**Proposition 3.4** For any two subsets  $A$  and  $B$  of  $(X, \tau)$ ,

- (i) If  $A \subset B$ , then  $Q^* - cl(A) \subset Q^* - cl(B)$ ,
- (ii)  $Q^* - cl(A \cap B) \subset Q^* - cl(A) \cap Q^* - cl(B)$ . ■

**Theorem 3.10.** If  $f : (X, \tau) \rightarrow (Y, \sigma)$  is a  $Q^*$ -homeomorphism, then  $Q^* - cl(f^{-1}(B)) = f^{-1}(Q^* - cl(B))$  for all  $B \subset Y$ .

**Proof.** Since  $f$  is a  $Q^*$ -homeomorphism,  $f$  is  $Q^*$ -irresolute. Since  $Q^* - cl(B)$  is a  $Q^*$ -closed set in  $(Y, \sigma)$ ,  $f^{-1}(Q^* - cl(B))$  is  $Q^*$ -closed in  $(X, \tau)$ . Now,  $f^{-1}(B) \subset f^{-1}(Q^* - cl(B))$  and so by Proposition 3.4,  $Q^* - cl(f^{-1}(B)) \subset f^{-1}(Q^* - cl(B))$ . Again since  $f$  is a  $Q^*$ -homeomorphism,  $f^{-1}$  is  $Q^*$ -irresolute. Since  $Q^* - cl(f^{-1}(B))$  is  $Q^*$ -closed in  $(X, \tau)$ ,  $(f^{-1})^{-1}(Q^* - cl(f^{-1}(B))) = f(Q^* - cl(f^{-1}(B)))$  is  $Q^*$ -closed in  $(Y, \sigma)$ . Now,  $B \subset (f^{-1})^{-1}(f^{-1}(B)) \subset (f^{-1})^{-1}(Q^* - cl(f^{-1}(B))) = f(Q^* - cl(f^{-1}(B)))$  and so

## International Journal for Research in Applied Science & Engineering Technology (IJRASET)

$Q^* \text{-cl}(B) \subset f ( Q^* \text{-cl}(f^{-1}(B)))$ . Therefore,  $f^{-1}( Q^* \text{-cl}(B)) \subset f^{-1}(f ( Q^* \text{-cl}(f^{-1}(B)))) \subset Q^* \text{-cl}(f^{-1}(B))$  and hence the equality holds. ■

Corollary 3.1 If  $f : (X, \tau) \rightarrow (Y, \sigma)$  is a  $Q^*$ -homeomorphism, then  $Q^* \text{-cl}(f(B)) = f ( Q^* \text{-cl}(B))$  for all  $B \subset X$ .

Proof. Since  $f : (X, \tau) \rightarrow (Y, \sigma)$  is a  $Q^*$ -homeomorphism,  $f^{-1} : (Y, \sigma) \rightarrow (X, \tau)$  is also a  $Q^*$ -homeomorphism. Therefore, by theorem 3.10 ,  $Q^* \text{-cl}((f^{-1})^{-1}(B)) = (f^{-1})^{-1}( Q^* \text{-cl}(B))$  for all  $B \subset X$ . i.e.,  $Q^* \text{-cl}(f(B)) = f ( Q^* \text{-cl}(B))$ . ■

Corollary 3.2 If  $f : (X, \tau) \rightarrow (Y, \sigma)$  is a  $Q^*$ -homeomorphism, then  $f ( Q^* \text{-int}(B)) = Q^* \text{-int}(f(B))$  for all  $B \subset X$ .

Proof. For any set  $B \subset X$ ,  $Q^* \text{-int}(B) = ( Q^* \text{-cl}(B^c))^c$ . Thus, by utilizing corollary 3.2 , we obtain  $f ( Q^* \text{-int}(B)) = f (( Q^* \text{-cl}(B^c))^c) = (f ( Q^* \text{-cl}(B^c)))^c = ( Q^* \text{-cl}(f(B^c)))^c = ( Q^* \text{-cl}((f(B))^c))^c = Q^* \text{-int}(f(B))$ . ■

Corollary 3.3 If  $f : (X, \tau) \rightarrow (Y, \sigma)$  is a  $Q^*$ -homeomorphism, then  $f^{-1}( Q^* \text{-int}(B)) = Q^* \text{-int}(f^{-1}(B))$  for all  $B \subset Y$ .

Proof. Since  $f^{-1} : (Y, \sigma) \rightarrow (X, \tau)$  is also a  $Q^*$ -homeomorphism, the proof follows from Corollary 3.2 . ■

### IV. $Q^*$ - HOMEOMORPHISM

Throughout this paper  $X$  and  $Y$  always represent nonempty topological spaces  $(X, \tau)$  and  $(Y, \sigma)$ . The family of all closed set in  $(X, \tau)$  is denoted by  $C(X)$ .

Theorem 4.1 Every  $Q^*$  homeomorphism is a homeomorphism.

Proof. Let  $f : (X, \tau) \rightarrow (Y, \sigma)$  be a  $Q^*$  - homeomorphism . Then  $f$  is bijective and both  $f$  and  $f^{-1}$  are  $Q^*$  - continuous . Since every  $Q^*$  continuous function is continuous we have  $f$  and  $f^{-1}$  are continuous . This shows that  $f$  is a homeomorphism. ■

Remark 4.1 The converse of the above theorem need not be true, as shown in the following example.

Example 4.1 Let  $X = Y = \{ a, b, c \}$ ,  $\tau = \{ \phi, X, \{ b \}, \{ a \}, \{ b, c \} \}$  and  $\sigma = \{ \phi, Y, \{ b \}, \{ b, c \} \}$ . Let  $f : X \rightarrow Y$  be the identity map . Therefore ,  $f$  is a homomorphism but not  $Q^*$  homeomorphism .

Definition 4.1 For a subset  $A$  of a space  $(X, \tau_1, \tau_2)$  we define the  $Q^*$  kernel of  $A$  ( briefly,  $Q^* \text{ker}(A)$  ) as follows :  $Q^* \text{ker}(A) = \cap \{ F : F \in Q^* O(X); A \subset F \}$  .  $A$  is said to be a  $Q^*$  -  $\Lambda$  set in  $X$  if  $A = Q^* \text{ker}(A)$  , or equivalently, if  $A$  is the intersection of  $Q^*$  open sets.  $A$  is said to be  $Q^*$   $\lambda$  - closed in  $X$  if it is the intersection of a  $Q^*$  -  $\Lambda$  set in  $X$  and clearly,  $Q^*$  -  $\Lambda$  sets and  $Q^*$  closed sets are  $Q^*$  -  $\lambda$  closed; complements of  $Q^*$  -  $\lambda$  closed sets in  $X$  are said to be  $Q^*$  -  $\lambda$  open in  $X$ .

Proposition 4.1 For a subset  $A$  of a space  $X$ , the following are equivalent:

- (i)  $A$  is  $Q^*$  -  $\lambda$  closed in  $X$  .
- (ii)  $A = L \cap Q^* \text{-cl}(A)$  , where  $L$  is a  $Q^*$  -  $\Lambda$  set in  $X$ .
- (iii)  $A = Q^* \text{-ker}(A) \cap Q^* \text{-cl}(A)$  . ■

Definition 4.2 A bijection  $f : X \rightarrow Y$  is called  $Q^*$ - homeomorphism , if  $f$  is  $Q^*$  irresolute and its inverse also  $Q^*$  irresolute .

Remark 4.2 We say that spaces  $X$  and  $Y$  are  $Q^*$ - homeomorphic if there exists a  $Q^*$  homeomorphism from  $(X, \tau)$  onto  $(Y, \sigma)$  . We denote the family of all  $Q^*$  homeomorphisms from  $(X, \tau)$  onto itself by  $Q^* \text{-H}(X)$  .

Theorem 4.2 Every  $Q^*$ - homeomorphism is a  $Q^*$  - homeomorphism.

Proof. Let  $f : (X, \tau) \rightarrow (Y, \sigma)$  be a  $Q^*$ - homeomorphism . Then  $f$  is bijective ,  $Q^*$  irresolute and  $f^{-1}$  is  $Q^*$  irresolute . Since every  $Q^*$  irresolute is  $Q^*$  continuous ,  $f$  and  $f^{-1}$  are  $Q^*$  continuous and so  $f$  is a  $Q^*$  homeomorphism. ■

Remark 4.3 The following example shows that the converse of the above theorem need not be true.

Example 4.3 Let  $X = Y = \{ a, b, c \}$ ,  $\tau = \{ \phi, X, \{ a, b \}, \{ b \} \}$  and  $\sigma = \{ \phi, Y, \{ b \}, \{ c \} \}$ . Let  $f : X \rightarrow Y$  be a map defined by  $f(a) = f(b) = c$  and  $f(c) = b$ . Therefore ,  $f$  is a  $Q^*$  - homomorphism but not  $Q^*$ - homeomorphism .

Remark 4.4 The concepts of homeomorphisms and  $Q^*$  - homeomorphism are independent.

Example 4.4 Let  $X = Y = \{ a, b, c \}$ ,  $\tau = \{ \phi, X, \{ b \}, \{ b, c \}, \{ a \} \}$  and  $\sigma = \{ \phi, Y, \{ a \}, \{ b \} \}$ . Let  $f : X \rightarrow Y$  be a identity map . Then  $f$  is homeomorphism but not  $Q^*$  homeomorphism.

Lemma 4.1 A function  $f : X \rightarrow Y$  is  $Q^*$  - irresolute if and only if  $f^{-1}(V)$  is a  $Q^*$  - closed set in  $X$  for every  $Q^*$  - closed set  $V$  in  $Y$  .

Theorem 4.3 If  $f : (X, \tau) \rightarrow (Y, \sigma)$  is a  $Q^*$ - homeomorphism, then  $Q^* \text{-cl}(f^{-1}(B)) = f^{-1}( Q^* \text{-cl}(B))$  for every  $B \subseteq Y$ .

Proof. Let  $f : (X, \tau) \rightarrow (Y, \sigma)$  be a  $Q^*$  - homeomorphism. Then by Definition 4.2, both  $f$  and  $f^{-1}$  are  $Q^*$  - irresolute and  $f$  is bijective. Let  $B \subseteq Y$ . Since  $Q^* \text{-cl}(B)$  is a  $Q^*$ -closed set in  $(Y, \sigma)$ , using Lemma 4.1,  $f^{-1}( Q^* \text{-cl}(B))$  is  $Q^*$ -closed in  $(X, \tau)$ . But

## International Journal for Research in Applied Science & Engineering Technology (IJRASET)

$Q^*\text{-cl}(f^{-1}(B))$  is the smallest  $Q^*$ -closed set containing  $f^{-1}(B)$ .

Therefore  $Q^*\text{-cl}(f^{-1}(B)) \subseteq f^{-1}(Q^*\text{-cl}(B))$ .  $\rightarrow (1)$ .

Again,  $Q^*\text{-cl}(f^{-1}(B))$  is  $Q^*$ -closed in  $(X, \tau)$ . Since  $f^{-1}$  is  $Q^*$ -irresolute,  $f(Q^*\text{-cl}(f^{-1}(B)))$  is  $Q^*$ -closed in  $(Y, \sigma)$ . Now,  $B = f(f^{-1}(B)) \subseteq f(Q^*\text{-cl}(f^{-1}(B)))$ . Since  $f(Q^*\text{-cl}(f^{-1}(B)))$  is  $Q^*$ -closed and  $Q^*\text{-cl}(B)$  is the smallest  $Q^*$ -closed set containing  $B$ ,  $Q^*\text{-cl}(B) \subseteq f(Q^*\text{-cl}(f^{-1}(B)))$  that implies  $f^{-1}(Q^*\text{-cl}(B)) \subseteq f^{-1}(f(Q^*\text{-cl}(f^{-1}(B)))) = Q^*\text{-cl}(f^{-1}(B))$ .

That is,  $f^{-1}(Q^*\text{-cl}(B)) \subseteq Q^*\text{-cl}(f^{-1}(B)) \rightarrow (2)$

From (1) and (2),  $Q^*\text{-cl}(f^{-1}(B)) = f^{-1}(Q^*\text{-cl}(B))$ . ■

Corollary 4.1 If  $f: (X, \tau) \rightarrow (Y, \sigma)$  is a  $Q^*$ -homeomorphism, then  $Q^*\text{-cl}(f(B)) = f(Q^*\text{-cl}(B))$  for every  $B \subseteq X$ .

Proof. Let  $f: (X, \tau) \rightarrow (Y, \sigma)$  be a  $Q^*$ -homeomorphism. Since  $f$  is  $Q^*$ -homeomorphism,  $f^{-1}$  is also a  $Q^*$ -homeomorphism. Therefore by Theorem 4.3, it follows that  $Q^*\text{-cl}(f(B)) = f(Q^*\text{-cl}(B))$  for every  $B \subseteq X$ . ■

Corollary 4.2 If  $f: (X, \tau) \rightarrow (Y, \sigma)$  is a  $Q^*$ -homeomorphism, then  $f(Q^*\text{-int}(B)) = Q^*\text{-int}(f(B))$  for every  $B \subseteq X$ .

Proof. Let  $f: (X, \tau) \rightarrow (Y, \sigma)$  be a  $Q^*$ -homeomorphism. For any set  $B \subseteq X$ ,  $Q^*\text{-int}(B) = (Q^*\text{-cl}(B^c))^c$ .  $f(Q^*\text{-int}(B)) = f((Q^*\text{-cl}(B^c))^c) = (f(Q^*\text{-cl}(B^c)))^c$ . Then using Corollary 4.2, we see that  $f(Q^*\text{-int}(B)) = (Q^*\text{-cl}(f(B^c)))^c = Q^*\text{-int}(f(B))$ . ■

Corollary 4.3 If  $f: (X, \tau) \rightarrow (Y, \sigma)$  is a  $Q^*$ -homeomorphism, then for every  $B \subseteq Y$ ,  $f^{-1}(Q^*\text{-int}(B)) = Q^*\text{-int}(f^{-1}(B))$ .

Proof. Let  $f: (X, \tau) \rightarrow (Y, \sigma)$  be a  $Q^*$ -homeomorphism. Since  $f$  is  $Q^*$ -homeomorphism,  $f^{-1}$  is also a  $Q^*$ -homeomorphism. Therefore by Corollary 4.2,  $f^{-1}(Q^*\text{-int}(B)) = Q^*\text{-int}(f^{-1}(B))$  for every  $B \subseteq Y$ . ■

Theorem 4.3 If  $f: (X, \tau) \rightarrow (Y, \sigma)$  and  $g: (Y, \sigma) \rightarrow (Z, \gamma)$  are  $Q^*$ -homeomorphisms, then their composition  $g \circ f: (X, \tau) \rightarrow (Z, \gamma)$  is also  $Q^*$ -homeomorphism.

Proof. Let  $U$  be a  $Q^*$ -open set in  $(Z, \gamma)$ . Since  $g$  is  $Q^*$ -homeomorphism,  $g$  is  $Q^*$ -irresolute and so  $g^{-1}(U)$  is  $Q^*$ -open in  $(Y, \sigma)$ . Since  $f$  is  $Q^*$ -homeomorphism,  $f$  is  $Q^*$ -irresolute and so  $f^{-1}(g^{-1}(U)) = (g \circ f)^{-1}(U)$  is  $Q^*$ -open in  $(X, \tau)$ . This implies that  $g \circ f$  is  $Q^*$ -irresolute. Again let  $G$  be  $Q^*$ -open in  $(X, \tau)$ . Since  $f$  is  $Q^*$ -homeomorphism,  $f^{-1}$  is  $Q^*$ -irresolute and so  $(f^{-1})^{-1}(G) = f(G)$  is  $Q^*$ -open in  $(Y, \sigma)$ . Since  $g$  is  $Q^*$ -homeomorphism,  $g^{-1}$  is  $Q^*$ -irresolute and so  $(g^{-1})^{-1}(f(G)) = g(f(G)) = (g \circ f)(G) = ((g \circ f)^{-1})^{-1}(G)$  is  $Q^*$ -open in  $(Z, \gamma)$ . This implies that  $(g \circ f)^{-1}$  is  $Q^*$ -irresolute. Since  $f$  and  $g$  are  $Q^*$ -homeomorphism  $f$  and  $g$  are bijective and so  $g \circ f$  is bijective. This completes the proof. ■

Theorem 4.4 The set  $Q^*H(X, \tau)$  is a group under composition of functions.

Proof. Let  $f, g \in Q^*H(X, \tau)$ . Then  $f \circ g \in Q^*H(X, \tau)$  by Theorem 3.17. Since  $f$  is bijective,  $f^{-1} \in Q^*H(X, \tau)$ . This completes the proof. ■

Theorem 4.5 If  $f: (X, \tau) \rightarrow (Y, \sigma)$  is a  $Q^*$ -homeomorphism, then  $f$  induces an isomorphism from the group  $Q^*H(X, \tau)$  onto the group  $Q^*H(Y, \sigma)$ .

Proof. Let  $f \in Q^*H(X, \tau)$ . Then define a map  $\psi_f: Q^*H(X, \tau) \rightarrow Q^*H(Y, \sigma)$  by  $\psi_f(h) = f \circ h \circ f^{-1}$  for every  $h \in Q^*H(X, \tau)$ . Let  $h_1, h_2 \in Q^*H(X, \tau)$ .

$$\begin{aligned} \text{Then } \psi_f(h_1 \circ h_2) &= f \circ (h_1 \circ h_2) \circ f^{-1} \\ &= f \circ (h_1 \circ f^{-1} \circ f \circ h_2) \circ f^{-1} \\ &= (f \circ h_1 \circ f^{-1}) \circ (f \circ h_2 \circ f^{-1}) \\ &= \psi_f(h_1) \circ \psi_f(h_2). \end{aligned}$$

Since  $\psi_f(f^{-1} \circ h \circ f) = h_1 \psi_f$  is onto. Now,  $\psi_f(h) = I$  implies  $f \circ h \circ f^{-1} = I$ . That implies  $h = I$ . This proves that  $\psi_f$  is one-one. This shows that  $\psi_f$  is an isomorphism. ■

Theorem 4.6 Every  $Q^*$ -open set is  $\Lambda_r$ -open.

Theorem 4.7 Every  $Q^*$ -homeomorphism is a  $\Lambda_r$ -homeomorphism.

Proof. Let  $f: (X, \tau) \rightarrow (Y, \sigma)$  be a  $Q^*$ -homeomorphism. Then  $f$  is bijective and both  $f$  and  $f^{-1}$  are  $Q^*$ -continuous. Since every  $Q^*$ -continuous function is  $\Lambda_r$ -continuous,  $f$  and  $f^{-1}$  are  $\Lambda_r$ -continuous. This shows that  $f$  is a  $\Lambda_r$ -homeomorphism.


Remark 4.5 The converse of the above theorem need not be true, as shown in the following example.

Example 4.4 Let  $X = Y = \{a, b, c\}$ ,  $\tau = \{X, \emptyset, \{a\}, \{b\}, \{a, b\}, \{b, c\}\}$  and  $\sigma = \{Y, \emptyset, \{b\}, \{c\}, \{b, c\}\}$ . Then  $\Lambda_r O(X, \tau) = \tau$  and  $\Lambda_r O(Y, \sigma) = \{Y, \emptyset, \{b\}, \{c\}, \{b, c\}, \{a, c\}, \{a, b\}\}$ . Let  $f: (X, \tau) \rightarrow (Y, \sigma)$  be defined by  $f(a) = c$ ,  $f(b) = b$  and  $f(c) = a$ . Then  $f$  is  $\Lambda_r$ -homeomorphism. Since  $f(\{b, c\}) = \{a, b\}$  is not  $Q^*$ -open in  $(Y, \sigma)$ ,  $f^{-1}$  is not  $Q^*$ -continuous that implies  $f$  is not a  $Q^*$ -homeomorphism.

### V. CONCLUSION

## International Journal for Research in Applied Science & Engineering Technology (IJRASET)

In this paper, we introduce two classes of maps called  $Q^*$  - homeomorphisms and  $Q^*$ -homeomorphisms and study their properties. From all of the above statements, we have the following diagram:



### REFERENCES

- [ 1 ] Chandrasekhara Rao.K , On semi homeomorphisms , International ResearchJournal of Pure Algebra , 3 ( 6 ) ( 2013 ) , 222 - 223 .
- [ 2 ] Crossley S.G and Hildebrand S. K, Semi-closure, Texas J. Sci. 22, pp. 99-112, (1971).
- [ 3 ] Devi R and Balachandran K (2001) Some generalizations of  $\alpha$  - homeomorphisms in topological spaces. Indian J. Pure appl. Math., 32(4) : 551-563.
- [ 4 ] Devi R, Balachandran K and Maki H, Semi-generalized homeomorphisms and generalized semi-homeomorphisms in topological spaces, Indian J. Pure Appl. Math., 26, pp. 271-284, (1995).
- [ 5 ] Devi.R and Vigneshwaran.M (2010)  $\ast g\alpha$  c-homeomorphisms in topological spaces. International Journal of Computer Applications 3( 6 ) : 17-19.
- [ 6 ] Devi R, Maki H and Balachandran K, Semi-generalized closed maps and generalized semi-closed maps, Mem. Fac. Kochi Univ. Ser. A. Math., 14, pp. 41-54, (1993). 18 M.
- [ 7 ] Garg. G . L and Sivaraj. D , Pre Semi Closed Mappings , Periodica Mathematica , Hungeriaca , 19 ( 2 ) , ( 1988 ) , 97 - 106 .
- [ 8 ] Janich K, Topologie, Springer-Verlag, Berlin, 1980 (English translation).
- [ 9 ] Jeyanthi , M.J. Pious Missier.S and Thangavelu.P ,  $\Lambda_r$ - homeomorphisms and  $\Lambda_r^*$  - homeomorphisms , Journal of Mathematical Sciences & Computer Applications 1 ( 2): 48-55, 2011.
- [ 10 ] Kokilavani.V and Basker.V , On  $\alpha\delta$  hms in topological spaces , international journal of advanced scientific and technical research , Issue 2 , volume 3 , 2012 .
- [ 11 ] Maki H, Sundram P and Balachandran K, On generalized homeomorphisms in topological spaces, Bull. Fukuoka Univ. Ed. Part III, 40, pp. 13-21, (1991).
- [ 12 ] Nagaveni.N , Studies on Generalizations of Homeomorphisms in Topological Spaces, Ph.D.,Thesis, Bharathiar University, Coimbatore(1999).
- [ 13 ] Malghan S. R, Generalized closed maps, J. Karnataka Univ. Sci., 27,pp. 82-88, (1982).
- [ 14 ] Mashhour A. S, Abd. El-Monsef M. E and El-Deeb S. N, On pre continuous and weak pre continuous mappings, Proc. Math. Phys. Soc. Egypt, 53, pp. 47-53, (1982).
- [ 15 ] M. Murugalingam and N. Lalitha , “ Q star sets “ , Bulletin of pure and applied Sciences , Volume 29E Issue 2 ( 2010 ) p. 369 - 376 .
- [ 16 ] M. Murugalingam and N. Lalitha , “  $Q^*$  sets in various spaces “ , Bulletin of pure and applied Sciences , Volume 3E Issue 2 ( 2011 ) p. 267 - 277 .
- [ 17 ] P.Padma and S .Udayakumar , “  $\tau_1 \tau_2 - Q^*$  continuous maps in bitopological spaces “ Asian Journal of Current Engineering and Mathematics , 1 : 4 Jul - Aug ( 2012 ) , 227 - 229 .
- [ 18 ] P . Padma and S .Udayakumar ,” Pairwise  $Q^*$ Homeomorphism in bitopological spaces , Research journal of Pure Algebra - 3 ( 3 ) , 2013 , page 111 - 118 .
- [ 19 ] Rajesh N and Ekici E, On a decomposition of  $T_{1/2}$ -spaces, Math.Maced. (to appear).
- [ 20 ] Reilly. I. L and Vamanamurthy .M.K , On semi compact spaces , Bull . Malaysian Math. Soc., ( 7 ) 2 ( 1984 ) , 61 - 67 .
- [ 21 ] Sreeja.D and Janaki.C , A New Type of Homeomorphism in Bitopological Spaces , International Journal of Scientific and Research Publications, Volume 2, Issue 7, July 2012
- [ 22 ] Stone M, Application of the theory of Boolean rings to general topology, Trans. Amer. Math. Soc. 41, pp. 374-481, (1937).
- [ 23 ] Vadivel.A and VairamanickamK ,  $\Lambda_r$ -homeomorphisms in topological spaces.International Journal of Math. Analysis, 4(18) : 881-890 , 2010 .





10.22214/IJRASET



45.98



IMPACT FACTOR:  
7.129



IMPACT FACTOR:  
7.429



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24\*7 Support on Whatsapp)