



IJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 7 Issue: VI Month of publication: June 2019

DOI: <http://doi.org/10.22214/ijraset.2019.6421>

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

Sufficient Condition for Wavelet Frame on Positive Half-Line

Abdullah¹, Afroz²

¹College of Engineering, Qassim University, Buraidah 51452, Al-Qassim, Saudi Arabia.

¹Department of Mathematics, Zakir Husain Delhi College, University of Delhi, JLN Marg, New Delhi-110 002, India.

²Department of Mathematics, School of sciences, Maulana Azad National Urdu University, Hyderabad-500032, India.

Abstract: In this paper, we present p -wavelet frames on positive half-line using Walsh-Fourier transform and also prove a sufficient condition for the system $\{\psi_{j,k}: j \in \mathbb{Z}, k \in \mathbb{Z}^+\}$ to be a wavelet frame in $L^2(\mathbb{R}^+)$.

I. INTRODUCTION

In recent years, wavelets have been generalized in many different settings, for example locally compact abelian group, abstract Hilbert spaces, locally compact Cantor dyadic group, Vilenkin group, local fields and positive half-line. In this paper our interest is in positive half-line. Farkov [12] has given general construction of compactly supported orthogonal p -wavelets in $L^2(\mathbb{R}^+)$. Farkov et al. [13] gave an algorithm for biorthogonal wavelets related to Walsh functions on positive half line. Shah and Debnath [19], constructed wavelet frame packets on the positive half-line \mathbb{R}^+ using the splitting trick for frames. Abdullah [3] has given characterization of nonuniform wavelet sets on positive half-line. The characterization of wavelets on positive half line by means of two basic equations in the Fourier domain established in [1]. A constructive procedure for constructing tight wavelet frames on positive half-line using extension principles was recently considered by Shah in [17], in which he has pointed out a method for constructing affine frames in $L^2(\mathbb{R}^+)$. Moreover, the author has established sufficient conditions for a finite number of functions to form a tight affine frames for $L^2(\mathbb{R}^+)$.

In this paper, we investigate wavelet frames on positive half-line \mathbb{R}^+ . The main result presented in this paper is sufficient condition for the system $\{\psi_{j,k}(x): j \in \mathbb{Z}, k \in \mathbb{Z}^+\}$ to be a wavelet frame in $L^2(\mathbb{R}^+)$.

II. NOTATIONS AND PRELIMINARIES

Let p be a fixed natural number greater than 1. As usual, let $\mathbb{R}^+ = [0, \infty)$, $\mathbb{N} = \{1, 2, \dots\}$ and $\mathbb{Z}^+ = \{0, 1, \dots\}$. Denote by $[x]$ the integer part of x . For $x \in \mathbb{R}^+$ and for any positive integer j , we set

$$x_j = [p^j x] \pmod{p}, \quad x_{-j} = [p^{1-j} x] \pmod{p}, \quad (2.1)$$

where $x_j, x_{-j} \in \{0, 1, \dots, p-1\}$.

Consider the addition defined on \mathbb{R}^+ as follows:

$$x \oplus y = \sum_{j < 0} \xi_j p^{-j-1} + \sum_{j > 0} \xi_j p^{-j} \quad (2.2)$$

with

$$\xi_j = x_j + y_j \pmod{p}, \quad j \in \mathbb{Z} \setminus \{0\}, \quad (2.3)$$

where $\xi_j \in \{0, 1, \dots, p-1\}$ and x_j, y_j are calculated by (2.1). Moreover, we write $z = x \ominus y$ if $z \oplus y = x$, where \ominus denotes subtraction modulo p in \mathbb{R}^+ .

For $x \in [0, 1)$, let $r_0(x)$ be given by

$$r_0(x) = \begin{cases} 1, & x \in \left[0, \frac{1}{p}\right) \\ \epsilon_p^j, & x \in [jp^{-1}, (j+1)p^{-1}), j = 1, 2, \dots, p-1, \end{cases} \quad (2.4)$$

where $\epsilon_p = \exp\left(\frac{2\pi i}{p}\right)$. The extension of the function r_0 to \mathbb{R}^+ is defined by the equality $r_0(x+1) = r_0(x)$, $x \in \mathbb{R}^+$. Then the generalized Walsh functions $\{\omega_m(x)\}_{m \in \mathbb{Z}^+}$ are defined by

$$\omega_0(x) = 1, \quad \omega_m(x) = \prod_{j=0}^k (r_0(p^j x))^{\mu_j},$$

where $m = \sum_{j=0}^k \mu_j p^j$, $\mu_j \in \{0, 1, 2, \dots, p-1\}$, $\mu_k \neq 0$.

For $x, \omega \in \mathbb{R}^+$, let

$$\chi(x, \omega) = \exp\left(\frac{2\pi i}{p} \sum_{j=1}^{\infty} (x_j \omega_{-j} + x_{-j} \omega_j)\right), \quad (2.5)$$

where x_j and ω_j are calculated by (2.1).

We observe that

$$\chi\left(x, \frac{m}{p^{n-1}}\right) = \chi\left(\frac{x}{p^{n-1}}, m\right) = \omega_m\left(\frac{x}{p^{n-1}}\right) \quad \forall x \in [0, p^{n-1}), \quad m \in \mathbb{Z}^+.$$

The Walsh-Fourier transform of a function $f \in L^1(\mathbb{R}^+)$ is defined by

$$\tilde{f}(\omega) = \int_{\mathbb{R}^+} f(x) \overline{\chi(x, \omega)} dx, \quad (2.6)$$

where $\chi(x, \omega)$ is given by (2.5).

If $f \in L^2(\mathbb{R}^+)$ and

$$J_a f(\omega) = \int_0^a f(x) \overline{\chi(x, \omega)} dx \quad (a < \infty), \quad (2.7)$$

then \tilde{f} is defined as limit of $J_a f$ in $L^2(\mathbb{R}^+)$ as $a \rightarrow \infty$.

The properties of Walsh-Fourier transform are quite similar to the classical Fourier transform. It is known that systems $\{\chi(\alpha, \cdot)\}_{\alpha=0}^{\infty}$ and $\{\chi(\cdot, \alpha)\}_{\alpha=0}^{\infty}$ are orthonormal bases in $L^2(0,1)$. Let us denote by $\{\omega\}$ the fractional part of ω . For $l \in \mathbb{Z}^+$, we have $\chi(l, \omega) = \chi(l, \{\omega\})$.

If $x, y, \omega \in \mathbb{R}^+$ and $x \oplus y$ is p -adic irrational, then

$$\chi(x \oplus y, \omega) = \chi(x, \omega) \chi(y, \omega), \quad \chi(x \ominus y, \omega) = \chi(x, \omega) \overline{\chi(y, \omega)}. \quad (2.8)$$

By a p -adic interval of range n in $[0,1)$, we mean intervals of the form

$$I_{(n)}^k = [k2^{-n}, (k+1)2^{-n}), \quad k \in \mathbb{Z}^+.$$

It is easy to verify that

$$I_{(n)}^k \cap I_{(n)}^l = \emptyset, \quad k \neq l \text{ and } \bigcup_{k=0}^{2^n-1} I_{(n)}^k = [0,1).$$

For each $x \in [0,1)$ and $n \in \mathbb{N}$, we denote the p -adic interval of length p^{-n} which contains x by $I_n(x)$. Thus

$$I_n(x) = I_{(n)}^k(x),$$

where $0 \leq k < p^n$ is uniquely determined by relationship $x \in I_n(x)$.

III. MAIN RESULTS

Let

$$\psi \in L^2(\mathbb{R}^+), \quad \psi_{j,k}(x) = p^{j/2} \psi(p^j x \ominus k), \quad j \in \mathbb{Z}, k \in \mathbb{Z}^+. \quad (3.1)$$

By taking Walsh-Fourier transform, we obtain

$$\hat{\psi}_{j,k}(\xi) = p^{-j/2} \overline{\chi(k, p^{-j} \xi)} \hat{\psi}(p^{-j} \xi). \quad (3.2)$$

Then, we call the function system $\{\psi_{j,k}(x) : j \in \mathbb{Z}, k \in \mathbb{Z}^+\}$ a wavelet frame for $L^2(\mathbb{R}^+)$ if there are two constants A and B , $0 < A \leq B < \infty$ such that

$$A \|f\|^2 \leq \sum_{j \in \mathbb{Z}} \sum_{k \in \mathbb{Z}^+} |\langle f, \psi_{j,k} \rangle|^2 \leq B \|f\|^2, \quad (3.3)$$

for all $f \in L^2(\mathbb{R}^+)$. The largest A and the smallest B for which (3.3) holds are called frame bounds. A frame is a tight frame if A and B are chosen so that $A = B$ and is a normalized tight frame if $A = B = 1$.

For $j \in \mathbb{Z}$, $l \in \mathbb{Z}^+$, we have

$$\int_{p^j l}^{p^{j(l+1)}} \omega_k(p^{-j} \xi) d\xi = \int_0^{p^j} \omega_k(p^{-j}(\xi + p^j l)) d\xi = \int_0^{p^j} \omega_k(p^{-j} \xi) d\xi.$$

Let $f \in L^2(\mathbb{R}^+)$ and $\psi \in L^2(\mathbb{R}^+)$, then

$$\langle f, \psi_{j,k} \rangle = p^{-j/2} \int_0^{p^j} \left[\sum_{l \in \mathbb{Z}^+} \hat{f}(\xi \oplus p^j l) \overline{\hat{\psi}(p^{-j} \xi \oplus l)} \right] \omega_k(p^{-j} \xi) d\xi.$$

Applying Parseval's formula and the fact that $\{\omega_n : n \geq 0\}$ forms an orthonormal basis for $L^2[0,1)$, we obtain

$$\sum_{k \in \mathbb{Z}^+} |f, \psi_{j,k}|^2 = \int_{\mathbb{R}^+} \overline{\hat{f}(\xi)} \hat{\psi}(p^{-j} \xi) \left\{ \sum_{l \in \mathbb{Z}^+} \hat{f}(\xi \oplus p^j l) \overline{\hat{\psi}(p^{-j} \xi \oplus l)} \right\} d\xi. \quad (3.4)$$

Now, we first prove a lemma, which will be used in the proofs of the main results.

1) *Lemma 3.1.* Let $\mathcal{D} = \{f \in L^2(\mathbb{R}^+) : \text{supp } \hat{f} \subset \mathbb{R}^+ \setminus \{0\}\}$ and let $f \in \mathcal{D}$ and $\psi \in L^2(\mathbb{R}^+)$. If $\text{ess sup}_{\xi \in \mathbb{R}^+} \sum_{j \in \mathbb{Z}} |\hat{\psi}(p^{-j} \xi)|^2 < +\infty$, then

$$\sum_{j \in \mathbb{Z}} \sum_{k \in \mathbb{Z}^+} |f, \psi_{j,k}|^2 = \int_{\mathbb{R}^+} |\hat{f}(\xi)|^2 \sum_{j \in \mathbb{Z}} |\hat{\psi}(p^{-j} \xi)|^2 d\xi + R_\psi(f), \quad (3.5)$$

where

$$R_\psi(f) = \sum_{j \in \mathbb{Z}} \sum_{l \in \mathbb{N}} \int_{\mathbb{R}^+} \overline{\hat{f}(\xi)} \hat{\psi}(p^{-j} \xi) \hat{f}(\xi \oplus p^j l) \overline{\hat{\psi}(p^{-j} \xi \oplus l)} d\xi. \quad (3.6)$$

Furthermore, the iterated series in (3.6) is absolutely convergent.

Proof. From (3.4), we have

$$\begin{aligned} \sum_{j \in \mathbb{Z}} \sum_{k \in \mathbb{Z}^+} |f, \psi_{j,k}|^2 &= \sum_{j \in \mathbb{Z}} \int_{\mathbb{R}^+} \{ |\hat{f}(\xi)|^2 |\hat{\psi}(p^{-j} \xi)|^2 + \overline{\hat{f}(\xi)} \hat{\psi}(p^{-j} \xi) \\ &\quad \times \sum_{l \in \mathbb{Z}^+} \hat{f}(\xi \oplus p^j l) \overline{\hat{\psi}(p^{-j} \xi \oplus l)} \} d\xi \\ &= \sum_{j \in \mathbb{Z}} \int_{\mathbb{R}^+} |\hat{f}(\xi)|^2 |\hat{\psi}(p^{-j} \xi)|^2 d\xi + R_\psi(f). \end{aligned}$$

Since $\text{ess sup}_{\xi \in \mathbb{R}^+} \sum_{j \in \mathbb{Z}} |\hat{\psi}(p^{-j} \xi)|^2 < +\infty$, and therefore, by the Levi lemma, we obtain

$$\sum_{j \in \mathbb{Z}} \sum_{k \in \mathbb{Z}^+} |f, \psi_{j,k}|^2 = \int_{\mathbb{R}^+} |\hat{f}(\xi)|^2 \sum_{j \in \mathbb{Z}} |\hat{\psi}(p^{-j} \xi)|^2 d\xi + R_\psi(f).$$

Now we claim that the iterated series in (3.6) is absolutely convergent. To do this, let

$$\begin{aligned} I &= \sum_{j \in \mathbb{Z}} \sum_{l \in \mathbb{N}} \int_{\mathbb{R}^+} |\hat{f}(\xi) \hat{\psi}(p^{-j} \xi) \hat{f}(\xi \oplus p^j l) \hat{\psi}(p^{-j} \xi \oplus l)| d\xi \\ &= \sum_{j \in \mathbb{Z}} \sum_{l \in \mathbb{N}} p^j \int_{\mathbb{R}^+} |\hat{f}(p^j \xi) \hat{\psi}(\xi) \hat{f}(p^j(\xi \oplus l)) \hat{\psi}(\xi \oplus l)| d\xi. \end{aligned}$$

Note that

$$|\hat{\psi}(\xi) \hat{\psi}(\xi \oplus l)| \leq \frac{1}{2} (|\hat{\psi}(\xi)|^2 + |\hat{\psi}(\xi \oplus l)|^2).$$

Therefore, it suffices to prove that

$$\sum_{j \in \mathbb{Z}} \sum_{l \in \mathbb{N}} p^j \int_{\mathbb{R}^+} |\hat{f}(p^j \xi) \hat{f}(p^j \xi \oplus p^j l)| |\hat{\psi}(\xi)|^2 d\xi < \infty. \quad (3.7)$$

Since $l \neq 0$ ($l \in \mathbb{N}$) and $f \in \mathcal{D}$, there exists $J > 0$ such that for all $|j| > J$,

$$\hat{f}(p^j \xi) \hat{f}(p^j \xi \oplus p^j l) = 0.$$

On the other hand, for each fixed $|j| \leq J$ and $\xi \in \mathbb{R}^+$, there exists a constant L such that for all $l > L$,

$$\hat{f}(p^j \xi \oplus p^j l) = 0.$$

Thus, it follows that only a finite number of terms of the iterated series in (3.7) are non-zero. Consequently, there exists a constant C such that

$$I \leq C \|f\|_{\infty}^2 \|\hat{\psi}\|_2^2.$$

This shows that the iterated series in (3.6) is absolutely convergent.

The following necessary condition for the system $\{\psi_{j,k} : j \in \mathbb{Z}, k \in \mathbb{Z}^+\}$ to be a frame was proved by Abdullah [2].

2) *Theorem 3.2.* If $\{\psi_{j,k}(x) : j \in \mathbb{Z}, k \in \mathbb{Z}^+\}$ is a wavelet frame for $L^2(\mathbb{R}^+)$ with bounds A and B , then

$$A \leq \sum_{j \in \mathbb{Z}} |\hat{\psi}(p^j \xi)|^2 \leq B, \quad a.e. \xi \in \mathbb{R}^+.$$

Now, in the following theorem we establish a sufficient condition of the system (3.1) to be a frame in $L^2(\mathbb{R}^+)$

3) *Theorem 3.3.* Let $\psi \in L^2(\mathbb{R}^+)$ such that

$$A(\psi) = \text{ess inf} \left\{ \sum_{j \in \mathbb{Z}} |\hat{\psi}(p^{-j} \xi)|^2 - \sum_{j \in \mathbb{Z}} \sum_{l \in \mathbb{N}} |\hat{\psi}(p^{-j} \xi) \overline{\hat{\psi}(p^{-j} \xi \oplus l)}|^2 : \xi \in \mathbb{R}^+ \right\} > 0,$$

$$B(\psi) = \text{ess sup} \left\{ \sum_{j \in \mathbb{Z}} \sum_{k \in \mathbb{Z}^+} |\hat{\psi}(p^{-j} \xi) \overline{\hat{\psi}(p^{-j} \xi \oplus k)}|^2 : \xi \in \mathbb{R}^+ \right\} < +\infty.$$

Then $\{\psi_{j,k}(x) : j \in \mathbb{Z}, k \in \mathbb{Z}^+\}$ is a wavelet frame for $L^2(\mathbb{R}^+)$ with frame bounds $A(\psi)$ and $B(\psi)$.

Proof. We can estimate $R_{\psi}(f)$ by rearranging the series in (3.6),

$$\begin{aligned} |R_{\psi}(f)| &= \left| \sum_{j \in \mathbb{Z}} \sum_{l \in \mathbb{N}} \int_{\mathbb{R}^+} \overline{\hat{f}(\xi)} \hat{\psi}(p^{-j} \xi) \hat{f}(\xi \oplus p^j l) \overline{\hat{\psi}(p^{-j} \xi \oplus l)} d\xi \right| \\ &\leq \sum_{j \in \mathbb{Z}} \sum_{l \in \mathbb{N}} \left\{ \int_{\mathbb{R}^+} |\hat{f}(\xi)|^2 |\hat{\psi}(p^{-j} \xi) \overline{\hat{\psi}(p^{-j} \xi \oplus l)}| d\xi \right\}^{\frac{1}{2}} \\ &\quad \left\{ \int_{\mathbb{R}^+} |\hat{f}(\xi \oplus p^j l)|^2 |\hat{\psi}(p^{-j} \xi) \overline{\hat{\psi}(p^{-j} \xi \oplus l)}| d\xi \right\}^{\frac{1}{2}} \\ &\leq \sum_{j \in \mathbb{Z}} \left\{ \sum_{l \in \mathbb{N}} \int_{\mathbb{R}^+} |\hat{f}(\xi)|^2 |\hat{\psi}(p^{-j} \xi) \overline{\hat{\psi}(p^{-j} \xi \oplus l)}| d\xi \right\}^{\frac{1}{2}} \\ &\quad \left\{ \sum_{l \in \mathbb{N}} \int_{\mathbb{R}^+} |\hat{f}(\xi \oplus p^j l)|^2 |\hat{\psi}(p^{-j} \xi) \overline{\hat{\psi}(p^{-j} \xi \oplus l)}| d\xi \right\}^{\frac{1}{2}} \\ &= \sum_{j \in \mathbb{Z}} \left\{ \sum_{l \in \mathbb{N}} \int_{\mathbb{R}^+} |\hat{f}(\xi)|^2 |\hat{\psi}(p^{-j} \xi) \overline{\hat{\psi}(p^{-j} \xi \oplus l)}| d\xi \right\}^{\frac{1}{2}} \\ &\quad \left\{ \sum_{l \in \mathbb{N}} \int_{\mathbb{R}^+} |\hat{f}(\omega)|^2 |\hat{\psi}(p^{-j} \omega \ominus l) \overline{\hat{\psi}(p^{-j} \omega)}| d\xi \right\}^{\frac{1}{2}}. \end{aligned}$$

Therefore

$$|R_{\psi}(f)| \leq \sum_{j \in \mathbb{Z}} \sum_{l \in \mathbb{N}} \int_{\mathbb{R}^+} |\hat{f}(\xi)|^2 |\hat{\psi}(p^{-j} \xi) \overline{\hat{\psi}(p^{-j} \xi \oplus l)}| d\xi.$$

By Levi Lemma, we have

$$|R_\psi(f)| \leq \int_{\mathbb{R}^+} |\hat{f}(\xi)|^2 \left\{ \sum_{j \in \mathbb{Z}} \sum_{l \in \mathbb{N}} |\hat{\psi}(p^{-j}\xi) \overline{\hat{\psi}(p^{-j}\xi \oplus l)}| \right\} d\xi.$$

Applying (3.5), we have

$$\int_{\mathbb{R}^+} |\hat{f}(\xi)|^2 \left\{ \sum_{j \in \mathbb{Z}} |\hat{\psi}(p^{-j}\xi)|^2 - \sum_{j \in \mathbb{Z}} \sum_{l \in \mathbb{N}} |\hat{\psi}(p^{-j}\xi) \overline{\hat{\psi}(p^{-j}\xi \oplus l)}| \right\} d\xi \leq \sum_{j \in \mathbb{Z}} \sum_{k \in \mathbb{Z}^+} |f, \psi_{j,k}|^2, \quad (3.8)$$

and

$$\sum_{j \in \mathbb{Z}} \sum_{k \in \mathbb{Z}^+} |f, \psi_{j,k}|^2 \leq \int_{\mathbb{R}^+} |\hat{f}(\xi)|^2 \left\{ \sum_{j \in \mathbb{Z}} \sum_{l \in \mathbb{N}} |\hat{\psi}(p^{-j}\xi) \overline{\hat{\psi}(p^{-j}\xi \oplus l)}| \right\} d\xi. \quad (3.9)$$

Taking infimum in (3.8) and supremum in (3.9), respectively, we obtain

$$A(\psi) \|f\|_2^2 \leq \sum_{j \in \mathbb{Z}} \sum_{k \in \mathbb{Z}^+} |f, \psi_{j,k}|^2 \leq B(\psi) \|f\|_2^2,$$

which holds for all $f \in \mathcal{D}$. This completes the proof of the theorem.

REFERENCES

- [1] Abdullah, Characterization of p -wavelets on positive half line using the Walsh–Fourier transform, *Int. J. Anal. Appl.*, 10 (2) (2016), 77-84.
- [2] Abdullah, Necessary condition of p -wavelet frame on positive half-line using the Walsh-Fourier transform, *Int. J. Comp. Math. Sci.*, 5(2) (2016), 105-109.
- [3] Abdullah, On the characterization of non-uniform wavelet sets on positive half line, *J. Info. Comp. Sc.*, 10 (1) (2015), 046-053.
- [4] Abdullah and Shah, F.A. Wave packet frames on local fields of positive characteristic. *Appl. Math. Comput.*, 249, 133-141 (2014).
- [5] Abdullah. Tight wave packet frames for $L^2(\mathbb{R})$ and $\mathcal{H}^2(\mathbb{R})$. *Arab J. Math. Sci.*, 19(2), 151–158 (2013).
- [6] P.G. Casazza and O. Christensen, Weyl-Heisenberg frames for subspaces of $L^2(\mathbb{R})$, *Proc. Amer. Math. Soc.*, 129 (2001), 145-154.
- [7] O. Christensen, Frames, Riesz and discrete Gabor/wavelet expansions, *Bull. Amer. Math. Soc.* 38 (2001), 273-291.
- [8] O. Christensen, *An introduction to Frames and Riesz Bases*, Birkhauser, Boston, 2003.
- [9] C.K. Chui and X. Shi, Inequalities of Littlewood-Paley type for frames and wavelets, *SIAM J. Math. Anal.*, 24 no. 1 (1993), 263-277.
- [10] I. Daubechies, A. Grossmann and Y. Meyer, Painless nonorthogonal expansions, *J. Math. Phys.*, 27 (1986), 1271-1283.
- [11] R. J. Duffin and A.C. Shaeffer, A class of nonharmonic Fourier series, *Trans. Amer. Math. Soc.* 72 (1952), 341-366.
- [12] Y. A. Farkov, Orthogonal p -wavelets on \mathbb{R}^+ , in *Proceedings of International Conference Wavelets and Splines*, St. Petersburg State University, St. Petersburg (2005), 4–26.
- [13] Y. A. Farkov, A. Y. Maksimov and S. A. Stroganov, On biorthogonal wavelets related to the Walsh functions, *Int. J. Wavelets, Multiresolut. Inf. Process.* 9(3) (2011), 485–499.
- [14] E. Hernández and G. Weiss, *A First Course on Wavelets*, CRC Press, New York, 1996.
- [15] F. A. Shah, Construction of wavelets packet on p -adic field, *Int. J. Wavelets, Multiresolut. Inf. Process.* 7(5) (2009), 553-565.
- [16] D. Li and H. Jiang, The necessary and sufficient conditions for wavelet frame on local fields, *J. Math. Anal. Appl.* 345 (2008), 500-510.
- [17] F. A. Shah, Tight wavelet frames generated by the Walsh polynomials, *Int. J. Wavelets, Multiresolut. Inf. Process.*, 11(6) (2013) 1350042, (15 pages).
- [18] F A Shah and Abdullah, A Characterization of Tight Wavelet Frames on Local Fields of positive characteristic, *J. Contem. Math. Anal.*, 49 (6) (2014), 251-259.
- [19] F A Shah and L. Debnath, p -Wavelet frame packets on a half-line using the Walsh–Fourier transform, *Integ. Trans. Special Funct.*, 22 (2011), 907–917.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)