



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 7 Issue: XI Month of publication: November 2019

DOI: <http://doi.org/10.22214/ijraset.2019.11107>

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

Review on Newton Raphson Method

Vishal Vaman Mehtre¹, Devender Kumar J Pal²

¹Assistant Professor, ²Department of Electrical Engineering, Bharti Vidyapeeth Deemed (to be) University, College Of Engineering, Pune

Abstract: The Newton's Raphson method is also known as Newton method. It is named after Isaac Newton and Joseph Raphson. This method is easy way to find an approximate to the roots of real value and also to solve the non-square and nonlinear problems. It also aims to represents a new approach of calculation of non-linear equation which is very similar to Newton Raphson method simple method and inverse Jacobian matrix will be used for further calculation and will also in some application. Self-derivative function in solving non-linear equation in scientific calculator, derivative Newton Raphson formula algorithm, uses and limitations of Newton Raphson method is been discussed below.

I. INTRODUCTION

Finding the solution of a set of no linear equation $f(x) = (f_1 \dots f_n) = 0$ is been a problem for past years. Newton Raphson method is to be considering an easy way to find the nonlinear equation solution. [2] Compare to other methods this method only requires one iteration and the derivative evaluation per iteration. Newton Raphson method is 7.678622465 times better than the bisection method and secant method is 1.389482397 times better Than Newton method[1]. This method is an easy way to find the roots of the nonlinear equation it provide good outcome with immediate convergence speed and matlab and C++ also adopted this method for finding the roots, scientific calculator is used for such calculation

II. NEWTONRAPHSON METHOD

Newton Raphson method is also known as Newton method, it is named after Issac Newton Joseph Raphson. This is a root finding algorithm which gives better approximations to the roots of real value[3]. It uses the idea that a continuous and differentiable function can be approximated by a straight line tangent to it. The idea is to start with a guess which is merely close to the true roots then a tangent line is drawn using calculus and where the it intersect the x axis. The point at which the x axis is intersected will be a better approximation to the original function root than the first guess and this method can be iterated.

Derivation of Newton method

X: $f(x) = 0$

Given a function f defined over the real x and its derivative f' , we begging with the first guess x_0 for root of the function f then x_1 is $X_1 = x_0 - f(x_0)/f'(x_0)$

IN graph tangent($x_0, f(x_0)$) intersect ($x_1, 0$) at x axis.

$X_{n+1} = f(X_n)/f'(X_n)$

Iteration is continued till the accurate value is reached. [6]




Fig 1.1 Graph illustrating root estimation using Newton Raphson method

1) Example

a) Find the real root of $2x^2 - 3\sin x - 6 = 0$ correct upto 4 decimal places with initial value of $x_0 = 1$ using Newton method

$$F(x) = 2x^2 - 3\sin x - 6$$

$$F'(x) = 4x - 3\cos x$$

1st Iteration

$$X_0 = 1$$

$$X_1 = x_0 - f(x_0)/f'(X_0)$$

$$F(x_0) = -5.424413$$

$$F'(x_0) = 2.279093$$

$$X_1 = 1 - (-5.424413)/2.279093$$

$$X_1 = 3.120668$$

2nd iteration

$$X_2 = 2.327631$$

3rd iteration

$$X_3 = 1.491784$$

4th iteration

$$X_4 = 1.950426$$

5th iteration

$$X_5 = 1.950157$$

6th iteration

$$X_6 = 1.950157$$

$$\text{Root of } f(x) = 1.950157$$

Table 1.1

Number	Iteration
X1	3.120668
X2	2.327631
X3	1.491784
X4	1.950426
X5	1.950157
X6	1.950157

b) Find the example, if one wishes to find the square root of 602, this is equivalent to finding the solution to [4]

$$X^2 = 602$$

The function to use in Newton's method is then,

$$F(x) = x^2 - 602$$

With derivative,

$$f'(x) = 2x$$

With the initial guess of 10, the sequence given by Newton's method is,

1st iteration

$$X_1 = x_0 - f(x_0)/f'(x_1)$$

$$= 10 - 10^2 - 602 / 2 * 10$$

$$= 35.6$$

2nd iteration

$$X_2 = x_1 - f(x_0)/f'(x_1)$$

$$= 35.6 - 35.6^2 - 602 / 2 * 35.6$$

$$X_2 = 26.38555671$$

3rd iteration

$$X_3 = 24.890065489$$

4th iteration



X4=24.738965784

5th iteration

X5= 24.73896584

Root of f(x) is 24.73896584

Table 2

Number	Iteration
X1	35.6
X2	26.385555671
X3	24.890065489
X4	24.738965784
X5	24.73896584

III. ADVANTAGES

- 1) This method is one of the quicker method which to iterate root fastly.
- 2) Converges of the root quadratic.
- 3) Approximation gets double when we get near to roots.
- 4) Multiple dimensions are easy to convert. [5]

IV. LIMITATION

- 1) This method may not work if there is point's inflection, local maxima or minima around x0 or the root.
- 2) Bad starting point
- 3) Iteration point is stationary
- 4) Derivative issue

V. CONCLUSION

Compare to others method Newton Raphson method is more accurate and fast. If the x axis and the tangent are merely parallel to each other this method is not useful in this situation. This method is most used in small molecules of calculation for large molecules of calculation this method is not used. This method is easy to convert multiple dimensions .It has a bad starting point and iteration is stationary. We can also say that the Newton Raphson method can be very useful to determine the intrinsic value based on measured permittivity.

VI. ACKNOWLEDGEMENT

We would like to show our gratitude & special thanks to Dr. D.S. Bankar, Head of Electrical department of electrical engineering for their support and for completing my research paper. I would also like my faculty members of the department of electrical engineering who helped us with extended support.

REFERENCES

- [1] Saba Akram, Qurrat ul Ann, "NewtonRaphson method", International Journal of Scientific & Engineering Research, Volume 6, Issue 7, July 2015
- [2] Ji Huan He, "A modified NewtonRaphson method", Volume 20, Issue 10,10 June 2004
- [3] Nicholas J Highman,& Hyunmin Kim "Numerical analysis for a quadratic matrix equation", Publication: 5 August 1999 from 13 December 1999
- [4] S.W.Ng &Y.S.Lee,"Variable Dimension NewtonRaphsonMethod",volume no 47, Issue no 6, June 2000
- [5] L.R.D.Reis, D.F.Novacki,"The NewtonRaphson method in the Extraction of Parameters of Pv modules", April 2017
- [6] Waqas Nazeer, Amir Naseem, "Generalized NewtonRaphson methods free from second derivative", Published in Journal of Nonlinear Science and Application, April 2016
- [7] Changbum Chun, "Iterative method improving Newton method by the decomposition method", March 2005.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)