



IJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 8 Issue: IV Month of publication: April 2020

DOI: <http://doi.org/10.22214/ijraset.2020.4129>

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

Roman Coloring of Bull Graphs and related Graphs

Preethi K Pillai¹, J. Suresh Kumar²

^{1,2}Assistant Professor, PG and Research Department of Mathematics, N.S.S. Hindu College, Changanacherry, Kerala, India
686102

Abstract: The Bull graph is a graph with 5 vertices and 5 edges consisting of a triangle with two disjoint pendant edges. In this paper, we obtain the Roman coloring of the Bull graph and related graphs such as Middle graph, Total graph, Splitting graph, Degree splitting graph, Shadow graph and Litact graph of the Bull graph.

Keywords: Graph, Roman coloring, Bull graph

I. INTRODUCTION

Graph coloring take a major stage in Graph Theory since the advent of the famous four color conjecture. Several variations of graph coloring were investigated [5] and still new types of coloring are available such as σ –(Sigma) coloring [4] and Roman coloring [6, 7]. In this Paper, we consider the proper coloring for the Bull graph and related graphs. By a graph, we mean a finite undirected graph without loops and parallel edges.

The Bull graph is a planar undirected graph with five vertices and five edges in the form of a triangle with two disjoint pendant edges. There are three variants of Bull graph, the name being derived from the fact that the structure of the graph is meant to represent the face of a “Bull”.




Figure.1. Bull Graph

The Bull graph was introduced by Weisstein [2]. It is a simple graph with chromatic number 3, chromatic index 3, radius 2, diameter 3 and girth 3. The concept of Bull-free graphs was studied, but there are no works related to the coloring of Bull graph and its related graphs. In this Paper, we investigate the coloring of Bull graph and its related graphs. We recall some basic definitions.

- 1) *Definition 1.1.* Roman coloring of a graph G is an assignment of three colors, namely $\{0, 1, 2\}$, to the vertices of G such that every vertex with the color, 0 must be adjacent to a vertex of color 2. The weight of a Roman coloring is defined as the sum of all vertex colors. The Roman Chromatic number of a graph G is defined as the minimum weight of a Roman coloring on G and denoted by $R(G)$.
- 2) *Definition. 1.2.* The Middle graph $M(G)$ of a graph G is the graph whose vertex set is $V(G) \cup E(G)$ and in which two vertices of $M(G)$ are adjacent if and only if either they are adjacent edges of G or one is a vertex of G and the other is an edge of G incident to it.




Figure.2. Middle graph of the Bull Graph

3) *Definition. 1.3.* The Total graph $T(G)$ of a graph G is the graph whose vertex set is $V(G) \cup E(G)$ and in which two vertices of $T(G)$ are adjacent whenever they are adjacent or incident in G .




Figure. 3. Total graph of the Bull graph

4) *Definition. 1.4.* The Splitting graph $S(G)$ of a graph G is obtained by adding a new vertex v' corresponding to each vertex v of G such that $N(v) = N(v')$ where $N(v)$ and $N(v')$ are the neighbourhood set of v and v' respectively.




Figure.4. Splitting graph of the Bull graph

5) *Definition 1.5.* Let $G = (V, E)$ be a graph with $V = S_1 \cup S_2 \cup \dots \cup S_t \cup T$, where each S_i is a set of vertices having at least two vertices and having the same degree and $T = V - \cup S_i$. The Degree Splitting graph $DS(G)$ of a graph G is obtained from G by adding vertices w_1, w_2, \dots, w_t and joining w_i to each vertex of S_i ($1 \leq i \leq t$)




Figure.5. Degree Splitting graph of Bull graph

- 6) *Definition.1.6.* The Shadow graph, $D_2(G)$ of a connected graph G is constructed by taking two copies of G , say G' and G'' and joining each vertex u' in G' to all the adjacent vertices of the corresponding vertex, u'' in G'' .




Figure.6. Shadow graph of Bull graph

- 7) *Definition.1.7.* Let $c(G)$ denotes the set of all cut vertices of Bull graph G . The Litact graph $m(G)$ of a graph G is the graph whose vertex set is $V(G) \cup c(G)$ in which two vertices are adjacent if they are adjacent edges of G or adjacent cut-vertices of G or one is an edge e_i of G and the other is a cut-vertex c_j of G and e_i is incident with c_j .




Figure.7. Litact graph of Bull graph

II. MAIN REESULTS

In this section, we discuss the Roman Coloring of Bull Graphs and related Graphs.

- 1) *Theorem. 2.1.* Roman Chromatic number of the Bull graph is 4.
 - a) *Proof.* Assign the color 0 to the two vertices of degree 3 each and the color 2 to the unique vertex of degree, 2 and assign the color 1 to the remaining 2 pendant vertices. Then it is a roman coloring with weight, 4. Clearly, it is minimal so that the Roman chromatic number of the Bull graph is 4.
- 2) *Theorem. 2.2.* Roman chromatic number of Middle graph of the Bull graph is 5.
 - a) *Proof.* In the Middle graph of Bull graph, let the vertices be v_i for $i = 1$ to 5 and e_j for $j = 1$ to 5. Assign the color 2 to the two vertices e_2 and e_4 and assign the color 0 to all the remaining tvertices except v_1 . We can assign the color 1 to v_1 . Then it is a roman coloring with weight, 5. Clearly, it is minimal so that the Roman chromatic number of the Middle graph of the Bull graph is 5.




Figure.8

3) *Theorem. 2.3.* The Roman chromatic number of the Total graph of the Bull graph is 4.

a) *Proof.* In the Total graph of Bull graph, let the vertices be v_i for $i = 1$ to 5 and e_j for $j = 1$ to 5. Assign the color 2 to the two vertices e_1 and v_4 and assign the color 0 to all the remaining vertices. Then it is a roman coloring with weight, 4. Clearly, it is minimal so that the Roman chromatic number of the Total graph of the Bull graph is 4.




Figure.9

4) *Theorem. 2.4.* The Roman chromatic number of Splitting graph of the Bull graph is 5.

a) *Proof.* Let the vertices of the Splitting graph of Bull graph be v_i for $i = 1$ to 5 and v'_j for $j = 1$ to 5. Assign the color 2 to the vertex v_4 , color 1 to the vertices v_1, v'_1, v'_4 and assign the color 0 to all the remaining vertices. Then it is a roman coloring with weight, 5. Clearly, it is minimal so that the Roman chromatic number of Splitting graph of the Bull graph is 5.




Figure.10

5) *Theorem. 2.5.* The Roman chromatic number of Degree Splitting graph of the Bull graph is 4.

a) *Proof.* Let the vertices of the Degree Splitting graph of Bull graph be v_i for $i = 1$ to 5 and w_j for $j = 1$ to 2. Assign the color 2 to the vertex v_4 , color 1 to the vertices v_1, w_1 and assign the color 0 to all the remaining vertices. Then it is a roman coloring with weight, 4. Clearly, it is minimal so that the Roman chromatic number of Degree Splitting graph of the Bull graph is 4.




Figure.11

6) *Theorem. 2.6.* The Roman chromatic number of shadow graph of the Bull graph is 5.

a) *Proof.* Let the vertices of the Shadow graph of Bull graph be v_i for $i = 1$ to 5 and v'_j for $j = 1$ to 5. Assign the color 2 to the vertex, v'_2 , color 1 to the vertices v_2, v_5, v'_5 and assign the color 0 to all the remaining vertices. Then it is a roman coloring with weight 5. Clearly, it is minimal so that the Roman chromatic number of shadow graph of the Bull graph is 5.




Figure.12

7) *Theorem. 2.7.* The Roman chromatic number of Litact graph of the Bull graph is 2.

a) *Proof.* Let the vertices of the Litact graph of Bull graph be e_i for $i = 1$ to 5 and c_j for $j = 1$ to 2. Assign the color 2 to the vertex, e_5 and assign the color 0 to all the remaining vertices. Then it is a roman coloring with weight 2. Clearly, it is minimal so that the Roman chromatic number of Litact graph of the Bull graph is 2.




Figure.13

REFERENCES

- [1] Gallian J.A, Dynamic Survey of Graph labelling, The Electronic Journal of Combinatorics, 2012.
- [2] Weisstein, Eric W , Bull Graph , Math World
- [3] Poulomy Ghosh, Sachchida N, Mishra, Anita pal, Various labelling on Bull graph and some related graphs, International ,Journal of Application of Fuzzy Sets and artificial intelligence (ISSN 2241-1240) vol:5 (2015) 23-35.
- [4] Gary Chartrand, Futaba Okamoto, Ping Zhang, The Sigma Chromatic Number of a Graph, Graphs and Combinatorics (2010) 26:755–773
- [5] Suresh Kumar J, Graph Coloring Parameters – A survey, International Journal for Research in applied Science and Engineering Technology, Vol. 7, Issue IV, April 2019.
- [6] Suresh Kumar J, Roman colouring of Graphs and Application to Military Strategy, International Journal for Research in Applied Science & Engineering Technology, Vol.8, Issue-III, March 2020.
- [7] Suresh Kumar J, and Preethi K Pillai, Proper Roman coloring of graphs, International Journal of Creative Research Thoughts, March, 2020
- [8] Suresh Kumar J, and Preethi K Pillai, Proper coloring of Bull graphs and related graphs, International Journal of Creative Research Thoughts, March, 2020 (Pre Print)
- [9] Suresh Kumar J, and Preethi K Pillai, Sigma coloring of Bull graph and related graphs, International Journal of Mathematics and its Applications, April, 2020 (Pre Print)
- [10] Harary Frank, Graph Theory, Addison Wesley, Reading Mass, 1969.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)