



# IJRASET

International Journal For Research in  
Applied Science and Engineering Technology



---

# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

---

**Volume: 9      Issue: IV      Month of publication: April 2021**

**DOI: <https://doi.org/10.22214/ijraset.2021.33677>**

**[www.ijraset.com](http://www.ijraset.com)**

**Call:  08813907089**

**E-mail ID: [ijraset@gmail.com](mailto:ijraset@gmail.com)**

# An Assessment of Earthworm Population Density of different Habitat Types in a Mango Cultivation Region of Palakkad District, Western Ghats

Chandini P K<sup>1</sup>, Dr. Jayasooryan K K<sup>2</sup>, Gayathri O S<sup>3</sup>, Sharon Mathew<sup>4</sup>, Dr. Rajathy Sivalingam<sup>5</sup>

<sup>2</sup>Scientist-B, Ecology and Environment Research Group, Center for Water Resource Development and Management, Kozhikode - 673 571

<sup>3,4</sup>Research Scholar, School of Environmental Studies, Cochin University of Science and Technology, Kochi, Kerala – 68 20 22

<sup>5</sup>Professor, School of Environmental Studies, Cochin University of Science and Technology, Kochi, Kerala – 68 20 22

**Abstract:** Earthworms are most important macrofauna in the soil, which plays a major role in soil fertility and nutrient cycling. The present study analysed population density of earthworms in four different habitats namely wooded grass land, irrigation canal, mango plantation with organic cultivation and mango plantation applying synthetic pesticides, in Muthalamada Panchayath, Palakkad District during June 2019 and June 2020. The study recorded five species of earthworms belonging to two families. Population density was varying in different habitats. From the study, it is observed that the intensive use agrochemicals have affected the earthworm diversity and population in the region.

**Keywords:** Earthworm, Population, Muthalamada, Mango Plantation, Pesticide

## I. INTRODUCTION

Earthworms are considered as an important soil macro fauna, having intense effect on ecosystem. Merely half a dozen species of earthworms in India are regularly used for vermiculture and composting (Julka and Paliwal, 2005). The Indian subcontinent has massive fauna of oligochaetes, which are represented by 509 species and 67 genera (Julka, 1993). Reproductive strategies of earthworms population helps to determine the genetic diversity of these creatures. The different spatial occurrence of earthworms roughly determine three ecological categories in the soil layer. Epigeic species also known as leaf litter dwellers, 2-6cm long, they decompose fresh organic matter near to the soil surface.

They are small and mostly dark red, mainly inhabits grassland, forest and compost, rarely found in cropland due to lack of permanent litter layers. They have a vigorous reproduction rate but a short lifespan. Endogeic species are shallow burrowers, small up to 18cm long, decompose organic substances in the soil and live in horizontal unstable galleries in the root area of the upper soil layer. Their reproduction rate is limited (8-12 cocoons/year) and lifespan is medium (3-5 years). Anecic species (deep burrowers, 15-45cm long) pull plant parts from the soil surface into their vertical stable burrows (diameter of 8-11mm), where they decompose partly and get ready for feeding. Because of their behavior, they are particularly sensitive to soil tillage at times, when they are active. Reproduction rate is limited and lifespan long. Adult animals deposit their faeces into the soil or above ground. Earthworms influence the supply of nutrients through their tissues but largely through their burrowing activities; produce aggregates and pores (i.e., bio structures) in the soil and/or on the soil surface, thus affecting its physical properties, nutrient cycling, and plant growth (Animesh *et al.*, 2014, Subin *et al.*, 2015, Bhaduarria *et al.*, 2000). Earthworms can indicate soil quality by (1) the abundance and species composition at a particular site, (2) the behavior of individuals in contact with a soil substrate (preference/avoidance/activity), (3) the accumulation of chemicals from the soil into the body, and (4) the biochemical/cytological stress-biomarkers in the earthworm (Ayten Karaca, 2011).

In an agricultural ecosystem, earthworms plays several beneficial roles for improving fertility of the soil and are called as cultivators of land. Their habit of burrowing and swallowing helps to increase the mobility and availability of nutrients in soil. Their burrow permits the penetration of air and moisture in the porous soil and improves the drainage; make the downward growth of the roots easier. The earthworms are continuously dragging dead leaves in to their burrows to eat them. They are only partially digested and their remains are thoroughly mixed with the castings. The earthworms consume the soil and organic matter and convert it into humus within short period of time and thereby increasing soil fertility. Reproduction and cocoon production is possible throughout the year.

The use of inorganic agriculture practices kill earthworms besides the target organisms. They are very susceptible to minimal dosages of agrochemicals, thus used as bio indicator of soil contamination providing an early warning of decline in soil quality. They serve as model organisms in toxicity testing. Earthworms are characterized by high ability to accumulate a lot of pollutants from soil in their tissues, thus they are used for studying of bioaccumulation potential of chemicals. A microcosm study conducted in orchards in the South Africa indicated adverse effects of spraying by pesticides (Chlorpyrifos and Azinphos methyl) on earthworm's biomass and cholinesterase activity. Authors concluded that earthworms were detrimentally affected by the pesticides due to chronic and intermittent exposure (Reinecke, 2007). A laboratory experiment that reproduced vineyard conditions in France showed that mixture of insecticides and/or fungicides at different environmental concentrations caused a neurotoxic effect in earthworms. After a long period of exposure or high concentrations, earthworms were physiologically damaged and could not cope with the high toxicity (Schreck *et al.*, 2008). The earth worm diversity and population density in Western Ghats was poorly studied especially in the intensive agricultural regions and mono culture plantations. The present study aims to estimate the population density of earthworms in different habitat types in and around the mango cultivating regions of Muthalamada, to understand the impacts of land use and cultivation practices on the diversity of earthworms.

## II. MATERIALS AND METHODS

### A. Study Area

Muthalamada is a village in Palakkad District in central Kerala, bordering with Tamil Nadu, is well known for mango plantation. Its geographical coordinates are 10° 38' 0" North, 76° 48' 0" East. The harvest season in Mango plantations in muthalamada begins by February end and goes on till the end of July. Muthalamada mango orchards well-known for the usage of extensive pesticides. The classification of study area were given in the table.1

### B. Earthworm Sampling and Preservation

The sampling location was located based on availability of worm caste on soil surface and humidity. Adult worms were collected by digging a pit of 25 X 25 cm to the depth of 30 cm and hand sorting method; the earthworms were sorted, washed with water and then preserved in 4% formalin (Julka, 1993, Julka, 1988). Collection was done during month of July 2019 and July 2020. The collected specimens were identified using systematic taxonomic procedures and keys with the help of experts.

Earthworm population density (EPD) was determined by the formula:

$$EPD = \text{Total no of earthworms in a sampling area} / \text{Sampling area (0.0625 m}^2\text{)}$$

## III. RESULT

In the present study, five different species of earthworms were identified which belonging to family Benhamiidae and Megascolicidae. During the study period (2019-2020) a total of 697 specimens were collected. The results are shown in Table 2 and 3. Among these specimens, 293 were identified with developed clitellum (mature worms). The study shows the presence five species namely *Dichogaster affinis* (Michaelsen, 1890), *Perionyx excavates* (Perrier, 1872), *Megascolex konkanensis* (Fedarb, 1897), *Metaphire houlleti* (Perrier, 1872) and *Megascolex sps*. The family Megascolex represents the highest species diversity with three Megascolex species followed. These results are comparable with the finding of Julka (2005). It is observed that earthworm population density is low in inorganic mango plantation, followed by grass covered portion with wooded area (chemmanampathi), water canal and organic mango plantation. Population density of *Perionyx excavates* was high in the study area and it was widely distributed in all the habitats. *Dichogaster affinis* is an exotic species which is widely distributed in the soils of mango plantation. The ecological categories of earthworms were given in the table 4.

## IV. DISCUSSION

Mariappan *et al.*, 2014 studied earthworm abundance in Rajapalayam cultivated lands, reported *Lampito mauritii* and *P. excavates* in paddy field, sugarcane farm, banana plantation, coconut grove and mango farm. They also assessed population density of earthworm and observed that lowest density in mango plantation while, coconut farm has highest density. Chaudhuri *et al.*, (2008) reported *Dichogaster affinis* in rubber plantations of Tripura. Julka, (1988) also reported *Dichogaster affinis* from Shasthankottah, Trivandrum (Thiruvananthapuram). *Metaphire houlleti* also recorded from Coconut plantation, manure heap, cocoa plantation, rubber plantations in Karnataka State (Siddaraju *et al.*, 2010). *Megascolex konkanensis*, *Metaphire houlleti*, *megascolex sp.* were recorded in the agro climatic regions of Wayanad District (John *et al.*, 2019). 10 different species of earthworms were obtained from different land areas like crop lands, agricultural lands, dense forests and river side's ecosystems in Wayanad which includes *Perionyx excavates* (Jijo *et al.*, 2019).



In the present study the maximum number of earthworms recorded were *Megascolex konkanensis* and *Perionyx excavates* from a water canal which is 2 km apart from the mango plantation. *Dichogaster affinis*, was evenly distributed all over the study area. Moreover these earthworms were distributed all over Kerala (Achuthan *et al.*, 2017, Harish kumar *et al.*, 2018, Prasanth narayanan *et al.*, 2014). *Eisenia fetida* (Savigny, 1826), *Pontoscolex corethrurus* (Müller, 1857), *Eudrilus eugeniae* (Kinberg, 1867), *Nematogonia panamaensis* (Eisen, 1900), *Metaphire houletti* (Perrier, 1872), *Polypheretima elongata* (Perrier, 1872), *Polypheretima taprobanae* (Beddard, 1892), *Ocnerodrilus occidentalis* (Eisen, 1878), *Pontodrilus litoralis* (Grube, 1855), *Dichogaster affinis* (Michaelsen, 1890), *Dichogaster annae* (Horst, 1893), *Dichogaster bolau* (Michaelsen, 1891), *Gordiodrilus elegans* (Beddard, 1892), and *Pithemera bicincta* (Perrier, 1875) are the exotic earthworms so far reported from Kerala state (Stephenson 1923, Aiyer 1929, Julka & Paliwal 1990, Kathireswari *et al.* 2005, Narayanan *et al.* 2012). *Metaphire houletti* (Perrier, 1872), *Dichogaster affinis* (Michaelsen, 1890) were recorded in the Muthalamada mango plantation and it recorded that native species was less compared to other habitats. The population density was high in mango plantation with organic cultivation practices during last samplings especially *Metaphire houletti* followed by *Dichogaster affinis* and *Megascolex konkanensis*. *Megascolex* sp. was also recorded in all habitats except mango plantation with inorganic cultivation. Sreelakshmi *et al.*, (2017), previously reported *Megascolex konkanensis* were recorded from Palakkad District. According to Julka (1993), family Megascolecidae, the distribution range of this family extends between Asia- and Australia. *Megascolex* and *perionyx* are confined to southern portion of Western Ghats. A few peregrine forms also introduced in soil in and around the roots of exotic plants which include Megascolecidae family. Successful colonization of peregrine species is mainly due to their tolerance to a wide range of ecological conditions. The reason for low population density in plantations is soil tillage and pesticide application. In water canals and mango plantation with organic cultivation practices the density was high which indicates better ecosystem services, such as decomposition of organic residues are carried out and the porosity is improved for infiltration. Endogeic species seem to adapt better to disturbances caused by ploughing and may benefit from soil inversion because of the incorporation of organic matter (Pelosi *et al.*, 2009). In arable soils, densities of epigeic species are low, depending on the amount of organic residues at the soil surface, since they are living in the mulch layer. They are better indicators of disturbances in soil. But in the present study total number of juvenile was 152, means a high reproduction rate and habitat suitability in canal area. The total earthworm density increases significantly with reduced tillage, due to the increased number of juveniles, while numbers of cocoons were many times higher with reduced tillage (Kuntz *et al.*, 2013). But in the present study no cocoons were observed in any of the sampling locations.

## V. CONCLUSION

From the study it was observed that organic cultivation farm followed by natural habitat have high population density and least in plantation with inorganic cultivation practices Earthworms are highly susceptible to insecticides causing immobility, rigidity and also show a significant effect on biomass reduction, growth and reproduction by disrupting various physiological activities leading to loss of earthworm population and soil biodiversity. So the application of chemical pesticides has significantly affected the population density of earth worms in Muthalamada mango plantations.

## VI. ACKNOWLEDGEMENT

First author acknowledges the research fellowship given by Government of Kerala.

## VII. CONFLICTS OF INTEREST

Authors declares no conflicts of interest

## REFERENCES

- [1] Achuthan Nair G., Vijayamohan TG., MaherHaeba H., Kathireswari Saminathan P. 2017. Diversity and Distribution of Earthworms in Relation to Altitude and Soil Factors of Kollam District, Kerala State, India, Environ. Sci. & Engg. Biodiversity and Conservation. 2:86-113
- [2] Animesh Dey., Chaudhuri P S. 2014. Earthworm community structure of pineapple (*Ananas comosus*) plantations under monoculture and mixed culture in West Tripura, India. Tropical Ecology. 55(1): 1-17
- [3] Aytan Karaca. 2010. Biology of Earthworms. Soil Biology. 2011th edition (9 November)
- [4] Bhadauria T., Ramakrishnan PS., Srivastava KN. 2000. Diversity and distribution of endemic and exotic earthworms in natural and regeneration ecosystems in central Himalaya, India. Soil Biology and Biochemistry. 32:2045-2054
- [5] Chaudhuri, PS., Nath S., Paliwal R. 2008. Earthworm population of rubber plantations (*Hevea brasiliensis*) in Tripura, India. Tropical Ecology. 49(2): 225-234
- [6] Fedarb SM. 1897. On some earthworms from India. Journal of the Bombay Natural History Society. 11(3): 431-437
- [7] Harish Kumar TS., Kanale S., Sreepada., Prasanth Narayanan S., John Reynolds W. 2018. Megadrile earthworms (annelida: oligochaeta) around udupi Power corporation limited (upcl), udupi district, karnataka, South-west coast of India. Megadrilogica. 23

[8] Jijo George., Deepthi MP., Saminathan KS and Kathireswari. 2019. Biodiversity and ecological category of earthworms in periyar of wayanad forest division, kerala. Life Science Journal.251-253

[9] Joseph John, Sreeraj PR., Asiya P., Kathireswari P., Rini Joseph and Jijo George.2019. Earthworm Population in Relation to Different Agro Climatic Regions of Wayanad District of Kerala. Mapana Journal of Sciences. 18: 4, 63-73

[10] Julka J M. 1988.The fauna of India and adjacent countries: Megadrile Oligochaeta (Earthworms). Haplotaxida: Lumbricina, Megascolecoidea: Octochaetidae, zoological survey of India. published by the director zoological survey of India, Calcutta. 1-415

[11] Julka JM., Paliwal R.1993. Collection, preservation and study of earthworms. Earthworm Resources and Vermiculture. 7-10.

[12] Julka J M., Paliwal R., and Kathireswari P.2009. Biodiversity of Indian earthworms-an overview. In: Edwards, C.A., Jayaraj, R., Indira,A(eds), Proceedings of indo US Workshop on Vermitechnology In Human Welfare. Rohini achagam, Coimbatore. 36-56

[13] Julka J,M.1993. Earthworm resources of India and their utilization in vermiculture. earthworm resources and vermiculture. 51-56

[14] Julka JM. 2014. Diversity and distribution of exotic earthworms (Annelida, Oligochaeta) in Indiaa review. In. Chaudhuri, P. & Singh, S.M. (Ed.) Biology and ecology of tropical earthworms,Discovery Publishing House Pvt. Ltd., New Delhi. 73-83.

[15] Kuntz M., Bernera A., Gattinger A., Scholberg JM., Mader P., Pfiffner L. 2013 Influence of reduced tillage on earthworm and microbial communities under organic arable farming. Pedobiologia. 56:251-260

[16] Mariappan V., Karthikairaj K., saiarasu L.2013. Relationship Between Earthworm Abundance and Soil Qualityof Different Cultivated Lands in Rajapalayam, Tamilnadu. World Applied Sciences Journal .27 (10): 1278-1281

[17] Muddaiah Siddaraju., Kanale S, Sreepada., John Warren Reynolds. 2010 checklist of earthworms (annelida: oligochaetes) from dakshina kannada, karnataka south west india. Megadrilogica. 14 (5):65-75

[18] Narayanan SP., Sathrumithra S., Christopher G., Thomas AP., Julka JM. 2016. Checklist of the earthworms (Oligochaeta) of Kerala, a constituent of Western Ghats biodiversity hotspot, India. Zootaxa. 4193 (1): 117-137.

[19] Pelosi C., Toutous L., Chiron F., Dubs, F., Hedde M., Muratet A., Ponge JF., Salmon S., Makowski D. 2013.Reduction of pesticide use can increase earthworm populations in wheat crops in a European temperate region Agriculture. Ecosystems and Environmen, 181:223-230

[20] Perrier E.1872. Recherches pour servir a l'histoire des Lombriciens terrestres. Nouv. Arch. Mus.Hist. Nat. 8: 5-198.

[21] Prasanth Narayanan S., Sathrumithra S., Dinu Kuriakose , Christopher G., Thomas AP. and Julka JM.2014. Earthworms (Oligochaeta: Megadrile) from the Mahatma Gandhi University campus, Kottayam, Kerala. Malabar Trogon Vol. 12:1-3

[22] Rashi Miglani.,Satpal Singh Bisht.2019. World of earthworms with pesticides and insecticides. Interdiscip Toxicol. 12: 71-82

[23] Reinecke SA., Reinecke AJ.2007. The impact of organophosphate pesticides in orchards on earthworms in the Western Cape, South Africa. Ecotoxicology and Environmental Safety. 66: 244-251

[24] Robert J., Blakemore A.2008. Series of Searchable Texts on Earthworm Biodiversity, Ecology and Systematics from Various Regions of the World- 3<sup>rd</sup> Edition. <http://www.annelida.net/earthworm/>

[25] Schreck E., Geret F., Gontier L., Treilhou M. 2008. Neurotoxic effect and metabolic responses induced by a mixture of six pesticides on the earthworm *Aporrectodea caliginosa nocturna*, Chemosphere. 71:1832-1839

[26] Sreelakshmi K., Sandra Rajan T., Danny Jacob P., Kathireswari P.2017. Earthworm Resources and Soil Parameters of Chittur, Palakkad and Kuvai, Coimbatore, India. International Journal of Current Microbiology and Applied Sciences. 6: 1860-1866

[27] Subin Kalu., Madan Koirala ., Udhab Raj., Khadaka.2015. Earthworm population in relation to different land use and soil characteristics. Journal of Ecology and the Natural Environment, 7:124-131

TABLES

Table 1. Sampling sites

Sampling site	Location	Characteristics
Grass covered portion with wooded area(S1)	10 <sup>0</sup> 35' 41.09'' 76 <sup>0</sup> 47' 36.20''	Standing crop field, zero tillage with grass cover
Irrigation canal(S2)	10 <sup>0</sup> 35' 29.43'' 76 <sup>0</sup> 45' 08.08''	Sandy soils was moist due to regular water supply for the fields
Organic Mango Plantation(S3)	10 <sup>0</sup> 35' 35.21'' 76 <sup>0</sup> 47' 36.20''	Mango plantation with weed cover but intensive management
Inorganic Mango Plantation(S4)	10 <sup>0</sup> 34' 37.77'' 76 <sup>0</sup> 45' 56.91''	Inorganic pesticide application with cultivation and intensive weed management

Table 2- Earthworm distribution in the soils of Muthalamada (2019)

study area	Species	Total number	EPD
S1	<i>Dichogaster affinis</i>	8	128
	<i>Perionyx excavatus</i>	33	528
	<i>Megascolex konkanensis</i>	60	960
	<i>Metaphire houlleti</i>	0	0
	<i>Megascolex sps.,</i>	0	0
S2	<i>Dichogaster affinis</i>	13	208
	<i>Perionyx excavatus</i>	65	1040
	<i>Megascolex konkanensis</i>	33	528
	<i>Metaphire houlleti</i>	0	0
	<i>Megascolex sps.,</i>	0	0
S3	<i>Dichogaster affinis</i>	3	48
	<i>Perionyx excavatus</i>	22	352
	<i>Megascolex konkanensis</i>	41	656
	<i>Metaphire houlleti</i>	0	0
	<i>Megascolex sps.</i>	4	64
S4	<i>Dichogaster affinis</i>	8	128
	<i>Perionyx excavatus</i>	0	0
	<i>Megascolex konkanensis</i>	4	64
	<i>Metaphire houlleti</i>	7	112
	<i>Megascolex sps.,</i>	0	0

Table -3 Earthworm distribution in different habitats of Muthalamada (2020)

study area	Species	Total number	EPD
S1	<i>Dichogaster affinis</i>	10	160
	<i>Perionyx excavatus</i>	40	640
	<i>Megascolex konkanensis</i>	70	1120
	<i>Metaphire houlleti</i>	0	0
	<i>Megascolex sps.,</i>	7	112
S2	<i>Dichogaster affinis</i>	15	240
	<i>Perionyx excavatus</i>	49	784
	<i>Megascolex konkanensis</i>	38	608
	<i>Metaphire houlleti</i>	0	0
	<i>Megascolex sps.,</i>	0	0
S3	<i>Dichogaster affinis</i>	3	48
	<i>Perionyx excavatus</i>	51	816
	<i>Megascolex konkanensis</i>	84	1344
	<i>Metaphire houlleti</i>	6	96
	<i>Megascolex sps.,</i>	0	0
S4	<i>Dichogaster affinis</i>	10	160
	<i>Perionyx excavatus</i>	0	0
	<i>Megascolex konkanensis</i>	6	96
	<i>Metaphire houlleti</i>	12	192
	<i>Megascolex sps.</i>	0	0

Table 4- Ecological categories of sampled earthworm

Sl No	Family	Species	Ecological Category	Native/Exotic	Habitat
1	Benhamiidae	<i>Dichogaster affinis</i>	Epegeic	Exotic	Grass areas
2	Megascolecidae	<i>Perionyx excavatus</i>	Epigeic	Exotic , Endemic peregrine	Water canal
3	Megascolecidae	<i>Megascolex konkanensis</i>	Endogeic	Native	Open ground areas
4	Megascolecidae	<i>Metaphire houlleti</i>	Epianecic	Exotic, Endemic peregrine	well wooded areas
5	Megascolecidae	<i>Megascolex sp.</i>	Epigeic	Native	Water canal





10.22214/IJRASET



45.98



IMPACT FACTOR:  
7.129



IMPACT FACTOR:  
7.429



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24\*7 Support on Whatsapp)