



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 4

Issue: 1

Month of publication: January 2016

DOI:

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

Phytochemical study of seeds of *Mucuna pruriens* (L.) DC.

Jyoti Pandey¹, Rajshree Pandey²

¹Research Scholar -Botany Deptt. ²Prof. of Botany,
Govt. Girls P.G. College, Rewa (M.P.), A.P.S. University, Rewa (M.P.) 486003

Abstract - The present paper deals to identify the phytochemical constituents and acute oral toxicity of methanolic extract of seeds of *Mucuna pruriens*. Fresh mature seeds were shade dried at room temperature, coarse powdered and extracted with methanol by Soxhlet's extraction method. Thereafter, the extract was concentrated using rotary flash evaporator to obtain semisolid crude extract with the yield of 09.534%. *Mucuna pruriens* seed extract was investigated for the presence of phytochemical constituents. The preliminary phytochemical evaluation of the *Mucuna pruriens* seed extract revealed the presence of steroids, alkaloids, tannins, carbohydrates, amino acid, resins and starch.

Keywords - *Mucuna pruriens*, phytochemical constituents.

I. INTRODUCTION

Nature always stands as a golden mark to exemplify the outstanding phenomenon of symbiosis. The biotic and abiotic elements of nature are all interdependent. The quest for long, healthy and happy life is as old as man himself. Nature has provided a complete storehouse of remedies to relieve the ailments of mankind. The consistent effects have resulted in many effective means of ensuring health care. The seers of Ayurveda were able to understand and record the various aspects regarding the drugs that even today are difficult to understand with modern available parameters (Anonymous, 1992 and Das, 1961). The medicinal plant products, which are derived from plant parts such as stem bark, leaves, fruits and seeds have been part of phytomedicine that produce a definite physiological action on the human body. The most important of these natural bioactive constituents of plants are alkaloids, tannins, flavonoids and phenolic compounds (Hill, 1952 & Gavishiddappa, *et al.* 2015). Itching bean *Mucuna pruriens* an underutilized legume species grown predominantly in Asia, Africa and in parts of America (Vadivel and Janardhanan, 2000).

Mature seeds, seeds from unripe pods and young pods of itching bean, *Mucuna pruriens* are soaked and boiled/roasted and eaten as such or mixed with salt by the North-East Indian tribes; North-Western parts of Madhya Pradesh tribes; South Indian tribes (Arora, 1991, Sahu, 1996 and Jain, 1981). To make this less- known legume palatable, tribal people follow a special processing method of continuous boiling and draining for about eight times until the boiled water changes from black to milky white. Consumption of improperly boiled seeds of itching bean is known to cause increase in body temperature and skin eruptions (Shankaranarayanan, 1978). It is attributed to the presence of high levels of 3, 4- dihydroxy-L-phenylalanine, L-Dopa, the aromatic non-protein amino acid (Jabadhas, 1980).

Hence, in the present study, the seeds of *Mucuna pruriens* were investigated and their chemical composition was investigated with a view to assess their phytochemical potential.

II. MATERIAL AND METHODS

A. Plant Materials

The seeds of *Mucuna pruriens* were procured locally after the seeds was authenticated by Dr. A.A. Khan, Retd. Prof. of Botany, Govt. Girls P.G. College, Rewa (M.P.). A voucher specimen has been deposited in our department.

B. Preparation Of Extract

Seeds were washed twice using tap water and then washed again in distilled water to remove the dust. The seeds were shade dried for 7–12 days at room temperature, until they were free from the moisture and then pulverized into coarse powder. The powdered material was extracted with methanol by Soxhlet's extraction method. Thereafter, the extract was concentrated using rotary flash evaporator to obtain semisolid crude extract. The percentage yield of the extract was found to be 09.534%. The extract was stored in airtight container in refrigerator below 10°C. Desired concentration of stock solution was prepared using distilled water for the following studies and then Preliminary phytochemical investigation were done.

C. Preliminary Phytochemical Screening

International Journal for Research in Applied Science & Engineering Technology (IJRASET)

Preliminary phytochemical tests were conducted on test extract to detect the presence of phytochemicals by following below mentioned the standard methods described in the Pharmacognosy text book of Trease and Evans.

D. Test for Steroids (Gibbs, 1974)

1) *Salkowski Test*: 2-3 drops of concentrated sulphuric acid was added to chloroform solution, shaken and allowed to stand, appearance of red color in lower layer indicates the presence of sterols.

2) *Liebermann-Burchard Test*: Extract was mixed with the chloroform and few drops of acetic anhydride and mixed well. Concentrated sulphuric acid was added from the sides of the test tube slowly until the ring appears; appearance of reddish brown ring indicates the presence of steroids.

E. Test for Flavonoids (Peach, 1956 and Rizk, 1982)

1) *Shinoda Test*: To the extract a few fragments of magnesium ribbon and concentrated hydrochloric acid was added. Appearance of red to pink color after few minutes indicates the presence of Flavonoids.

2) *Lead Acetate Test*: To the extract added few drops of aqueous basic lead acetate solution. Formation of yellow precipitate indicates presence of flavonoids.

3) *Alkaline Reagent Test/ NaOH Test*: few drops of sodium hydroxide solution was added to extract. Intense yellow color disappeared after adding dilute HCl which indicates the presence of flavonoids.

F. *Test For Alkaloids*: The extract was basified with ammonia and extracted with chloroform. The chloroform solution was acidified with dilute hydrochloric acid, shaken well and filtered. The filtrate was used for testing the alkaloids.

1) *Hager's Test*: The filtrate was treated with few drops of Hager's reagent. Formation of yellow precipitate indicates the presence of alkaloids (Varadarajan, *et al.* 2008).

2) *Wagner's Test (Iodine In Potassium Iodide)*: The acid layer was treated with few drops of Wagner's reagent. Formation of reddish brown precipitate indicates the presence of alkaloids.

3) *Mayer's Test (Potassium Mercuric Iodine Solution)*: The acid layer was treated with few drops of Mayer's reagent. Formation of creamy white precipitate indicates the presence of alkaloids.

4) *Dragendorff's Reagent (Potassium Bismuth Iodide)*: The acid layer was treated with few drops of Dragendorff's reagent. Formation of reddish brown precipitate indicates the presence of alkaloids.

G. Test For Tannins (Kokate, 1994)

1) *Gelatin Test*: To the extracts of the drug added 1% solution of gelatin containing 10% sodium chloride. Formation of white precipitate indicates the presence of tannins.

2) *Ferric Chloride Test*: To extracts few drops of 1% neutral ferric chloride solution were added, formation of blackish blue color indicates the presence of tannins.

H. Test For Saponins (Trease, 2002 And Sofowara, 1993)

1) *Foam Test*: Small amount of extract of the drug was shaken with little quantity of water, if foam produced persists for 10 minutes; it indicates the presence of saponins.

2) *Froth Test*: To 5 ml of extract of the drug added single drop of sodium bicarbonate solution. Shaken the mixture vigorously and left for 3 minutes. Formation of honey comb like froth indicates presence of saponins.

I. Test For Carbohydrates

Small amount of extracts of the drug were dissolved in little quantity of distilled water and filtered separately. The filtrates were used to test presence of carbohydrates.

1) *Molisch's Test*: The filtrate of the drug was treated with Molisch reagent and concentrated sulphuric acid was added from the sides of the test tube to form a layer. A reddish violet ring shows the presence of carbohydrates.

International Journal for Research in Applied Science & Engineering Technology (IJRASET)

2) *Benedict's Test*: to the filtrate added 2 ml Benedict's reagent and boiled in water bath. Formation of Green or reddish brown precipitate indicates presence of carbohydrates.

3) *Fehlings Test*: Filtrates were hydrolyzed with dilute hydrochloric acid, neutralized with alkali and heated with equal amount of Fehling's A and B solutions. Formation of green to yellow to red precipitate indicated the presence of reducing sugars.

J. Test For Amino Acid/ Protein

1) *Ninhydrin Test*: Heated the 3 ml of extract of the drug and 3 drops of ninhydrin solution in boiling water bath for 10 minutes. Appearance of purple color shows the presence of amino acids.

2) *Biuret Test*: To 3 ml of extract of the drug added 4% NaOH and few drops of 1% copper sulphate solution. Formation of violet color confirms the presence of protein.

3) *Millon's Reagent Test*: Mixed the extract with millon's reagent. Formation of brick red precipitate indicates the presence of protein.

K. Test For Resins

Dissolved the extract in acetone and pour the solution in to distilled water. Turbidity indicates the presence of resins (Harborne, 2007).

L. Test For Starch

Dissolved 0.015 gm of iodine and 0.075 gm of potassium iodide in 5 ml of distilled water and add 2-3 ml of an aqueous extract of drug, blue color is produced.

III. RESULTS

A. Preliminary Phytochemical Screening

Preliminary Phytochemical screening of methanolic extract of seeds of *Mucuna pruriens* revealed the presence of different kind of phytochemical components that are summarized in table 1

Table 1: Preliminary phytochemical screening of Methanolic *Mucuna pruriens* seed extract

S.No.	Phytochemical	Test	Result
1.	Test for Steroids	Salkowski test:	Present
		Liebermann-Burchard test	Present
2.	Test for Flavonoids	Shinoda test	Absent
		Lead acetate test	Absent
		Alkaline reagent test/ NaOH test	Absent
3.	Test for Alkaloids	Hager's test	Present
		Wagner's test	Present
		Mayer's test	Present
		Dragendorff's reagent	Present
4.	Test for Tannins	Gelatin test	Present
		Ferric chloride test	Present
5.	Test for Saponins	Foam test	Absent
		Froth test	Absent
6.	Test for Carbohydrates	Molisch's test	Present
		Benedict's test	Present
		Fehlings test	Present
7	Test for Amino acid/ Protein	Ninhydrin test	Present
		Biuret test	Present
		Millon's reagent test	Present

International Journal for Research in Applied Science & Engineering Technology (IJRASET)

8.	Test for Resins		Present
9.	Test for starch		Present

IV. DISCUSSION

These results obtained in the present study are in good consonance with the earlier reports of *Mucuna pruriens*. Steroids, alkaloids, tannins, carbohydrates, amino acid and resins were present in methanolic extract of seeds of *Mucuna pruriens*. The medicinal values of the seeds may be related to their constituent phytochemicals. According to Varadarajan *et al.*, (2008) the secondary metabolites (phytochemicals) and other chemical constituents of medicinal plants account for their medicinal value. For example, saponins are glycosides of both triterpene and steroids having hypotensive and cardiodepressant properties, while anthraquinones possess astringent, purgative, anti-inflammatory, moderate antitumor, and bactericidal effects (Olaleye, 2007 and Muzychkina, 1998). *Mucuna pruriens* indicates presence of both carbohydrates and starch this indicate they are polysaccharides. Presence of proteins in *Mucuna pruriens* has been described by Fathima *et al.* (2010) *Mucuna pruriens* contain higher crude protein when compared with commonly consumed pulse crops such as black gram, green gram, pigeon pea, chick pea and cow pea (Nagmain, *et al.* 2012).

V. CONCLUSION

In conclusion, the findings of the present study suggest that seeds extract of the *Mucuna pruriens* possesses steroids, alkaloids, tannins, carbohydrates, amino acid, resins and starch.

VI. ACKNOWLEDGEMENTS

The author is greatly indebted to Principal and Head of Botany Deptt. of Govt. Girls P.G. College, Rewa (M.P.) who permitted to carryout this work.

REFERENCES

- [1] Anonymous (1992). Charaka Samhita, Chakrapani Datta Teeka Chowkambha Prakashan, Varanasi.
- [2] Das S.R. Govind (1961). Bhaishajya Ratnavali. Chowkambha Sanskrit Series office, Varanasi.
- [3] Hill, A.F. (1952). Economic Botany. A textbook of useful plants and plant products. 2nd edn. New York: McGraw Hill Book Company Inc; P. 205
- [4] Vadivel, V., and Janardhanan, K. (2000). Nutritional and anti-nutritional composition of velvet bean: an underutilized food legume in South India. *Int. J. Food Sci.Nutr.*, 51: 279-287
- [5] Arora, K.R. (1991). Native food plants of the northeastern India. In: Contributions to Ethnobotany of India (Ed. SK Jain), Scientific Publishers, Jodhpur, and India. P.137-152.
- [6] Sahu, T.R. (1996). Life support promising food plants among aboriginal of Baster (M.P.), India. In: Ethnobiology in Human Welfare (Ed. SK. Jain), Deep Publications, New Delhi, India, P. 26-30.
- [7] Jain, S.K. (1981). Glimpses of Indian ethnobotany. New Delhi, India: Oxford and IBH Publishing Co. P. 1-230
- [8] Shankaranarayanan, A.S. (1978). Studies in the chemistry and pharmacology of Indian medicinal plants. Ph.D. Thesis, Madras University, Madras, India.
- [9] Jabadhas, A.W. (1980). Ethnobotanical studies on some hill tribes of South India. Ph.D. Thesis, Madras University, Madras, India.
- [10] Harborne, J.B. (2007). Phytochemical methods: A guide to modern techniques of plant analysis, 3rd edition, Chapman and Hall, London, 135-203.
- [11] Gibbs, R.D. (1974). Chemotaxonomy of Flowering Plants, Vol.1, McGill Queen's University Press, Montreal and London.
- [12] Peach, K. and Tracey, M.V. (1956). Modern methods of plant analysis, Vol 3, Springer Verlag, Berlin.
- [13] Rizk, A.M. (1982). *Fitoterapia*, 52:2: 35-42.
- [14] Trease, G.E. and Evans, W.C. (2002). Pharmacognosy, 15th edition, WB Saunders Publishers, London.
- [15] Kokate, C.K. (1994). Practical Pharmacognosy. 4th edition, Vallabh Prakashan, New Delhi, India.
- [16] Sofowara, A. (1993). Medicinal plants and traditional medicine in Africa. Chichester John Wiley & Sons, New York.
- [17] Veeraraghavan, Prema (2000). Expert consultant, CPCSEA, OECD Guideline no. 420.
- [18] Varadarajan, P., Rathinaswamy, G. and Asirvatham, D. (2008). "Antimicrobial properties and phytochemical constituents of Rheo discolor," *Ethnobotanical Leaflet*, 12: 841-845.
- [19] Olaleye, M.T. (2007). "Cytotoxicity and antibacterial activity of methanolic extract of Hibiscus sabdariffa," *Journal of Medicinal Plants Research*, 1:1: 9-13.
- [20] Muzychkina, R.A. (1998). Natural Anthraquinones: Biological and Physiological Properties, G. A. Tolstikov, Ed., PHASIS, Moscow, Russia.
- [21] Nagamani, J., Suresh, J., Ahuja, V. Reddy (2012). Comparative phytochemical screening of Vatashunga, Shatavari and Shatapushpa claimed for Prajasthapana activity; *Annals of Biological Research*, 3:3:1294-1304
- [22] Fathima, K.R., P. Tresina Soris, and V.R. Mohan (2010). Nutritional and Antinutritional Assessment of *Mucuna pruriens* Adv. Biores. vol 1:2:79-83.
- [23] Gavishiddappa, A. Hadimani, S.D. Desai, Prakash, Biradar, Nanjappaiah, H.M., Shivakumar Hugar, Ishwar, B. Bgoji (2015). Evaluation of Acute Oral Toxicity and Phytoconstituents of Methanolic Extract of *Mucuna pruriens*. *Phcog common*, Vol 5, Issues.1.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)