



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 4 Issue: VII Month of publication: July 2016

DOI:

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

Solar and UV Photolysis of Ampicillin Antibiotic Solution

Gayatri Devi Singh¹, Dr. K. C. Gupta², Dr. Archana Shrivastawa³

²Department of Microbiology, VRG Girls College, Morar Gwalior, (M.P.) India

^{1,3}Department of Microbiology, College of Life Science, Cancer Hospital and Research Institute Gwalior, (M.P.) India

Abstract-The ampicillin shows broad spectrum antibacterial activity mostly against Gram positive bacterial group but it also shows such activities against some Gram negative bacterial group. The solution of concentration ranging from 50-500 ppm of the ampicillin was exposed to the sun light and UV radiation. The duration of the exposure, for the degradation by UV radiation, was 1 to 10 hrs. The UV rays (15W and 30W) lamps were used for the study. The degradation was measured by the zone of inhibition. The maximum degradation was recorded after 5 hrs. of duration by the exposure of UV 30W. The maximum degradation of the solar degradation was after 72 hrs. of the direct sun light exposure. The UV and the solar degradation proved to be the significant mode to control the environmental pollution caused by ampicillin residues.

Key words- Photolysis, Ampicillin, UV radiation and Environment.

I. INTRODUCTION

Antibiotics are molecules that kill or inhibit the growth of microorganisms, including bacteria and fungi both (Cundcliffe et. al., 2000; Dunkle et. al., 2010). Ampicillin is a semi-synthetic antibiotic belonging to the penicillin group, which consist β -lactam ring, thus it belongs to β -lactam group of antibiotics (Nascimento et. al., 2013). It shows broad spectrum antibacterial activity against Gram positive bacterial and some Gram negative bacterial group. This antibiotic is used for the treatment of meningitis, enterococcal endocarditis urinary tract infection and prophylaxis for bacterial endocarditis (Sharma et al 2013). The antibiotics as human and veterinary medicine, has led to wide spreaded toxic pollution in the environment. The biological resistant antibiotics have been recorded in pre-treated sewage, industrial effluent, aquatic environment and also in drinking water (Klavarioti et al., 2009). Emolla et al., (2009) reported the photo-degradation of ampicillin, amoxicillin, and cloxacillin antibiotics with the help of photo-fenton process in aquatic medium. Elmolla et al., (2010) focussed on the aqueous degradation of amoxicillin, ampicillin and cloxacillin antibiotics in different operating condition such as UV, zinc oxide, concentration, pH and irradiation time and observed the rate of degradation of amoxicillin, ampicillin and cloxacillin antibiotics was highly affected by the pH. Wirzal et al., (2013) recorded that the degradation of ampicillin penicillin-G by oxidation method by using metal (mixed) oxide electrode as anode. Alalm (2016) recorded the photo-degradation of ampicillin antibiotic with the help of UV radiations in the presence of Ru, WO₃ and ZrO₂ catalyst in liquid medium. The main objective of this study was to determine the maximum rate the degradation of ampicillin antibiotic by sun light and UV radiation of different intensities.

II. MATERIAL AND METHOD

A. Chemicals and Media

Ampicillin 500mg (manufactured by Candila pharmaceuticals Ltd. Dholka, Ahmedabad, India) was used in the present study. The Muller-Hinton agar media (High media) was used for the determination of the zone of inhibition.

B. Experimental Setup


1) *Effects of Sun Light on Antibiotics:* Different concentrations of ampicillin antibiotic solutions (50, 100, 200, 300, 400 and 500 ppm) were prepared and poured in the petriplates for experiments. These plates were exposed to sun light during day time in the month of March in the Gwalior region (Madhya Pradesh, India) the temperature was 30^oC, the sun light exposure was period ranging from 12-108 hrs. was observed. The antibiotic discs were prepared by dipping them in the different concentrations of antibiotic, which were then exposed to the sun light. The controls discs dipped in untreated antibiotic solutions were also prepared. The bacterial culture was prepared in the petriplates containing the Muller-Hinton agar media. The treated and control discs were placed in these petriplates containing media. The petriplates, containing treated and control discs, were exposed to the environment of the institution for bacterial culture for five minutes. After exposure these plates were incubated at 37^oC for 48 hrs. The inhibition zones, appeared in the petriplates, were measured in millimeters (Jasim et al., 2010; Chen et al., 2011;

International Journal for Research in Applied Science & Engineering Technology (IJRASET)

Hernandez et al., 2012).

- 2) Effects of UV Radiation on Antibiotics: Different concentrations of ofloxacin antibiotic solutions (50, 100, 200, 300, 400 and 500 ppm) were prepared for experiments as discuss above. These plates were exposed to UV radiation in laboratory at 30⁰C temperature for the period ranging from 1to 10 hrs. The UV-lamps (Philips, 15W and 30W) were used as source for UV radiation. The antibiotic discs were prepared by dipping in the different concentrations of antibiotic, which were exposed to the UV radiation. The controls discs containing untreated antibiotic solutions were used as controls. The petriplates containing the Muller-Hinton agar media were also prepared for the bacterial culture. The treated and control discs were placed in these petriplates. These petriplates were exposed to the environment of the institution for bacterial culture for five minutes. These plates were incubated at 37⁰C for 48 hrs. The inhibition zones appeared in the petriplates were measured in millimeters (Jasim et al., 2010; Chen et al., 2011; Hernandez et al., 2012).

Structure of Ampicillin



Chemical fomula: C₁₆H₁₉N₃O₄S

Molecular weight: 349.41g/mol

IUPAC name: (2*S*,5*R*,6*R*)-6-([(2*R*)-2-amino-2 phenylacetyl] amino)-3,3-dimethyl-

7-oxo-4 thia-1 azabicyclo [3.2.0] heptane-2- carboxylic acid.

III. RESULT AND DISCUSSION

Emolla et al., (2009) have demonstrated the complete degradation of ampicillin, cloxacillin and amoxicillin at the concentration 105, 103 and 104 mg/l respectively after 5hrs. UV exposure and at pH 5. The percentage of ampicillin, cloxacillin and amoxicillin degradation was 3.8%, 4.9% and 2.9% respectively. Emolla et al. (2010) also studied the degradation of ampicillin, cloxacillin and amoxicillin at the concentration 105, 103 and 104 mg/l respectively but when the antibiotics were treated with UV/ZnO photocatalysis they recorded that the complete degradation took place after 180 minute of exposure. Dehghani et al., (2014) have recorded that the UV radiations were most effective for the penicillin G degradation. They also noticed the photo-catalytic degradation rate of penicillin G was increased by UV radiation treatment.

In the present study the degradation efficiency of ampicillin was measured by the UV exposure of two intensities (15W and 30W). The maximum degradation of AMP was recorded at 30W of UV exposure after 5 hours but 15W exposure of UV indicated the maximum degradation was possible after 7hrs of exposure (Table 1, 2, fig. 1, 2, 3, 4, 5 and 6).The different concentrations (50 to 500ppm) of ampicillin antibiotic were degraded by the solar exposure at different time intervals (12hrs to 108hrs). The maximum degradation of ampicillin was observed after 72 hrs. (Table3 and fig.7).

International Journal for Research in Applied Science & Engineering Technology (IJRASET)

Table-1:- Zone of inhibition (in mm) of the Ampicillin with UV (15W) treatment.

Con. → Exposure Time ↓	Concentration					
	50ppm	100ppm	200ppm	300ppm	400ppm	500ppm
0hrs	9.0	11	13	15	18	20
1hrs	8.0	9.0	11	13	16	18
2hrs	6.0	8.0	10	12	14	16
3hrs	5.0	7.0	9.0	10	13	15
4hrs	4.0	5.0	7.0	9.0	11	13
5hrs	3.6	4.0	6.0	8.0	8.0	9.0
6hrs	2.9	3.0	4.5	6.0	7.0	7.6
7hrs	2.7	2.9	4.0	5.5	6.4	6.9
8hrs	2.6	2.8	3.9	5.4	6.3	6.8
9hrs	2.5	2.7	3.8	5.3	6.2	6.7
10hrs	2.4	2.6	3.7	5.2	6.1	6.6

Table-2:- Zone of inhibition (in mm) of the Ampicillin with UV (30W) treatment.

Con. → Exposure Time ↓	Concentration					
	50ppm	100ppm	200ppm	300ppm	400ppm	500ppm
0hrs	9.0	11	13	15	18	20
1hrs	7.0	8.0	10	11	14	16
2hrs	5.0	7.0	9.0	10	12	14
3hrs	4.0	5.0	7.0	9.0	10	11
4hrs	3.0	3.0	6.0	8.0	9.0	9.5
5hrs	2.2	2.3	4.0	5.0	6.5	7.0
6hrs	2.1	2.2	3.9	4.9	6.4	6.9
7hrs	2.0	2.1	3.8	4.8	6.3	6.8
8hrs	1.9	2.0	3.7	4.7	6.2	6.7
9hrs	1.8	1.9	3.6	4.6	6.1	6.6
10hrs	1.7	1.8	3.5	4.5	6.0	6.5

International Journal for Research in Applied Science & Engineering Technology (IJRASET)

Table-3:- Zone of inhibition (in mm) of the Ampicillin with sun light treatment.

Con. →	50ppm	100ppm	200ppm	300ppm	400ppm	500ppm
Exposure Time ↓						
Non treated	9	11	13	15	18	20
12hrs	8	10	12	14	17	19
24hrs	7	9	11	13	15	17
36hrs	6	7	10	11	13	16
48hrs	5	6	8	10	11	14
60hrs	4	5	7	9	8	10
72hrs	3	3.8	4	5.0	7.0	8
84hrs	2.9	3.7	3.9	4.9	6.9	7.8
108hrs	2.8	3.6	3.8	4.8	6.8	7.6




Fig 1: 50 ppm concentration of ampicillin showing UV degradation.

International Journal for Research in Applied Science & Engineering Technology (IJRASET)




Fig2: 100ppm concentration of ampicillin showing UV degradation.




Fig 3: 200ppm concentration of ampicillin showing UV degradation.

International Journal for Research in Applied Science & Engineering Technology (IJRASET)




Fig4: 300ppm concentration of ampicillin showing UV degradation.




Fig 5: 400ppm concentration of ampicillin showing UV degradation.

International Journal for Research in Applied Science & Engineering Technology (IJRASET)




Fig 6: 500ppm concentration of ampicillin showing UV degradation.




Fig 7: Ampicillin showing degradation with sun light treatment.

IV. CONCLUSION

The semi synthetic antibiotic ampicillin is affective against the harmful bacteria but it is also a killing agent of useful bacteria. When this antibiotic is released untreated in the soil through nursing home and hospital garbage then it causes the soil as well as water

International Journal for Research in Applied Science & Engineering Technology (IJRASET)

pollution. For the remediation of such pollutant, the solar and UV exposure were studied. The UV exposure was more effective than the solar exposure. The optimum degradation was observed after 5 hrs. of 30 W UV exposure. The maximum solar degradation of ampicillin was noticed after 72 hrs. of exposure.

REFERENCES

- [1] Alalm, M.G., Ookawara, S., Fukushi, D., Sato, A., and Tawfik, A., (2016). Improved WO₃ photocatalytic efficiency using ZrO₂ and Ru for the degradation of carbofuran and ampicillin. *Journal of Hazardous Materials*. 302: 225-231.
- [2] Chen, Dong-hui, Wu, Yang, Kookana, Rai, (2011). Aqueous Photodegradation of Selected Antibiotics under Different Conditions. *International Conference on Biology, Environ. Chem.*, 1: 191- 194.
- [3] Cundcliffe, E., (2000) Antibiotic biosynthesis: Some thoughts on “why” and “how”. *The Ribosome: Structure, Function, Antibiotics and Cellular Interactions*, eds RA Garrett, A Liljas, AT Matheson, PB Moore, and HF Noller (ASM Press, Washington, DC), pp 409–417.
- [4] Dehghani, M., Nasser, S., Ahmadi, M., Samaei, M.R., and Anushiravani, A., (2014). Removal of penicillin G from aqueous phase by Fe³⁺- TiO₂/UV -A process. *Journal of Environmental Health Science and Engineering*. 12:56.
- [5] Dunkle A. Jack, Xiong Liqun, Mankin S. Alexander, and Cate H.D. Jamie (2010). Structures of the Escherichia coli ribosome with antibiotics bound near the peptidyl transferase center explain spectra of drug action. *PNAS Early Edition* pp. 17152-17157.
- [6] Elmolla, E.S., and Chaudhuri, M., (2010). Degradation of amoxicillin, ampicillin and cloxacillin antibiotics in aqueous solution by the UV/ZnO photocatalytic process. *Journal of Hazardous Materials*. 173(1-3): 445-449.
- [7] Elmolla, E.S., and Chaudhuri, M., (2009). Degradation of the antibiotics amoxicillin, ampicillin and cloxacillin in aqueous solution by the photo-fenton process. *Journal of Hazardous Materials*. 172: 1476-1481.
- [8] Hernandez, F., Rivera, A., Ojeda, A., Zayas, T., and Cedillo, L., (2012). Photochemical Degradation of the Ciprofloxacin Antibiotic and Its Microbiological Validation. *Journal of Environmental Science and Engineering A*.1:448-453.
- [9] Jasim, H. A., Ali, S., and Abdulla, N., madher, (2010). Evaluation of Antibacterial Activity of Tetracycline and Cephalexine Decomposed by Sunlight. *Journal of Basrah Researches (Sciences)*.36(5):73-78. *Journal of Environmental Science and Engineering A*.1:621-649.
- [10] Klavarioti, M., Mantzavinos, D., Kassianos, D., Envi .Inter. 35 (2009) 402.
- [11] Nascimento, T.G., Oliveira, E.J., Junior, I.D.B., Araujo-Junior, J.X., and Macedo, R.O., (2013). Short term stability studies of ampicillin and cephalixin in aqueous solution and human plasma: Application of least squares method in arrhenius equation. *Journal of Pharmaceutical and Biomedical Analysis*. 73: 59-64.
- [12] Navarro, P.G., Blazquez, I.H., Osso, B.Q., Martinnez de las Parras, P.J., Puentedura, M.I.M., and Garcia, A.A.M., (2003). Penicillin degradation catalyzed by Zn(II) ions in methanol. *International Journal of Biological Macromolecules*. 33: 159-166.
- [13] Sharma, S. K., Singh, L., and Singh, S., (2013). Comparative study between penicillin and ampicillin. *Scholars Journal of Applied Medical Sciences (SJAMS)*. 1(4): 291-294.
- [14] Wirzal, M.D.H., Yusoff, A.R.M., Zima, J., and Barek, J., (2013). Degradation of ampicillin and penicillin G using anodic oxidation. *International Journal of Electrochemical Science*. 8: 8978-8988.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)