



# IJRASET

International Journal For Research in  
Applied Science and Engineering Technology



---

# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

---

**Volume: 5      Issue: VI      Month of publication: June 2017**

**DOI:**

**[www.ijraset.com](http://www.ijraset.com)**

**Call:  08813907089**

**E-mail ID: [ijraset@gmail.com](mailto:ijraset@gmail.com)**

# **Utilization of Controlled Wastewater Nutrients for Lipid Production from *Botryococcus Braunii***

Dr. A. Swaroopa Rani<sup>1</sup>

<sup>1</sup>Dept of Biotechnology, Jawaharlal Nehru Technological University Anantapur, College of Engineering, Pulivendula- A.P

**Abstract -** Diminishing oil reserves, rising oil prices and a significant increase in atmospheric carbon dioxide levels have led to an increasing demand for alternative fuels. Microalgae have been suggested as a means for fuels production because of their advantages related to higher growth rates, higher photosynthetic efficiency and higher biomass production, compared to other terrestrial energy crops. During photosynthesis, microalgae can fix carbon dioxide from different sources, including the atmosphere, industrial exhaust gases and soluble carbonate salts. To determine the most optimal conduction for the growth of *Botryococcus braunii* in order in order to produce lipids that can be transformed into biodiesel fuel, different nutritional conduction were investigated. For this purpose sewage water is used as a media; and effect of nutrients viz., carbon (C), nitrogen (N), phosphorus (P) and potassium (K) etc.. on biomass growth and lipid accumulation was elucidated and the results achieved are most effective and further analyzed for biomass production chlorophyll content and lipid content. The best growth resulted in an enriched solution from the N+P media condition and the best results were obtained when medium was maintained at 16 hrs light/8 hrs darkness cycles and mechanical stirring at 120 rpm (only during the lighting hrs) with higher lipid content of 70% was observed with (C) condition.

**Keywords--** Biofuels, microalgae, *Botryococcus braunii*, sewage water, batch culture.

## **I. INTRODUCTION**

The colonial alga *Botryococcus braunii* is widespread in freshwater environments and can be also found in brackish lakes, reservoirs, ponds in subtropical or tropical zones [1-7]. This alga is famous for a notable ability to synthesize a variety of lipids. It produces a high amount of hydrocarbons, excretes them outside the cells, and accumulates them in its extracellular matrix [8, 9]. Some strains of *Botryococcus braunii* can also produce several certain ether lipids [6, 10]. The hydrocarbon content in *Botryococcus braunii* is significantly higher than that in other oil-producing algae and it has already been proposed as a future renewable source of fuel

Biotechnological value of *Botryococcus braunii*: It is known that hydrocarbons from *Botryococcus braunii* can be used as direct substitute of petroleum-based fuel products without the complexity of tranesterification and low temperature gelling, which showed an inherent superiority of producing hydrocarbons over biodiesel. *Botryococcus braunii* has a thicker cell wall than most strains, and usually the wet biomass contains a lot of water which cause the efficiency of hydrocarbon recovery to be rather low. Enzymes regulates the unique pathways of triterpene biosynthesis in *Botryococcus braunii* have attracted many interests.

Biodiesel fuel is becoming more promising as it is produced from non toxic, biodegradable and renewable resources and its use leads to a decrease in the emission of harmful air pollutants [11].

Microalgae are a group of fast growing unicellular or simple multicellular micro organisms, which have the ability to fix CO<sub>2</sub> while capturing solar energy with efficiency 10 to 50 times greater than that of terrestrial plants and higher biomass production compared to energy crops [12]. Microalgae have several advantages, including higher photosynthetic efficiency as well as higher growth rates and higher biomass production compared to other energy crops. Several microalgae strains have been reported to have the ability to accumulate large quantities of lipids.

Microalgae *Botryococcus braunii* was chosen as a subject for this research due to its easy growth in a sewage water as a media without the necessity of utilizing very specific compounds and its significant lipid content, as lipids are the most desirable component from the energy point of view.

The purpose of this work was to establish the growth of *B. braunii* in a medium where the nutrients and the other organic compounds existing in the wastewater can serve as a source of nutrient medium for the growth of algae. The process also provides economic mode of lipid production from wastewater. Therefore an attempt was made in this study to evaluate the influence of sewage waste water in association with nutrients supplementation on growth and lipid accumulation using *Botryococcus braunii* and

## International Journal for Research in Applied Science & Engineering Technology (IJRASET)

sewage wastewater as substrate. Influence of nutrient conditions was studied individually and in combinations. Biomass growths were estimated while lipids were extracted and analyzed. An investigation was based on batch experiments carried out in a lab in order to optimize the medium composition for the best microalgal biomass and lipid yield, and consequently for the economic efficiency of the process if used in an up-scale for biofuel production.

### II. MATERIALS AND METHODS

#### A. Microalgae strain and medium

*Botryococcus braunii* was isolated from mixed microalgae culture. *B. braunii* was grown on CHU media.

#### B. Cultivation

The cells in exponential period were inoculated (10%, v/v) in a sewage culture medium (sewage water was autoclaved at 121<sup>0</sup> C for 20 min and then twice filtered through whatt man filter paper) to start the culture. In addition to this different nutrients viz., nitrogen (N), phosphorus (P), carbon (C) and NPK. Nine experimental variations were studied by varying the nutrients as described in Table 1. A control condition without the addition of external nutrients was operated parallelly to elucidate the effect of native nutrients present in the sewage water.

All the experiments were carried out at a photo period of 16 h light:08 h dark. Light intensity during the experiment was measured using a light meter (Li- Cor measuring device). In the light phase, flasks were placed in a rotary shaker at 120 rpm for mixing. The cultures were maintained at room temperature (25-27°C) at pH: 8.2 on a fluorescent light with a light dark photoperiod of 16h: 8h. Sterile-air containing 2% (v/v) CO<sub>2</sub> was aerated into the flask through an air sparger at the bottom of the flask. The strains were checked for 25 days growth period. All experiments were conducted in duplicates.

#### C. Growth measurements

The growth of *B. braunii* was measured via spectrophotometry and biomass dry weight. Optical density for biomass factor was determined at wavelength 550 nm. One ml of sample was appropriately diluted with deionized water and the absorbance of the sample was read at 550 nm.

The cultures were determined gravimetrically and growth was expressed in terms of dry weight (mg/L) [13]. The cultures were harvested by centrifugation at 3000g for 10 min and the cells were washed with distilled water. Then the pellet was freeze dried. The dry weight of algal biomass was determined gravimetrically and growth was expressed in terms of dry weight (g/l).

The biomass yield was calculated. Biomass yield (mg ml<sup>-1</sup>) = 
$$\frac{\text{Final weight (g)} - \text{Initial weight (g)}}{\text{Sample taken (ml)}}$$

#### D. Chlorophyll and carotenoid content

The isolation of pigments from algae cells included the following procedures: harvesting 2 ml of microalgae cells by centrifugation at 10000 rpm, two times for 3 min and discarding the supernatant, suspension of cells in 2 ml methanol/water 90:10 v/v and mixing of Vortex for 1 min., heating of the suspension for half an hour in a water bath at 60°C, cooling of the samples at room temperature, centrifugating the suspension (10000 rpm for 3 min) and discarding the supernatant with dissolved pigments. The absorbance of the pigments extract (665, 652 nm for chlorophyll content (a+b) and 470, 666 nm for carotenoids content) was recorded by using spectrophotometer. The chlorophyll content was computed (mg/l) according Porra et al. [14] and carotenoid content was computed (mg/l) according Lichtenthaler [15].

Chlorophyll (a) and chlorophyll (b) were estimated as given below.

Chlorophyll a (µg ml<sup>-1</sup>) = 16.75 x A<sub>665.2</sub> - 9.16 x A<sub>652.2</sub>

Chlorophyll b (µg ml<sup>-1</sup>) = 34.09 x A<sub>652.4</sub> - 15.28 x A<sub>665.2</sub>

Total Chlorophyll (a+b) content was calculated.

#### E. Lipid content

The total lipids were extracted from microalgae biomass using a modified method of Bligh & Dyer, 1959 [16]. The lipids were extracted using a mixture of chloroform/methanol (1:2 v/v). The quantity of lipid residue was measured gravimetrically and expressed as dry weight percentage.

### III. RESULTS AND DISCUSSION

#### A. Culture of *Botryococcus braunii*

## International Journal for Research in Applied Science & Engineering Technology (IJRASET)

Much research has been carried out on investigating the optimal culture conditions for *Botryococcus braunii*. It is recognized that the production of hydrocarbons in *Botryococcus braunii* is positively correlated to the cell growth, regardless of the particular culture conditions and the media used. According to the algal biochemistry, it is such a substantial commitment for *Botryococcus braunii* to produce a considerable amount of hydrocarbons, the energetically expensive molecules. Therefore, the observed slow growth rate of this microalgae compared to other strains is apparently not the consequence of nutrient limitation, but the energetic consumption to synthesize and accumulate hydrocarbons.

*Botryococcus braunii* culture requires water, light, CO<sub>2</sub>, and inorganic nutrients. Factors that govern the algae productivity include pH, pCO<sub>2</sub>, light intensity, salinity and temperature. Modified Chu-13 medium is often used as growth medium for *Botryococcus braunii*. This medium has the following composition (g·L<sup>-3</sup>): KNO<sub>3</sub> (0.05), K<sub>2</sub>HPO<sub>4</sub> (0.01), MgSO<sub>4</sub>·7H<sub>2</sub>O (0.025), CaCl<sub>2</sub>·6H<sub>2</sub>O (0.02), ferric citrate (0.0025), citric acid (0.025), boron (0.125 ppm), manganese (0.125 ppm), copper (0.005 ppm), cobalt (0.005 ppm), and molybdenum (0.005 ppm). The pH in the media is adjusted to 7.5 before sterilization [17].

CO<sub>2</sub> is necessary for Photosynthetic cultures of *Botryococcus braunii*. CO<sub>2</sub> enrichment favors the formation of lighter botryococcenes (C30–C32), while a substantial portion of heavier botryococcenes (C33–C34) are discovered in the cells sparged with ambient air [18].

Although under autotrophic cultivation, *Botryococcus braunii* could also utilize exogenous carbon sources for enhanced cell growth, as well as hydrocarbon production. A variety of carbon sources, such as C1–C6, lactose and sucrose, have been demonstrated to achieve a decrease in mass doubling time of the microalgae from 6–7 days to less than 2–3 days [19]. Even though we found that a deficiency of nitrogen supply favors lipid accumulation [20], nitrogen is still essential for growth. Studies on the nitrogen source as NO<sub>3</sub><sup>-</sup>, NO<sub>2</sub><sup>-</sup>, and NH<sub>3</sub> indicate that the critical factor in regulating nitrogen metabolism in *Botryococcus braunii* is the nitrate uptake system. Thereby nitrogen is generally supplied as nitrate salts. An initial NO<sub>3</sub><sup>-</sup> concentration of ≥0.2 g·L<sup>-3</sup> favors hydrocarbon production [21].

Brenckman et al. found that with 1 g·L<sup>-3</sup> KNO<sub>3</sub> the hydrocarbon production after 30 days was 4.8 g·L<sup>-3</sup>, while about the same level of hydrocarbon production (4.5 g·L<sup>-3</sup>) was obtained after 30 days if the initial concentration of KNO<sub>3</sub> in the culture media was 3 g·L<sup>-3</sup>. This is because high concentration of nitrate interferes with hydrocarbon production of *Botryococcus braunii* [22]. Phosphorus is also essential for *Botryococcus braunii* growth, and usually supplied in the form of K<sub>2</sub>HPO<sub>4</sub>. However, active growth persists after complete depletion of phosphate in the medium. In the early exponential phase, phosphate levels in the medium can drop to below the detection limit (0.5 mg·L<sup>-3</sup>) [23].

*Botryococcus braunii* can rapidly assimilate phosphate even over the cellular metabolic requirement. Cells store the excessive phosphate as intracellular granules. The excessive phosphate is stored as intracellular granules. When the extracellular phosphate is depleted, the cells will begin to use this phosphate reserve. As discussed before, high light intensity may increase the carotenoid/chlorophyll ratio, and thus induce the color change in cell colony [24]. Experiments showed that when *Botryococcus braunii* are exposed constantly to a light intensity within the range of 25 to 72 μE·m<sup>-2</sup>·s<sup>-1</sup>, carbohydrate concentration, intracellular nitrogen, and phosphorus content drop [25].

*Botryococcus braunii* that is exposed to a high light intensity during early stage of cell culture could reach a higher biomass density (7 g·L<sup>-3</sup>) and hydrocarbon content (approximately 50% of cell dry weight) compared to cells that had been exposed to low light intensity (3 klx, or ~42 μE·m<sup>-2</sup>·s<sup>-1</sup>) [26]. However, it is showed that high intensity would induce accumulation of carotenoids over chlorophyll, where photoinhibition took place during the exponential phase of cell growth. In controlled close system, There are two approaches to overcome the overexposure of *Botryococcus braunii*;

1). Partial shading of the bioreactor [26]. 2). Adopting a diurnal light cycle (16 h light/8 h dark).

The pH of culture medium is usually adjusted to between pH 7 and 7.5 before cultivation. A rise in pH is observed during active growth of *Botryococcus braunii*, and then a slight drop will be seen later [23]. Similar trends in pH are commonly observed in CO<sub>2</sub> enriched cultures during exponential growth. Experiments showed that the optimal temperature for cell growth is 25°C [27].

### *B. Influence of nutrients on micro algal growth*

The influence of the media constituent's potassium nitrate, magnesium sulphate, dihydrogen potassium phosphate and ferric citrate on growth and lipid production in *B. braunii* was investigated using response surface methodology (RSM) [31]. In this study nutrients in combination showed positive effect on the biomass growth compared to the individual nutrients operation [Tab:1].



## International Journal for Research in Applied Science & Engineering Technology (IJRASET)

Table 1: effect of variation in nutrient sources on biomass and lipid production.

#	Experimental deviation	Nutrient source	Chemical Concentration
1	Control	Sewage Water (SW)	138 mg nitrates/L, 58 mg phosphates/L
2	Carbon	SW +glucose	600 mg glucose/L
3	Nitrogen	SW +Potassium nitrate	600 mg KNO <sub>3</sub> /L
4	Phosphorus	SW +inorganic phosphate	600 mg K <sub>2</sub> PO <sub>4</sub> /L
5	Carbon + nitrogen	SW + glucose +potassium nitrate	300 mg C <sub>6</sub> H <sub>12</sub> O <sub>6</sub> + 300 mg KNO <sub>3</sub>
6	Nitrogen +phosphorus	SW + pot. nitrate + inorg. phosphate	300 mg/L KNO <sub>3</sub> + 300 mg/L K <sub>2</sub> PO <sub>4</sub>
7	Carbon + phosphorus	SW + glu + inorganic phosphate	300 mg/L C <sub>6</sub> H <sub>12</sub> O <sub>6</sub> +300 mg K <sub>2</sub> PO <sub>4</sub> /L
8	Carbon + nitrogen + phosphorus	SW +glu +potassium nitrate+ inorganic phosphate	200 mg/L C <sub>6</sub> H <sub>12</sub> O <sub>6</sub> + 200 mg/L KNO <sub>3</sub> + 200 mg/L K <sub>2</sub> PO <sub>4</sub>

Among the individual conditions, presence of phosphorus (P) showed good biomass growth (1.2 g/l) followed by carbon and nitrogen (C+N) conditions (0.9 g/l). While among the nutrients in combination, N +P condition (1.4g/l) showed high biomass growth followed by C+P (1.2 g/l), C + N+ P (1.0 g/l) & C+N (0.7 g/l) as shown in [Fig:1]. Control condition with sewage water showed comparatively less biomass growth (0.5g/l). Highest biomass concentration observed in P as a individual nutrient condition might be due to the biomass enriching ability of P in its inorganic form. [28,30]. The next best conditions noticed were with N and C. Where carbon is considered as a key factor governing the growth pattern of microalgae. While nitrogen in the form of nitrates get assimilated in the microalgal cells and are used as a nutrient source for enhancing algal growth. Higher biomass growth resulting in N + P as seen in [Fig:2] condition might be due to the involvement of both the nutrients in the growth enhancement. Good algal biomass growth observed with C + P combination might be due to the individual function of both the nutrients facilitating cell division of algae cells. Likewise, C + N combination showed its influence on biomass increment due to the involvement of both the components in the synthesis of amino acids, chlorophyll and other nitrogen containing compounds. C + N + P combinations comprises of all the major nutrients viz., C, N, P and K essential for growth.




Fig: 1 Biomass yield of B.braunii culture grown in waste waster under the influence of different nutrients.

## International Journal for Research in Applied Science & Engineering Technology (IJRASET)




Fig: 2 Comparison of Biomass yield of B.braunii culture grown in waste waster under the influence of N+P with that of control.

Dayananda et al.,[32]; reported the biomass yield of 2.0 and 2.8 g/l in B. braunii culture (SAG 30.81 and LB-572) treaded with different levels of BG11 media. The increase in the biomass yield of B. braunii under light and dark conditions was reported by [33]. According to [34] the maximum biomass of B. braunii is 2.3 g/l. Shen et al.[35] reported dry biomass concentration of up to 2.543 g L-1 for B. braunii. Nitrates & phosphates have positive effects on algal growth [36]. These variables affect algae growth independent of each other without any interaction among them.

Lipid productivities at the end of growth were high compared to initial values which might be due to initialization of reserve food material conversion to lipid granules along with the synthesis of carbohydrates as a part of algal metabolism. While at the end of growth, all the available nutrients initiated towards deprivation and stimulated the lipid synthetic pathways enhancing lipid accumulation. The lipid productivities were noticed to be maximum with C condition (70.0%) followed by N (55%), N+P (50%), control (45%), NPK (42%), C+ N +P (40%), P (31%), C +N (30%), and C +P (30%) as seen in [Fig:3]




Fig: 3 Lipid yield of B.braunii culture grown under the influence of different experiemtal vaiation.

Individual nutrients showed relatively higher lipid accumulation compared to nutrients operated in combinations. However, both individual nutrients and nutrients in combination showed uniformity in the order of lipid productivity. When each individual nutrient condition is considered, C condition evidenced higher lipid productivity which might be due to the storage of total lipids as neutral lipids instead utilizing for the formation of photosynthetic apparatus.

Dayananda et al.[37] researching Botryococcus in Chu 13 media and 2% CO2 receives similar results in terms of chlorophyll (9.8) and carotenoids (1.8). Anitha et al. [38] reveals that at decreasing concentration of nitrogen sources there was a decreased growth, chlorophyll and biomass. Nitrogen starvation also triggered a rapid decline in nitrogen containing compound such as photosynthetic pigments, causing complete loss of photosynthetic efficiency. Rao et al. [39] also receive close to our values of chlorophyll (11.6) and carotenoids (2.5) as shown in [Fig: 4 & 5]. A high intensity of light increases the carotenoid-to-chlorophyll ratio, and this affects the color of algal colonies [40].

## International Journal for Research in Applied Science & Engineering Technology (IJRASET)




Fig: 4 Chlorophyll content of *B. braunii* culture grown in waste water.




Fig: 5 Carotenoids content of *B. braunii* culture grown in waste water.

Taking into account the carotenoids of *B. braunii* again higher values (2.5) occurred in cultures grown in enriched with nitrates medium [Fig:5]. Lutein is the major carotenoid among the total carotenoids from *B. braunii* as reported by [41]. The green algae *B. braunii* has received much attention because it contained unusually high levels of hydrocarbons ranging from 15-75% of dry wt [42].

#### IV. CONCLUSION

Microalgae are termed as ideal candidates for biofuel production due to their ability to utilize sewage water, nutrients and CO<sub>2</sub> to make the process economically viable and carbon neutral. Experiments performed with variation in nutrient supplements showed positive influence on both biomass growth and lipid accumulation. The obtained results show that the *B. braunii* develops better in sewage water, as larger values are observed in the biomass and in the percentage of lipids. Chlorophyll content in all cultures follows the dynamics of variation of the curves of growth.

#### REFERENCES

- [1] Wake LV, Hillen LW. Study of a "bloom" of the oil-rich alga *Botryococcus Braunii* in the DarwinRiver Reservoir. *Biotechnol. Bioeng.*, 22:1637–56 (1980)
- [2] Wake LV, Hillen LW. Nature and hydrocarbon content of blooms of the alga *Botryococcus Braunii* occurring in Australian freshwater lakes. *Aust. J. Mar. Freshwater Res.*, 32(3): 353–367 (1981)
- [3] Aaronson S, Berner T, Gold K, Kushner L, Patni NJ, Repak A, Rubin D. Some observations on the green planktonic alga, *Botryococcus Braunii* and its bloom form. *J. Plankton Res.*, 5: 693–700 (1983)
- [4] Huszar VLM, Reynolds CS. Phytoplankton periodicity and sequences of dominance in an Amazonian flood-plain lake (Lago Bata, Parà, Brazil): responses to gradual environmental change. *Hydrobiologia*, 346: 169–181 (1997)

## International Journal for Research in Applied Science & Engineering Technology (IJRASET)

- [5] Huang Y, Street-Perrott FA, Perrott RA, Metzger P, Eglinton G. Glacial-interglacial environmental changes inferred from molecular and compound-specific  $\delta^{13}\text{C}$  analyses of sediments from Sacred lake, Mt. Kenya. *Geochim. Cosmochim.*, 63: 1383–140476 (1999)
- [6] Metzger P, Largeau C. Chemicals of *Botryococcus Braunii*. In: Cohen Z (ed) *Chemicals from microalgae*. Taylor & Francis, London, pp 205–260 (1999)
- [7] Volova TG, Kalacheva GS, Zhila NO. Specificity of lipid composition in two *Botryococcus* strains, the producers of liquid hydrocarbons. *Russ. J. Plant Physiol.*, 50: 627–633 (2003)
- [8] Brown AC, Knights BA. Hydrocarbon content and its vrelationship to physiological state in the green alga *Botryococcus Braunii*. *Phytochemistry*, 8: 543–547 (1969)
- [9] Knights BA, Brown AC, Conway E, Middleditch BS. Hydrocarbons from the green form of the freshwater alga *Botryococcus Braunii*. *Phytochemistry*, 9: 1317–1324 (1970)
- [10] Metzger P, Casadevall E. Botryococcoid ethers, ether lipids from *Botryococcus Braunii*. *Phytochemistry*, 30: 1439–1444 (1991).
- [11] Gouveia L, Oliveira AC. 2009. Microalgae as raw material for biofuels production. *J. Ind. Microbiol. Biotechnol.*, 36: 269-274.
- [12] Wang B, Li Y, Wu N, Lan CQ. 2008. CO<sub>2</sub> bio-mitigation using microalgae. *Appl. Microbiol. Biotechnol.*, 79: 707–718.
- [13] Rao AR, Dayananda C, Sarada R, Shamala TR, Ravishankar GA. 2007. Effect of salinity on growth of green alga *Botryococcus braunii* and its constituents. *Biores. Technol.*, 98: 560–564.
- [14] Porra RJ, Thomson WA, Kriedemann PE. 1989. Determination of accurate extinction coefficients and simultaneous equations for assaying chlorophylls a and b extracted with four different solvents: verification of the concentration of chlorophyll standards by atomic absorption spectroscopy. *Biochimica et Biophysica Acta*, 975: 384-394.
- [15] Lichtenthaler HK. 1987. Chlorophylls and carotenoids: Pigments of photosynthetic biomembranes. *Methods in Enzymology*, 148: 350-382.
- [16] Bligh EG, Dyer WJ. 1959. A rapid method of total lipid extraction and purification. *Can. J. Biochem. Physiol.*, 37: 911-917.
- [17] Largeau C, Casadevall E, Berkaloff C, Dhamelincourt P. Sites of accumulation and composition of hydrocarbons in *Botryococcus Braunii*. *Phytochemistry*, 19: 1043–1051 (1980)
- [18] Wolf, F. R., Nanomura, A. M., and Bassham, J. A. Growth and branched hydrocarbon production in a strain of *Botryococcus Braunii*. *J. Phycol.*, 21: 388 (1985)
- [19] Weetal, H. H. 1985. Studies on nutritional requirements of the oil producing alga *Botryococcus Braunii*. *Appl. Biochem. Biotechnol.*, 11: 377 (1985)
- [20] Ben-Amotz, A., Torbene, T. G., and Thomas, W. H. Chemical profile of selected species of microalgae with emphasis on lipids. *J. Phycol.*, 21: 72 (1985)
- [21] Casadevall, E., Largeau, C., Metzger, P., Chirac, C., Berkaloff, C., and Coute, A. Hydrocarbon production by unicellular microalga *Botryococcus Braunii*. *Biosciences*, 2: 129 (1983)
- [22] Brenckman, F., Largeau, C., Casadevall, E., and Berkaloff, C. Effect of nitrogen nutrition on growth and hydrocarbon production of the unicellular microalga *Botryococcus Braunii*. *Comm. Eur. Communities, Energy Biomass*, 717 (1989)
- [23] Casadevall, E., Dif, D., Largeau, C., Gudin, C., Chamount, D., and Desanti, O. Studies on batch and continuous culture of *Botryococcus Braunii*: hydrocarbon production in relation to physiological state, cell ultrastructure and phosphate nutrition. *Biotechnol. Bioeng.*, 27: 286 (1985)
- [24] Wolf, F. R., Nanomura, A. M., and Bassham, J. A. Growth and branched hydrocarbon production in a strain of *Botryococcus Braunii*. *J. Phycol.*, 21: 388 (1985)
- [25] h, H. M., Kim, S., Park, E. R., Lee, S. T., Kwon, J. S., and Yoon, B. D. Effects of light intensity and nutrients on the growth of *Botryococcus* sp. *Misaengmul Hakhoechi*, 25: 339 (1997)
- [26] Kojima, E. and Zhang, K. Growth and hydrocarbon production by microalga *Botryococcus Braunii* in bubble column photobioreactor. *J. Bioscience Bioeng.*, 87: 811 (1999)
- [27] Lupi, F. M., Fernandes, H. M. L., Sa Correia, I., and Novais, J. M. Temperature profiles of cellular growth exopolysaccharide synthesis by *Botryococcus Braunii*. *J. Phycol.*, 3: 35 (1991) 127
- [28] Chen M, Tang H, Ma H, Holland TC, Simon Ng KY, Salley SO. Effect of nutrients on growth and lipid accumulation in the green algae *Dunaliella tertiolecta*. *Bioresour Technol* 2011;102:1649-55.
- [29] Pick U, Weiss M. Polyphosphate hydrolysis within acidic vacuoles in response to amine-induced alkaline stress in the halotolerant alga *Dunaliella salina*. *Plant Physiol* 1991;97:1234-40
- [30] Liu J, Huang J, Sun Z, Zhong Y, Jiang Y, Chen F. Differential lipid and fatty acid profiles of photoautotrophic and heterotrophic *Chlorella zofingiensis*: assessment of algal oils for biodiesel production. *Bioresour Technol* 2011;102: 106-10
- [31] Dayananda C, Sarada R, Bhattacharya S, Ravishankar G. 2005. Effect of media and culture conditions on growth and hydrocarbon production by *Botryococcus braunii*. *Process Biochem.*, 40: 3125–3131
- [32] Dayananda C, Sarada R, Bhattacharya S, Ravishankar G. 2007. Effect of media and culture conditions on growth and hydrocarbon production by *Botryococcus braunii*. *Process Biochemistry*, 40(9): 3125-3131
- [33] Tanoi T, Kawachi M, Watanabe M. 2011. Effects of carbon source on growth and morphology of *Botryococcus braunii*. *Journal of Applied Phycology*, 23(1): 25-33.
- [34] Ge Y, Liu J, Tian G. 2010. Growth characteristics of *Botryococcus braunii* 765 under high CO<sub>2</sub> concentration in photobioreactor. *Bioresour. Technol.*, 102(1): 130-13
- [35] Shen Y, Yuan W, Pei Z, Mao E. 2008. Culture of microalga *Botryococcus* in livestock wastewater. *ASABE*, 51(4): 1395-1400.
- [36] Fried S, Mackie B, Nothwehr E. Nitrate and phosphate levels positively affect the growth of algae species found in Perry Pond. *Tillers* 2003;4:21e4.
- [37] Dayananda C, Sarada R, Kumar V, Ravishankar G. 2007a. Isolation and characterization of hydrocarbon producing green alga *Botryococcus braunii* from Indian freshwater bodies. *Electronic Journal of Biotechnology*, 10(1): 78-91.
- [38] Anitha FS, Sergio OL, Ricardo MC. 2009. Effects of nitrogen starvation on the photosynthetic physiology of a tropical marine microalga *Rhodomonas* sp. (Cryptophyceae). *Aquatic botany*, 91: 291-297.
- [39] Rao AR, Dayananda C, Sarada R, Shamala TR, Ravishankar GA. 2007. Effect of salinity on growth of green alga *Botryococcus braunii* and its constituents.



## International Journal for Research in Applied Science & Engineering Technology (IJRASET)

- Biores. Technol., 98: 560–564.
- [40] Wolf FR, Nanomura AM, Bassham JA. 1985. Growth and branched hydrocarbon production in a strain of *Botryococcus braunii*. *J. Phycol.*, 21: 388.
- [41] Ranga Rao A, Sarada R, Bascaran V, Ravishankar G. 2006. Antioxydant activity of *Botryococcus braunii* extract elucidated in vitro models. *Journal of Agricultural and Food Chemistry*, 54: 4593-4599.
- [42] Sawayama S, Inoue S, Yokoyama S. 1994. Continuous culture of hydrocarbon-rich microalga *Botryococcus braunii* in secondarily treated sewage Appl. *Microbiol. Biot.*, 41: 729-731.



10.22214/IJRASET



45.98



IMPACT FACTOR:  
7.129



IMPACT FACTOR:  
7.429



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24\*7 Support on Whatsapp)