



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 5 Issue: VII Month of publication: July 2017

DOI:

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

Antibacterial Effect of *Allium Sativum* and *Zingiber Officinale* Extracts on Some Urinary Tract Infection Causing Bacteria

Sudeshna Pattanayak¹, Pradip Samanta^{2,3}, Smarajit Maiti³, Nirmalya Kumar Sinha⁵, Shweta Parida⁶

^{1,6}Centre for Food Science & Technology, Sambalpur University, Sambalpur, India. ,

²Dept. of Microbiology, Raja N.L. Khan Women's College, Midnapore, India,

²Herbal Medicine Research Unit, Kankabati Rishi Arabinda Rural Development and Social Welfare Institute, Kankabati, PaschimMedinipur, West Bengal, India,

³Dept. of Biochemistry and Biotechnology, Cell and Molecular Therapeutics Laboratory, Oriental Institute of Science and Technology, Midnapore, West Bengal, India,

³Founder Secretary, Midnapore T^AAgricure Biotech Research Society, Mahatabpur, Midnapore, West Bengal, India,

⁵Dept. of Nutrition, Raja N.L. Khan Women's College, Midnapore, India,

⁵Public Health Research Unit, Kankabati Rishi Arabinda Rural Development and Social Welfare Institute, Kankabati, PaschimMedinipur, West Bengal, India.

Abstract: Urinary tract infections (UTIs) are an age old problem in the human civilization. UTIs are generally caused by several pathogens, amongst them the most commons are *Candida albicans*, *Enterococcus faecalis*, *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus mirabilis*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Staphylococcus saprophyticus*. UTI pathogens are gradually becoming drug resistant for the indiscriminate use of synthetic drug and also for self medication. A number of medicinal plants may have potential effect on this disease. The present study was carried out to investigate the antibacterial activities of two medicinal plants viz. *Allium sativum*, *Zingiber officinale* against three UTI causing bacteria such as *Escherichia coli*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*. The antimicrobial activity of the plant extracts were carried out by agar well diffusion assay. The study indicated that the diameter of inhibition zones (DIZ) ranging from 9 to 29 mm with the highest zone observed against *E. coli* of *Z. officinale* and *A. Sativum*. Preliminary phytochemical analysis of the plant parts showed the presence of active compounds such as phenolics, steriods, alkaloids, glycosides, and flavonoids. This study clearly indicated that there are some human edible plants in the worlds, having antimicrobial properties nearly similar to that of the market available drug.

Key words: Antibacterial activity; Diameter of inhibition zones; Phytochemicals; Urinary Tract Infection

I. INTRODUCTION

Urinary tract infections (UTIs), accounting for 25% of all infections^[1], are caused by a range of pathogens, most commons are *Candida albicans*, *Enterococcus faecalis*, *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus mirabilis*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Staphylococcus saprophyticus*^[2-5]. UTIs are commonly caused by the bacteria which enter through the urethra^[6]. UTIs commonly develop in the bladder and spread to renal tissues^[6]. This can occur in any part of the urinary tract such as bladder, kidney, ureters, and urethra^[6]. UTIs affect nearly 150 million people each year worldwide and cost the global economy in more than 6 billion US dollars.^[7] UTIs can occur in any populations and in any age groups but more prevalent among the females of reproductive age group.^[1,8] some recent study reported that 40-50% women suffer at least one clinical episode during their lifetime.^[9] The risk factors of UTIs are personal hygiene, lower socio economic status, increased parity, increased age, sickle cell trait and anaemia, lack of prenatal care, diabetes mellitus and the functional urinary tract abnormalities.^[10,11] It is responsible for poor self-esteem, impaired quality-of-life, social isolation, and depression.^[12,13]

Up till now, there are so many antimicrobial agents that have beneficial role on UTIs. However, most of them cause different adverse effects and therefore, much attention has been paid to the search for effective therapeutic agents with negligible adverse effects.^[6] Moreover, most of the isolated UTI pathogens are now resistant to most of the commonly used antibiotics especially the ampicillin, amoxicillin, cotrimoxazole, amoxicillin-clavulinate, floxacin because of unwarranted use.^[14,15] Drug resistance among gram-negative pathogens is a risk factor for inappropriate empiric treatment (IET), which in turn increases the risk for mortality.^[16]

So, there is some urgent need to restrict the unnecessary use of antibiotics and also to search out some new drugs which are inexpensive, non-invasive and side-effect less. In this regard, the traditional medicines are the best alternatives, as these medicines are used from the time of immemorial with same efficacy.^[17-19] The antimicrobial efficacy of some plants for the treatment of UTIs has been beyond belief. The plants bio-constituents have been a good source of antimicrobial agents but still many of the plant species remained unexplored. A recent study reported that there are nearly 500,000 plant species in the world, of which only 1% has been phytochemically investigated.^[20]

Keeping this in view, the present study was initiated with an aim to study the antibacterial effect of *Allium sativum* and *Zingiber officinale* extracts on some urinary tract infection causing bacteria.

II. MATERIALS AND METHODS

A. Plant Materials

Allium sativum L. (rhizome) and *Zingiber officinale* Roscoe (rhizome), commonly human edible medicinal plants, were used in this study. The plants used in this study were taxonomically identified in the Dept. of Botany, Raja N.L. Khan Women's College, Midnapore, West Bengal.

B. Preparation of Plant Extract

The collected plant materials were rinsed with sterile distilled water and kept at an incubator for 24h. The plant materials were dried (air-dry) and powdered by grinder by maintaining the standard protocol.^[21] 60 g of dried powder of each plant parts were taken and mixed separately with water and ethanol solvent respectively in 1:5 ratio. The extraction procedure was carried out in a Rotary Shaker at 37°C for 24 h. Then the extracts were filtered using Whatman No. 1 filter paper. The extracts were concentrated to dryness by a rotary evaporator at 50°C and those are kept at 4°C for drying.

C. Bacterial Strains

Three UTI causing bacterial strains (*Escherichia coli* [MTCC No. 40], *Staphylococcus aureus* [MTCC No. 3160] and *Pseudomonas aeruginosa* [MTCC No. 424]), collected from IMTECH, Chandigarh, were used in this study. All the microorganisms were incubated into Mueller-Hinton broth for 24 h at 37°C. All the cultures were kept at 4°C until further use.

D. In Vitro Antibacterial Assay by Agar Well Diffusion Method

Bacterial suspensions were spread on the Mueller-Hinton agar (MHA) plates. The wells were cut on the agar plates using a sterile cork borer. The extracts were poured into the well. To compare the antibacterial potentiality with the plant extracts, one standard antibiotic (Chloramphenicol) was used against three UTI causing bacteria.^[22] The plates were incubated at 37°C for 24 h. The diameter of inhibition zones (DIZ) were measured (in mm) and recorded.^[21]

E. Phytochemical Analysis

The phytochemicals (viz. phenolics, alkaloids, flavonoids, tannins, saponins, steroids and glycosides) present in the plant extracts were determined by following standard methods described by Harborne^[23] and Kolkate et al.^[24]

III. RESULTS

Fig 1 shows the antibacterial activity of aqueous and ethanol extract of *Z. officinale*, *A. Sativum*, against three UTI causing bacteria (viz. *E. coli*, *S. aureus*, *P. aeruginosa*). The study indicated that the DIZ ranging from 9 to 29 mm with the highest zone observed against *E. coli* of *Z. officinale* and *A. Sativum* (Fig - 1). It was also noted that this plant is highly effective in the case of *S. aureus*, but no such effectiveness was observed in the cases of *P. aeruginosa*. Phytochemical studies revealed that alkaloids, steroids, glycosides are present in these selected plants (Table 1).

IV. DISCUSSION

Medicinal plants have been used for traditional treatments of numerous human diseases from ancient to modern times.^[25] Even herbal medicines were used 60,000 years back by Neanderthal man.^[26] But, it has lost its importance in twentieth century because of the advent of modern synthetic treatments.^[27] Recently, multiple drug resistance in both human as well as plant pathogens has been developed due to improper use of synthetic drugs.^[28] To combat the diminishing efficacy of existing antibiotics a diverse arsenal of new antibacterial agents is essentially required.^[29] According to World Health Organization, medicinal plants would be the best source to obtain variety of drugs.^[30] Clinical microbiologists have great interest in screening of medicinal plants for new

therapeutics.^[31] All pharmaceutical industries are now searching for medicinal plants contain some organic bio-active molecules including tannins, alkaloids, carbohydrates, terpenoids, steroids, flavonoids, phenols etc.^[32]

In the present study aqueous and ethanolic plant extracts was exhibited antibacterial activity towards all UTI pathogens, with more activity observed with ethanolic extracts. Ethanol extracts may have the potential role of extracting plant biomolecules that are actually responsible for higher antimicrobial activity than the aqueous one.^[33] Phytochemicals exert this antimicrobial activity through different mechanisms, like act by iron deprivation, hydrogen bonding or non-specific interactions with vital proteins such as enzymes.^[34] Sawyer et al.^[35] demonstrated that the main indoloquinoline alkaloid, cryptolepine, causes cell lysis and morphological changes of *S. aureus*, but the antimicrobial effects of the alkaloid may be through another mechanism, since the compound is known to be a DNA intercalator and an inhibitor of DNA synthesis through topoisomerase inhibition.

V. CONCLUSION

UTIs are an age old problem in the human civilization. UTI pathogens are gradually becoming drug resistant for the indiscriminate use of synthetic drug and also for self medication. The situation becomes worse and phytomedicines are considered to be the best useful alternatives. It has low/no side effects, low cost, easily available and used from the time of immemorial with same efficacy. But the young generation has less knowledgeable about the indigenous medicine and this knowledge is going to be lost with the aged people who got this knowledge from their forefathers.

There are some human edible plants in the worlds, having antimicrobial properties nearly similar to that of the market available laboratory use drug. The presence of phytochemicals including phenols, tannins and flavonoids may be responsible for the antibacterial activity. These are the products of secondary metabolism of the plant. The secondary metabolism of the plant can vary considerably depending on factors such as: seasonality, temperature, water availability, UV radiation, nutrients, altitude, air pollution and even induction by mechanic stimuli or attack by pathogens. Further more there are a strict procedure while collecting the plant or plant materials for the phytomedicine as the phytochemicals are altered with the times, age of the plant and diurnal variation. A proper awareness is needed.

REFERENCES

- [1] Ahmed MA, Shukla GS, Bajaj HK. Incidence of Urinary Tract Infections and determination of their susceptibility to antibiotics among Pregnant Women. *Int J Cell Sci Biotech*. 2016; 5:12-16.
- [2] Sharma A, Chandraker S, Patel VK, Ramteke P. Antibacterial Activity of Medicinal Plants Against Pathogens causing Complicated Urinary Tract Infections. *Indian J Pharm Sci*. 2009; 71(2): 136-139.
- [3] Svanborg C, Godaly G. Bacterial virulence in urinary tract infection. *Infect Dis Clin N Am* 1997;11:513-29.
- [4] Samanta P, Sinha NK. Antimicrobial activity of five traditionally used medicinal plants on bacterial infection of urinary tract. *Int Res J Basic Appl Sci* 2016;1(2): 24-28.
- [5] Divya P, Pavithra DP. Antimicrobial susceptibility pattern of *Escherichia coli* isolates causing urinary tract infections at tertiary care hospital. *Int J Pharm Bio Sci*. 2015; 6(1): (B) 84 – 87.
- [6] Nabavi SF, Sureda A, Daglia M, Izadi M, Nabavi SM. Cranberry for Urinary Tract Infection: From Bench to Bedside. *Curr Top Med Chem*. 2017;17(3):331-339.
- [7] Stamm WE, Norrby SR. Urinary tract infections: disease panorama and challenges. *J Infect Dis*. 2001;183 (Suppl 1):S1-S4
- [8] Karki A, Tiwari BR, Pradhan SB. Study of Bacteria Isolated from Urinary Tract Infections and Their Sensitivity Pattern. *J Nepal Med Assoc*. 2004; 43: 200-203.
- [9] Singh B, Tilak R, Srivastava RK, Katiyar D. Urinary tract infection and its risk factors in women: An appraisal. *J Pure Appl Microbiol*. 2014; 8(5):1-8.
- [10] Nowiciki B. Urinary tract infections in pregnant women: Old dogmas and current concepts regarding pathogenesis. *Curr Infect Dis Rep*. 2002; 4(6): 529-535.
- [11] Manjula NG, Math GC, Patil SA, Gaddad SM, Shivannavar CT. Incidence of Urinary Tract Infections and Its Aetiological Agents among Pregnant Women in Karnataka Region. *Advances in Microbiology*. 2013;3: 473-478.
- [12] Fonda D. Promoting continence as a health issue. *Eur Urol*. 1997;32:28-32
- [13] Vyas S, Varshney D, Sharma P, Juyal R, Nautiyal V, Shrotriya VP. An Overview of the Predictors of Symptomatic Urinary Tract Infection Among Nursing Students. *Ann Med Health Sci Res*. 2015; 5(1): 54-58.
- [14] Kidwai SS, Nageen A, Ghaznavi S, Bashir F, Ara J. Antibiotic susceptibility in commonly isolated pathogens from urinary tract infection in a cohort of subjects from low socioeconomic strata. *Pak J Med Sci*. 2017;33(2):254-259.
- [15] Masika WG, O'Meara WP, Holland TL, Armstrong J. Contribution of urinary tract infection to the burden of febrile illnesses in young children in rural Kenya. *PLoS One*. 2017;12(3):e0174199.
- [16] Zilberberg MD, Nathanson BH, Sulham K, Fan W, Shorr AF. Carbapenem resistance, inappropriate empiric treatment and outcomes among patients hospitalized with Enterobacteriaceae urinary tract infection, pneumonia and sepsis. *BMC Infect Dis*. 2017;17(1):279.
- [17] Das DC, Sinha NK, Chattopadhyay JC, Das M, Samanta P. The use of Medicinal Plants for the treatment of Gonorrhoea and Syphilis in South West Bengal of India. *Int J Phytomed*. 2013;5(1): 14-7.
- [18] Das DC, Sinha NK, Das M. The use of medicinal plants for the treatment of Gynaecological disorders in the Eastern parts of India. *Indian J ObstetGynaecol Res*. 2015; 2(1):16-27.

- [19] Das DC, Sinha NK, Patsa MK, Das M. Investigation of herbals for the treatment of leucorrhoea from south West Bengal, India. *Int J Bioassay*. 2015; 4(11): 4555-9.
- [20] Kim HS. Do not put too much value on conventional medicines. *J Ethnopharmacol*. 2005; 100(1-2): 37-39.
- [21] Al-Talib H, Ali NDM, Suhaimi MH, Rosli SSN, Othman NH, Mansor NAS, Shah AKS, Ariffin NS, Al-Khateeb A. Antimicrobial effect of Malaysian vegetables against enteric bacteria. *Asian Pac J Trop Biomed* 2016; 6(3): 211–215
- [22] Sharma A, Verma R, Ramteke P. Antibacterial activity of some medicinal plants used by tribals against UTI causing pathogens. *World Appl Sci J*. 2009; 7(3): 332-339.
- [23] Harborne J B. *Phytochemical methods: a guide to modern techniques of plant analysis*. 3rd ed. New York, NY: Chapman and Hall. 1998. pp. 49–188.
- [24] Kolkate CK, Purohit AP, Gokhale SB. *Pharmacognosy*. 22nd edition. Pune, India: Nirani Publication. 2003.
- [25] Gaherwal S, Johar F, Wast N, Prakash MM. Anti-Bacterial Activities of *Allium sativum* Against *Escherichia coli*, *Salmonella Ser. Typhi* and *Staphylococcus aureus*. *Int J Microbiol Res*. 2014;5 (1): 19-22.
- [26] Solecki RS. Shanidar IV, a neanderthal flower burial in northern Iraq. *Science*. 1975;190(28): 880-881.
- [27] Iyabo OG, Vianne OZ, Ikhiwili OM, Temitope AB, Adetutu AA. In vitro antibacterial and antifungal activities of *Chrysophyllum albidum* and *Diospyros monbutensis* leaves. *J Pharmacognosy Phytother*. 2016;8(1):1-7.
- [28] Hart CA, Karriuri S. Antimicrobial resistance in developing countries. *BMJ*. 1998;317:421-452.
- [29] Chopra I, Hodgson J, Metcalf B and Poste G. The search for antibacterial agents effective against bacteria resistance to multiple antibiotics. *Antimicrob Agent Chemother*. 1997;41:497-503.
- [30] Akeel RA, Al-Sheikh Y, Mateen A, Syed R, Janardhan K, Guptab VC. Evaluation of antibacterial activity of crude protein extracts from seeds of six different medical plants against standard bacterial strains. *Saudi J Biol Sci*. 2014; 21(2): 147–151.
- [31] Periyasamy A, Rajkumar, Mahalingam K. Phytochemical screening and antimicrobial activity from five Indian medicinal plants against human pathogens. *Middle-East J Sci Res*. 2010;5(6):477-482.
- [32] Mandal S, Patra A, Samanta A, Roy S, Mandal A, Das Mahapatra T, Pradhan S, Das K, Nandi DK. Analysis of phytochemical profile of *Terminalia arjuna* bark extract with antioxidative and antimicrobial properties. *Asian Pac J Trop Biomed*. 2013; 3(12): 960-966.
- [33] Samanta P, Sinha NK. Antimicrobial activity of five traditionally used medicinal plants on bacterial infection of urinary tract. *Int Res J Basic Appl Sci*. 2016;1(2):24-28.
- [34] Scalbert A. Antimicrobial properties of tannins. *Phytochemistry*. 1991;30, 3875-3883.
- [35] Sawyer IK, Berry MI, Ford JL. The killing effect on *Staphylococcus aureus*. *Let Appl Microbiol*. 2005;40: 24-29.

Table 1. Phytochemical analysis of two medicinal plants in ethanolic extracts

Phytochemicals	<i>Zingiberofficinale</i>	<i>Allium sativum</i>
Phenolics	-	-
Alkaloids	+	+
Flavonoids	-	-
Tannins	-	-
Saponins	-	-
Steroids	+	+
Glycosides	+	+




Fig 1. The yield of extract of *Zingiberofficinale* and *Allium sativum* against three UTI causing bacteria



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)