



# IJRASET

International Journal For Research in  
Applied Science and Engineering Technology



---

# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

---

**Volume: 5      Issue: VII      Month of publication: July 2017**

**DOI:**

**[www.ijraset.com](http://www.ijraset.com)**

**Call:  08813907089**

**E-mail ID: [ijraset@gmail.com](mailto:ijraset@gmail.com)**

# Certain Transformation Formulae for Basic Hypergeometric Series

Rajesh Pandey<sup>1</sup>, Pravesh Chandra Srivastava<sup>2</sup>

<sup>1</sup>Department of Applied Science, Institute of Engineering & Technology, Sitapur Road, Lucknow 226021 India.

<sup>2</sup>Department of Applied Science, IMS Engineering College, Ghaziabad 201013, India.

**Abstract:** In this paper, general transformation formulae for basic hypergeometric series of two variables have been established. Special cases have also been studied.

**Keywords:** Basic hypergeometric series, Transformation formulae, Summation, Basic Analogue, Parameters

## I. INTRODUCTION

Jeugt, Pitre and Srinivasa Rao [1] obtain certain summation theorems for double and triple hypergeometric functions. The following

interesting summation formula for double hypergeometric function has been established  $F_{i;l}^{0;3} \left[ \begin{matrix} \delta - \alpha\beta + \gamma, -p; \alpha - \delta, \beta + p, -\gamma; 1, 1 \\ \beta : \beta + \gamma ; \alpha + p \end{matrix} \right] =$

$$\frac{(\alpha)_p(\delta)_r}{(\partial)_p(\alpha)_r} \tag{1.1}$$

The basic analogue of (1.1) has been mentioned as

$$F_{i;l}^{0;3} \left[ \begin{matrix} \delta / \alpha, \beta q^r, q^{-p}; \alpha / \delta, \beta q^p, q^{-r}; q, q \\ \beta : \delta q^r ; \alpha q^p \end{matrix} \right] = \left( \frac{\delta}{\alpha} \right)^{p-r} \frac{(\alpha; q)_p(\delta; q)_r}{(\alpha; q)_p(\alpha; q)_r} \tag{1.2}$$

## II. DEFINITIONS AND NOTATIONS

The Gauss hypergeometric function is represented as:

$${}_2F_1 \left[ \begin{matrix} a, b; z \\ c \end{matrix} \right] = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n z^n}{(c)_n n!}, \tag{2.1}$$

Where,

$$(a)_n = a(a+1)\dots(a+n-1) = \frac{\Gamma(a+n)}{\Gamma(a)}, \quad (a)_0 = 1 \tag{2.2}$$

The generalised hypergeometric function is defined as:

$${}_A F_B \left[ \begin{matrix} (a); z \\ (b) \end{matrix} \right] = \sum_{n=0}^{\infty} \frac{[(a)]_n z^n}{[(b)]_n n!} \tag{2.3}$$

Where (a) stands for A-parameters of the form  $a_1, a_2, \dots, a_A$ . A double hypergeometric function is defined by

$$F_{C:D}^{A:B:B'} \left[ \begin{matrix} (a) : (b); (b'); x, y \\ (c) : (d); (d') \end{matrix} \right] = \sum_{m,n=0}^{\infty} \frac{[(a)]_{m+n} [(b)]_m [(b')]_n x^m y^n}{[(c)]_{m+n} [(d)]_m [(d')]_n m! n!} \tag{2.4}$$

And in case of  $B=B', D=D'$ , we simply write the function as,

$$F_{C:D}^{A:B} \left[ \begin{matrix} (a) : (b); (b'); x, y \\ (c) : (d); (d') \end{matrix} \right]$$

The basic analogue of (2.3) known as generalized basic hypergeometric function is defined by,

$${}_A F_B \left[ \begin{matrix} (a); z \\ (b); q^\lambda \end{matrix} \right] = \sum_{n=0}^{\infty} \frac{[(a)]_n z^n q^{\lambda n(n-1)/2}}{[(b)]_n (q)_n}, \tag{2.5}$$

Where (a) stands for A-parameters of the form  $a_1, a_2, \dots, a_A$  : and

$$(a)_n = (a; q)_n = (1-a)(1-aq)\dots(1-aq^{n-1}); (a; q)_0 = 1 .$$

The basic double hypergeometric function is defined as:

$$F_{C;D:D'}^{A:B:B'} \left[ \begin{matrix} (a) : (b); (b'); x, y \\ (c) : (d); (d'); q^\lambda, q^\mu \end{matrix} \right] = \sum_{m,n=0}^{\infty} \frac{[(a)]_{m+n} [(b)]_m [(b')]_n x^m y^n}{[(c)]_{m+n} [(d)]_m [(d')]_n (q)_m (q)_n} q^{\lambda m(m-1)/2 + \mu n(n-1)/2} \tag{2.6}$$

### III.MAIN RESULTS

#### A. Analytic Proof of (1.2)

In this section we shall give analytic proof of (1.2). Let us represent the left hand side of (1.2) by  $\Omega$ , then

$$\begin{aligned} \Omega &= \sum_{m,n=0}^{p,r} \frac{\left(\frac{\delta}{\alpha}\right)_m (\beta q^r)_m (q^{-p})_m \left(\frac{\alpha}{\delta}\right)_n (\beta q^p)_n (q^{-r})_n q^{m+n}}{(\beta)_{m+n} (\delta q^r)_m (\alpha q^p)_n (q)_m (q)_n} \\ &= \sum_{m=0}^p \frac{(\delta/\alpha)_m (\beta q^r)_m (q^{-p})_m q^m}{(\beta)_m (\delta q^r)_m (q)_m} {}_3\Phi_2 \left[ \begin{matrix} \alpha/\delta, \beta q^p, q^{-r}; q \\ \beta q^m, \alpha q^p \end{matrix} \right] \end{aligned} \tag{3.1}$$

Now, transforming the inner  ${}_3\Phi_2$  series, we get

$$\Omega = \sum_{m=0}^p \frac{\left(\frac{\delta}{\alpha}\right)_m (\beta q^r)_m (q^{-p})_m q^m \left(\frac{\beta \delta q^m}{\alpha}\right)_r \left(\frac{\alpha}{\delta}\right)^r}{(\beta)_m (\delta q^r)_m (q)_m (\beta q^m)_r} \times {}_3\Phi_2 \left[ \begin{matrix} q^{-r}, \alpha/\delta, \alpha/\beta; q^{1+p-m} \\ \alpha q^p, \alpha/\beta \delta, q^{1-m-r} \end{matrix} \right] \tag{3.2}$$

Summing the inner  ${}_3\Phi_2$  series with the help of Saalschütz summation formula, we get,

$$\Omega = \frac{(\delta)_{p+r} (\beta)_{p+r} (\alpha)_p \left(\frac{\beta \delta}{\alpha}\right)_p \left(\frac{\beta \delta}{\alpha}\right)_r \left(\frac{\alpha}{\delta}\right)^r}{(\alpha)_{p+r} (\beta \delta/\alpha)_{p+r} (\delta)_p (\beta)_p (\beta)_r} \times {}_3\Phi_2 \left[ \begin{matrix} q^{-r}, \delta/\alpha, \beta \delta/\alpha, q^r; q \\ \delta q^r, \beta \delta/\alpha \end{matrix} \right] \tag{3.3}$$

Again, applying the transformation formula and then summing the  ${}_3\Phi_2$  series on the right hand side of (3.3) with the help of, we get the right hand side of (1.2).

#### B. General Transformation Formula

We shall establish the following general transformation formula:

$$\begin{aligned} \Phi_{C;D+1;D'+1}^{A:B+1;B'+1} \left[ \begin{matrix} (a) : (b), \alpha; (b'), \delta; \delta z_1/\alpha, \alpha z_2/\alpha \\ (c) : (d), \alpha; (d'), \alpha; q, q \end{matrix} \right] &= \sum_{m,n=0}^{\infty} \frac{[(a)]_{m+n} [(b)]_m [(b')]_n (\beta)_{m+n} (\alpha)_m (\delta/\alpha)_m (\alpha/\delta)_n (-z_1)^m (-z_2)^n}{[(c)]_{m+n} [(d)]_m [(d')]_n (\alpha)_{m+n} (\delta)_{m+n} (\beta)_m (\beta)_n (q)_m (q)_n} \times \\ &\times \Phi_{C;D+2;D'+2}^{A:B+2;B'+2} \left[ \begin{matrix} (a)q^{m+n} : (b)q^m, \beta q^{m+n}, \alpha q^m; (b')q^n, \beta q^{m+n}, \delta q^n; z_1, z_2 \\ (c)q^{m+n} : (d)q^m, \beta q^m, \alpha q^{m+r}; (d')q^n, \beta q^m, \delta q^{m+n}; q, q \end{matrix} \right] \end{aligned} \tag{3.4}$$

Proof :

Let us represent the left hand side of (3.4) by  $\wedge$ , then

$$\wedge = \sum_{p,r} \frac{[(a)]_{p+r} [(b)]_p [(b')]_r z_1^p z_2^r q^{p(p-1)/2 + r(r-1)/2}}{[(c)]_{p+r} [(d)]_p [(d')]_r (q)_p (q)_r} \left\{ \frac{(\alpha)_p (\delta)_r}{(\delta)_p (\alpha)_r} \left(\frac{\delta}{\alpha}\right)^{p-r} \right\}.$$

Putting the value of  $\left\{ \frac{(\alpha)_p (\delta)_r}{(\delta)_p (\alpha)_r} \left(\frac{\delta}{\alpha}\right)^{p-r} \right\}$ , in the form of double series from (1.2) we get,

$$\wedge = \sum_{p,r} \frac{[(a)]_{p+r} [(b)]_p [(b')]_r z_1^p z_2^r q^{p(p-1)/2+r(r-1)/2}}{[(c)]_{p+r} [(d)]_p [(d')]_r (q)_p (q)_r} \times \sum_{m=0}^p \sum_{n=0}^r \frac{(\delta/\alpha)_m (\beta q^r)_m (q^{-p})_m (\alpha/\delta)_n (\beta q^p)_n (q^{-r})_n}{(\beta)_{m+n} (\delta q^r)_m (\alpha q^p)_n (q)_m (q)_n} q^{m+n}$$

Now changing the order of summations and putting  $p+m$ ,  $r+n$  for  $p$  and  $r$  respectively, we get the right hand side of (3.4) after some simplifications.

**IV. SPECIAL CASES OF (3.4) AND RESULTS:**

A. Putting  $A = C = 0, B = B' = D = D' = 1, b_1 = \delta, d_1 = \alpha$  and  $d_1' = \delta$  in (3.4) we get,

$$\sum_{u,v=0}^{\infty} \frac{q^{u(u-1)/2+v(v-1)/2}}{(q)_u (q)_v} \left(\frac{\delta}{\alpha}\right)^{u-v} z_1^u z_2^v$$

Certain transformation formulae for basic hypergeometric series

$$= \sum_{m,n=0}^{\infty} \frac{(\delta)_m (\alpha)_n (\beta)_{m+n} (\delta/\alpha)_n (\alpha/\delta)_n (-z_1)^m (-z_2)^n}{(\alpha)_{m+n} (\delta)_{m+n} (\beta)_m (q)_m (q)_n} {}_2\Phi_2 \left[ \begin{matrix} \delta q^m, \beta q^{m+n}; z_1 \\ \beta q^m, \alpha q^{m+n}; q \end{matrix} \right] {}_2\Phi_2 \left[ \begin{matrix} \alpha q^n, \beta q^{m+n}; z_2 \\ \beta q^n, \delta q^{m+n}; q \end{matrix} \right] \quad (3.5)$$

Taking  $z_1 = z_2$  and then equating the coefficients of  $z^{u+v}$  of both sides we get the following summation formula:

$$\sum_{r=0}^u \sum_{s=0}^v \frac{(q^{-u})_r (q^{-v})_s \left(\frac{q^{1-u-v}}{\beta}\right)_{r+s} \left(\frac{q^{1-u-v}}{\alpha}\right)_s \left(\frac{q^{1-u-v}}{\delta}\right)_r (\alpha q^{u+v})^r (\delta q^{u+v})^s}{\left(\frac{q^{1-u-v}}{\beta}\right)_r \left(\frac{\alpha}{\delta} q^{1-u}\right)_r \left(\frac{q^{1-u-v}}{\beta}\right)_s \left(\frac{\alpha}{\delta} q^{1-u}\right)_s} (q)_r (q)_s q^{rs}$$

$$= \frac{(\alpha)_{u+v} (\delta)_{u+v} (\beta)_u (\beta)_v (-)^{u+v} q^{(u/2)+(v/2)} (\delta/\alpha)^{u+v}}{(\beta)_{u+v} (\delta)_u (\delta/\alpha)_u (\alpha)_v (\alpha/\delta)_v} \quad (3.6)$$

B. PUTTING  $A = C = 0, B = B' = D = D' = 1, b_1 = \beta, d_1 = \alpha, b_1' = \beta, d_1' = \alpha$  IN (3.4) WE GET :

$${}_1\Phi_1 \left[ \begin{matrix} \beta; z_1 & \delta/\alpha \\ \partial; q \end{matrix} \right] {}_1\Phi_1 \left[ \begin{matrix} \beta; z_2 & \alpha/\delta \\ \alpha; q \end{matrix} \right]$$

$$= \sum_{m,n=0}^{\infty} \frac{(\beta)_{m+n} (\delta/\alpha)_m (\alpha/\delta)_n (-z_1)^m (-z_2)^n}{(\delta)_{m+n} (\alpha)_{m+n} (q)_m (q)_n} \times {}_1\Phi_1 \left[ \begin{matrix} \beta q^{m+n}; z_1 \\ \alpha q^{m+n}; q \end{matrix} \right] {}_1\Phi_1 \left[ \begin{matrix} \beta q^{m+n}; z_2 \\ \delta q^{m+n}; q \end{matrix} \right] \quad (3.7)$$

Taking  $z_1 = -\alpha/\beta$ , and  $z_2 = -\delta/\beta$ , in (3.7) and summing  ${}_1\Phi_1$  series of both sides we get an identity:

$$\sum_{m,n=0}^{\infty} \frac{(\beta)_{m+n} (\delta/\alpha)_m (\alpha/\delta)_n (\alpha/\beta)^m (\delta/\beta)^n}{(q)_m (q)_n} = 1 \quad (3.8)$$

Taking  $\beta \rightarrow \infty$  in (3.8) we get :

$$\sum_{m,n=0}^{\infty} \frac{(\delta/\alpha)_m (\alpha/\delta)_n (\alpha)^m (\delta)^n (-)^{m+n} (q)^{(m+n)(m+n-1)/2}}{(q)_m (q)_n} = 1 \quad (3.9)$$

Replacing  $q$  by  $q^2$  in (3.9) and then taking  $\alpha = \delta q$  and finally putting  $\delta = 1$ , we get:

$$\sum_{m,n=0}^{\infty} \frac{(q^{-1}; q^2)_m (q; q^2)_n q^m (-)^{m+n} q^{(m+n)(m+n-1)}}{(q^2; q^2)_m (q^2; q^2)_n} = 1 \quad (3.10)$$

C. PUTTING  $z_1/\beta, z_2/\beta$  FOR  $Z_1$  AND  $Z_2$  IN (3.7) AND THEN TAKING  $\beta \rightarrow \infty$  WE GET :

$${}_0\Phi_1 \left[ \begin{matrix} -; -z_1 \delta/\alpha \\ \delta; q^2 \end{matrix} \right] {}_0\Phi_1 \left[ \begin{matrix} -; -z_2 \alpha/\delta \\ \delta; q^2 \end{matrix} \right]$$

$$= \sum_{m,n=0}^{\infty} \frac{(\delta/\alpha)_m (\alpha/\delta)_n z_1^m z_2^n q^{(m+n)(m+n-1)/2} (-)^{m+n}}{(\delta)_m (\alpha)_{m+n} (q)_m (q)_n} \times {}_0\Phi_1 \left[ \begin{matrix} -; -z_1 q^{m+n} \\ \alpha q^{m+n}; q^2 \end{matrix} \right] {}_0\Phi_1 \left[ \begin{matrix} -; -z_2 q^{m+n} \\ \delta q^{m+n}; q^2 \end{matrix} \right] \quad (3.11)$$

Again taking  $\delta = q^{1+v_1}, \alpha = q^{1+v_2}, z_1 = \frac{x^2}{4} q^{1+v_2}, z_2 = \frac{y^2}{4} q^{1+v_1}$  in (3.11), we get:

$$\begin{aligned} & {}_0\Phi_1 \left[ \begin{matrix} -; \frac{x^2}{4} q^{1+v_1} \\ q^{1+v_1}; q^2 \end{matrix} \right] {}_0\Phi_1 \left[ \begin{matrix} -; \frac{y^2}{4} q^{1+v_2} \\ q^{1+v_2}; q^2 \end{matrix} \right] \\ &= \sum_{m,n=0}^{\infty} \frac{(q^{v_1-v_2}; q)_m (q^{v_2-v_1}; q)_n \left(\frac{x}{2}\right)^{2m} \left(\frac{y}{2}\right)^{2n} q^{(1+v_2)m} q^{(1+v_1)n}}{(q^{1+v_1}, q^{1+v_2}; q)_{m+n} (q)_m (q)_n} \times \\ & q^{(m+n)(m+n-1)/2} {}_0\Phi_1 \left[ \begin{matrix} -; \frac{x^2}{4} q^{1+v_2+m+n} \\ q^{1+v_2+m+n}; q^2 \end{matrix} \right] {}_0\Phi_1 \left[ \begin{matrix} -; \frac{y^2}{4} q^{1+v_1+m+n} \\ q^{1+v_1+m+n}; q^2 \end{matrix} \right] \end{aligned} \quad (3.12)$$

Changing  ${}_0\Phi_1$  series into Bessel function of second kind defined by,

$$J_{v_1}^{(2)}(x; q) = \frac{(q^{1+v_1}; q)_{\infty} (x/2)^v}{(q; q)_{\infty}} \sum_{n=0}^{\infty} \frac{(-)^n (x^2/4)^n (q^{1+v_1})^n q^{n^2-n}}{(q; q)_n (q^{1+v_1}; q)_n}$$

We get :

$$\begin{aligned} J_{v_1}^{(2)}(x; q) J_{v_2}^{(2)}(y; q) &= \sum_{m,n=0}^{\infty} \frac{(q^{v_1-v_2}; q)_m (q^{v_2-v_1}; q)_n (-)^{m+n} (x/y)^{m-n}}{(q)_m (q)_n} \\ &\times q^{(1+v_1)n+(1+v_2)m} J_{v_1+m+n}^{(2)}(y) J_{v_2+m+n}^{(2)}(x) \end{aligned} \quad (3.13)$$

A number of similar other interesting results can also be deduced.

### V. CONCLUSION

In this paper, an attempt has been made to give the analytic proof of (1.2) we shall also make use of (1.2) to establish a general transformation formula for basic hypergeometric series of two variables. Special cases have also been studied and some very interesting and new results have been obtained.

### VI. ACKNOWLEDGEMENT

My special thanks to my Ph.D supervisor Dr. Rajesh Pandey Department of Applied Science, Institute of Engineering & Technology (IET) Lucknow for his guidance and encouragement. It was not possible to complete this research paper without his necessary support. I am extremely grateful for his constructive support.

### REFERENCES

- [1] Gasper, G. and Rahman, M. (1991): Basic hypergeometric series, Cambridge University Press.
- [2] Agarwal, R.P., Manocha, H.L. and Rao, K.Srinivas (2001); Selected Topics in special functions, Allied Publisher Limited, New Delhi.
- [3] Agarwal, R.P.: Generalized hypergeometric series and its application to the theory of combinatorial analysis and partition theory (Unpublished monograph).
- [4] L.J. Slater: Generalized Hypergeometric Functions, Cambridge University Press, (1966).
- [5] S. Ramanujan : Notebook, Vol. II, Tata Institute of Fundamental Research, Bombay, (1957).
- [6] Jeugt, J. Van der, Pitre, Sangita N, Srinivasa Rao, K. (To appear in J. Phys. A.).





10.22214/IJRASET



45.98



IMPACT FACTOR:  
7.129



IMPACT FACTOR:  
7.429



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24\*7 Support on Whatsapp)