



# IJRASET

International Journal For Research in  
Applied Science and Engineering Technology



---

# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

---

**Volume: 5      Issue: VIII      Month of publication: August 2017**

**DOI: <http://doi.org/10.22214/ijraset.2017.8174>**

**[www.ijraset.com](http://www.ijraset.com)**

**Call:  08813907089**

**E-mail ID: [ijraset@gmail.com](mailto:ijraset@gmail.com)**

# Solution of Higher Order Partial Differential Equation by using Homotopy Analysis Method

Vimal P. Gohil

Assistant professor, Department of Mathematics, Government Engineering College, Bhavnagar

**Abstract:** In this paper, various types of higher order non linear partial differential equations are discussed. Also shown that homotopy analysis method applied successfully for solving higher order non linear partial differential equations including homogeneous and non-homogeneous.

**Keywords:** Homotopy analysis method, partial differential equation, homogeneous, nonlinear.

## I. INTRODUCTION

Homotopy analysis method (HAM) has been successfully employed to solve many types of non linear, homogeneous or non homogeneous, equations and systems of equations as well as problems in science and engineering . Very recently, Ahmad Bataineh et al. ([2]) presented two modifications of HAM to solve linear and non linear ODEs. The HAM contains a certain auxiliary parameter  $h$  which provides us with a simple way to adjust and control the convergence region and rate of convergence of the series solution. Moreover, by means of the so-called  $h$ -curve, it is easy to determine the valid regions of  $h$  to gain a convergent series solution. Thus, through HAM, explicit analytic solutions of non linear problems are possible.

Systems of partial differential equations (PDEs) arise in many scientific models such as the propagation of shallow water waves and the Brusselator model of the chemical reaction-diffusion model. Very recently, Batiha et al. [2] improved Wazwaz [9] results on the application of the variational iteration method (VIM) to solve some linear and non linear systems of PDEs. In [8], Saha Ray implemented the modified Adomian decomposition method (ADM) for solving the coupled sine-Gordon equation.

## II. HOMOTOPY ANALYSIS METHOD

We consider the following differential equations,

$$N_i [S_i(x, t)] = 0, i = 1, 2, \dots, n$$

Where  $N_i$  are nonlinear operators that the represents the whole equations,  $x$  and  $t$  are independent variables and  $S_i(x, t)$  are unknown functions respectively.

By means of generalizing the traditional homotopy method, Liao constructed the so-called zero-order deformation equations

$$(1 - q)L[\varphi_i(x, t, ;q) - S_{i,0}(x, t)] = qh_i N_i[\varphi_i(x, t, ;q)] \quad (1)$$

Where  $q \in [0, 1]$  is an embedding operators,  $h_i$  are nonzero auxiliary functions,  $L$  is an auxiliary linear operator,  $S_{i,0}(x, t)$  are initial guesses of  $S_i(x, t)$  and  $\varphi_i(x, t, ;q)$  are unknown functions.

It is important to note that, one has great freedom to choose auxiliary objects such as  $h_i$  and  $L$  in HAM.

When  $q = 0$  and  $q = 1$  we get by (1),

$$\varphi_i(x, t, ;0) = S_{i,0}(x, t) \text{ and } \varphi_i(x, t, ;1) = S_i(x, t)$$

Thus  $q$  increase from 0 to 1, the solutions  $\varphi_i(x, t, ;q)$  varies from initial guesses  $S_{i,0}(x, t)$  to  $S_i(x, t)$ .

Expanding  $\varphi_i(x, t, ;q)$  in Taylor series with respect to,

$$\varphi_i(x, t, ;q) = S_{i,0}(x, t) + \sum_{m=1}^{\infty} S_{i,m}(x, t) \cdot q^m \quad (2)$$

Where

$$S_{i,m}(x,t) = \left[ \frac{1}{m!} \cdot \frac{\partial^m \varnothing_i(x,t;;q)}{\partial q^m} \right]_{q=0} \tag{3}$$

If the auxiliary linear operator, initial guesses, the auxiliary parameter and auxiliary functions are properly chosen than the series equation (2) converges at  $q = 1$ .

$$\varnothing_i(x,t;;1) = S_{i,0}(x,t) + \sum_{m=1}^{\infty} S_{i,m}(x,t). \tag{4}$$

This must be one of solutions of the original nonlinear equations.

According to (3), the governing equations can be deduced from the zero-order deformation equations (1).

Define the vectors

$$\overrightarrow{S_{i,n}} = \{S_{i,0}(x,t), S_{i,1}(x,t), S_{i,2}(x,t), \dots, S_{i,n}(x,t)\}$$

Differentiating (1) m times with respect to the embedding parameter and the setting  $q = 0$  and finally dividing them by  $m!$

We have the so-called  $m^{th}$  order deformation equations

$$L[S_{i,m}(x,t) - \chi_m S_{i,m-1}(x,t)] = h_i R_{i,m}(\overrightarrow{S_{i,m-1}}) \tag{5}$$

Where

$$R_{i,m}(S_{i,m-1}) = \left[ \frac{1}{(m-1)!} \cdot \frac{\partial^{m-1} N_i[\varnothing_i(x,t;;q)]}{\partial q^{m-1}} \right]_{q=0} \tag{6}$$

$$\chi_m = \begin{cases} 0, & m \leq 1 \\ 1, & m > 1 \end{cases}$$

#### HOMOGENEOUS NON LINEAR PARTIAL DIFFERENTIAL EQUATION

Consider homogeneous linear differential equation

$$\frac{\partial u}{\partial t} - \frac{\partial u}{\partial x} \frac{\partial^2 u}{\partial x^2} + 2u = 0 \tag{7}$$

Subject to the initial condition

$$u(x,0) = e^x \tag{8}$$

To solve this system (7) to (8) by HAM, first we choose initial approximation

$$u_0(x,t) = e^x$$

And the linear operator

$$L(\phi_1(x,t;q)) = \frac{\partial \phi_1(x,t;q)}{\partial t}$$

With the property  $L(C) = 0$  where  $C$  is integral constant.

We define system of non-linear operator as

$$N(\phi_1(x,t;q)) = \frac{\partial \phi_1(x,t;q)}{\partial t} - \frac{\partial \phi_1(x,t;q)}{\partial x} \frac{\partial^2 \phi_1(x,t;q)}{\partial x^2} + 2\phi_1(x,t;q) \tag{9}$$

Using the above definition, we construct the zeroth-order deformation equations

$$(1-q)[\phi_1(x,t;q) - u(x,t)] = qhN(\phi_1(x,t;q)) \tag{10}$$

Obviously, when  $q = 0$  and  $q = 1$  we get

$$\phi_1(x, t; 0) = u_0(x, t) \text{ and } \phi_1(x, t; 1) = u(x, t) \tag{11}$$

As  $q$  increase 0 to 1,  $\phi_1$  varies from  $u_0(x, t)$  to  $u(x, t)$  Expanding  $\phi_1(x, t; q)$  in Taylor series with respect to  $q$ ,

$$\phi_1(x, t; q) = u_0(x, t) + \sum_{m=1}^{\infty} u_m(x, t) \cdot q^m \tag{22}$$

Where

$$u_m(x, t) = \left[ \frac{1}{m!} \frac{\partial^m \phi_1(x, t; q)}{\partial q^m} \right]_{q=0} \tag{23}$$

If the auxiliary linear operator, initial guesses, the auxiliary parameter  $h$  and auxiliary functions are properly chosen than the series equation (23) converges at  $q = 1$ .

$$\phi_1(x, t; 1) = u_0(x, t) + \sum_{m=1}^{\infty} u_m(x, t)$$

$$\text{i.e. } u(x, t) = u_0(x, t) + \sum_{m=1}^{\infty} u_m(x, t)$$

This must be one of solutions of the original non linear equations as proved by Liao Define the vectors

$$\overline{u}_n = (u_0(x, t), u_1(x, t), u_2(x, t), \dots, u_{n-1}(x, t)) \tag{24}$$

We have the so-called  $m^{\text{th}}$  order deformation equations

$$L[u_m(x, t) - \chi_m u_{m-1}(x, t)] = h R_m(\overline{u}_{m-1}) \tag{25}$$

Where

$$R_{1,m}(\overline{u}_{m-1}) = \left[ \frac{1}{(m-1)!} \frac{\partial^{m-1} \phi_1(x, t; q)}{\partial q^{m-1}} \right]_{q=0} \tag{26}$$

$$\text{i.e. } R_{1,m}(\overline{u}_{1,m-1}) = \frac{\partial u_{m-1}}{\partial t} - \frac{\partial u_{m-1}}{\partial x} \frac{\partial^2 u_{m-1}}{\partial x^2} + 2u_{m-1} \tag{27}$$

$$u_m(x, t) = \chi_m u_{m-1}(x, t) + h \int_0^t R_m(\overline{u}_{m-1}) dt + c \tag{28}$$

Now we will calculate

$$u_1(x, t) = \chi_m u_0(x, t) + h \int_0^t R_1(\overline{u}_0) dt \tag{29}$$

Where

$$R_1(\overline{u}_0) = 2e^x - 2e^{2x}$$

So

$$u_1(x, t) = 2ht [e^x - e^{2x}]$$

Now, the  $N^{\text{th}}$  order approximation can be expressed by

$$S_N(x, t) = u_0(x, t) + \sum_{m=1}^{N-1} u_m(x, t) \tag{30}$$

As  $N \rightarrow \infty$  we get  $S_N(x, t) \rightarrow u(x, t)$  with some appropriate assumption of  $h$

### III. NON HOMOGENEOUS NON LINEAR PARTIAL DIFFERENTIAL EQUATION

Consider non homogeneous non linear differential equation

$$\frac{\partial^2 u}{\partial t^2} + \frac{\partial u}{\partial x} \frac{\partial^2 u}{\partial x^2} + 2u + 3 = 0 \tag{45}$$

Subject to the initial condition

$$u(x, 0) = e^x \tag{46}$$

To solve this system (45) to (46) by HAM, first we choose initial approximation

$$u_0(x, t) = e^x$$

And the linear operator

$$L(\phi_1(x, t; q)) = \frac{\partial^2 \phi_1(x, t; q)}{\partial t^2}$$

With the property  $L(C) = 0$  where  $C$  is integral constant.

We define system of non-linear operator as

$$N(\phi_1(x, t; q)) = \frac{\partial \phi_1(x, t; q)}{\partial t^2} + \frac{\partial \phi_1(x, t; q)}{\partial x} \frac{\partial^2 \phi_1(x, t; q)}{\partial x^2} + 2\phi_1(x, t; q) + 3 \tag{47}$$

Using the above definition, we construct the zeroth-order deformation equations

$$(1 - q)[\phi_1(x, t; q) - u(x, t)] = qhN(\phi_1(x, t; q)) \tag{48}$$

Obviously, when  $q = 0$  and  $q = 1$  we get

$$\phi_1(x, t; q) = u_0(x, t) \text{ and } \phi_1(x, t; 1) = u(x, t) \tag{49}$$

As  $q$  increase 0 to 1,  $\phi_1$  varies from  $u_0(x, t)$  to  $u(x, t)$  Expanding  $\phi_1(x, t; q)$  in Taylor series with respect to  $q$ ,

$$\phi_1(x, t; q) = u_0(x, t) + \sum_{m=1}^{\infty} u_m(x, t) \cdot q^m \tag{50}$$

Where

$$u_m(x, t) = \left[ \frac{1}{m!} \frac{\partial^m \phi_1(x, t; q)}{\partial q^m} \right]_{q=0} \tag{51}$$

If the auxiliary linear operator, initial guesses, the auxiliary parameter  $h$  and auxiliary functions are properly chosen than the series equation (51) converges at  $q = 1$ .

$$\phi_1(x, t; 1) = u_0(x, t) + \sum_{m=1}^{\infty} u_m(x, t)$$

$$\text{i.e. } u(x, t) = u_0(x, t) + \sum_{m=1}^{\infty} u_m(x, t)$$

This must be one of solutions of the original non linear equations as proved by Liao Define the vectors

$$\overline{u}_n = (u_0(x, t), u_1(x, t), u_2(x, t), \dots, u_n(x, t)) \tag{52}$$

We have the so-called  $m^{\text{th}}$  order deformation equations

$$L[u_m(x, t) - \chi_m u_m(x, t)] = hR_m(\overline{u}_{m-1}) \tag{53}$$

Where

$$R_m(\overline{u_{m-1}}) = \left[ \frac{1}{(m-1)!} \frac{\partial^{m-1} \phi_1(x, t; q)}{\partial q^{m-1}} \right]_{q=0} \tag{54}$$

$$\text{i.e. } R_m(\overline{u_{m-1}}) = \frac{\partial^2 u_{m-1}}{\partial t^2} + \frac{\partial u_{m-1}}{\partial x} \frac{\partial^2 u_{m-1}}{\partial x^2} + 2u_{m-1} + 3 \tag{55}$$

$$u_m(x, t) = \chi_m u_{m-1}(x, t) + h \int_0^t \int_0^t R_m(\overline{u_{m-1}}) dt + c \tag{56}$$

Now we will calculate

$$u_1(x, t) = \chi_m u_0(x, t) + h \int_0^t \int_0^t R_1(\overline{u_0}) dt + c \tag{57}$$

Where

$$R_1(\overline{u_0}) = e^{2x} + 2e^x + 3$$

So

$$u_1(x, t) = \frac{ht^2}{2} (e^{2x} + 2e^x + 3)$$

Now the  $N^{th}$  order approximation can be expressed by

$$S_N(x, t) = u_0(x, t) + \sum_{m=1}^{N-1} u_m(x, t) \tag{58}$$

As  $N \rightarrow \infty$  we get  $S_N(x, t) \rightarrow u(x, t)$  with some appropriate assumption of  $h$

#### IV. CONCLUSION

Various types of homogeneous and non homogeneous non linear partial differential equation can be solved by using homotopy analysis method. Using freedom of choosing parameter  $h$ , numerical result can be derived.

#### REFERENCES

- [1] Abbasbandy S., "The application of homotopy analysis method to non linear equations arising in heat transfer", Phys. Lett. A 360 (2006) 109113
- [2] Batiha B., Noorani M.S.M., Hashim I., Numerical simulations of systems of PDEs by variational iteration method, Phys. Lett. A, 372 (2008) 822829
- [3] Bataineh A.Sami, Noorani M.S.M., Hashim I., Modified homotopy analysis method for solving systems of second-order BVPs, Commun. Nonlinear Sci. Numer. Simul., in press (doi:10.1016/j.cnsns.2007.09.012).
- [4] S.J Liao., "On the homotopy analysis method for non linear problems", Applied Mathematics and Computation, vol.147, no. 2, pp.499513,2004.
- [5] S.J. Liao and Y. Tan, "A general approach to obtain series solutions of non linear diferential equations", Studies in Applied Mathematics, vol.119, no.4, pp.297354,2007
- [6] S.J. Liao., Beyond Perturbation: Introduction to the Homotopy Analysis Method, Chapman
- [7] S.J. Liao, Comparison between the homotopy analysis method and homotopy perturbation method, Appl. Math. Comput. 169 (2005) 11861194
- [8] Saha R.S., A numerical solution of the coupled sine-Gordon equation using the modified decomposition method, Appl. Math. Comput. 175 (2006) 10461054.
- [9] Wazwaz A.M., The variational iteration method for solving linear and non linear systems of PDEs, Comput. Math. Appl. 54 (2007) 895902.
- [10] Z. Wang ,I. Zou ,H. Zhang "Applying homotopy analysis method for solving diferential-diference equation". Phys Lett A 2007;371:72-82



10.22214/IJRASET



45.98



IMPACT FACTOR:  
7.129



IMPACT FACTOR:  
7.429



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24\*7 Support on Whatsapp)